

An intestinal *Candida albicans* model for monomicrobial and polymicrobial biofilms and the Bgl2 ligand

by Rini Purbowati

Submission date: 25-Dec-2022 11:13PM (UTC+0700)

Submission ID: 1986539844

File name: 2._logo-2_dan_naskah.pdf (2.42M)

Word count: 5061

Character count: 28201



An intestinal *Candida albicans* model for monomicrobial and polymicrobial biofilms and effects of hydrolases and the Bgl2 ligand

Masfufatun¹, Rini Purbowati², Nira A. Arum³, Mey S. Yasinta³, Sri Sumarsih¹ and Afaf Baktir⁴

10

1. Department of Biochemistry, Faculty of Medicine, University of Wijaya Kusuma Surabaya, Surabaya, Indonesia;
2. Department of Biomedicine and Biomolecular, Faculty of Medicine, University of Wijaya Kusuma Surabaya, Surabaya, Indonesia;
3. Department of Chemistry, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia;
4. Department of Chemistry, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia.

Corresponding author: Masfufatun Masfufatun, e-mail: masfufatun@uwks.ac.id

Co-authors: RP: rini.purbowati@uwks.ac.id, NAA: nira.ambar.arum-2016@fst.unair.ac.id,
 MSY: r42irli.yasinta-2016@fst.uniar.ac.id, SS: sri-s@fst.unair.ac.id, AB: afaf-b@fst.unair.ac.id

Received: 16-11-2021, **Accepted:** 14-03-2022, **Published online:** 29-04-2022

doi: [www.doi.org/10.14202/vetworld.2022.1134-1140](https://doi.org/10.14202/vetworld.2022.1134-1140) **How to cite this article:** Masfufatun M, Purbowati R, Arum NA, Yasinta MS, Sumarsih S, Baktir A (2022) An intestinal *Candida albicans* model for monomicrobial and polymicrobial biofilms and effects of hydrolases and the Bgl2 ligand, *Veterinary World*, 15(4): 1134-1140.

Abstract

20

Background and Aim: *Candida albicans* is the most prevalent human fungal pathogen. In biofilms, *C. albicans* becomes more resistant to antifungal agents because of the production of an extracellular matrix (ECM) that protects the yeast cells. This study aimed to determine the effects of hydrolase enzymes and the Bgl2 ligand on monomicrobial and polymicrobial biofilms.

Materials and Methods: Biofilm induction in rats was carried out using streptomycin (25 mg/kg) and gentamicin (7.5 mg/kg) administered orally once per day for 5 days. Rats were injected subcutaneously with cortisone acetate (225 mg/kg) as an immunosuppressant on day 5. In addition, rats were orally administered *C. albicans* for the single microbial model and a combination of *C. albicans* with *Escherichia coli* for the polymicrobial model. Following the biofilm production, the groups were treated with glucosamine (8.57 mg/kg body weight) and *Achatina fulica* hydrolases (1.5 μ g) orally for 2 weeks. The reduction of the biofilm was measured using confocal laser scanning microscopy (CLSM). Data were analyzed using a t-test, with a significance value of 95%.

Results: CLSM images revealed a strong association between *C. albicans* and *E. coli* in the polymicrobial biofilm. On the contrary, the combination treatment using glucosamine and *A. fulica* hydrolases reduced the ECM of the single microbial biofilm (53.58%). However, treatment effectiveness against the matrix (19.17%) was reduced in the polymicrobial model.

Conclusion: There is a strong association between *C. albicans* and *E. coli* in the formation of polymicrobial biofilms. The combination of glucosamine and the *A. fulica* enzyme can reduce the single microbial biofilm ECM; however, it is ineffective in the polymicrobial model.

Keywords: *Achatina fulica* hydrolases, Bgl2 ligand, *Candida albicans*, *Escherichia coli*, intestinal polymicrobial biofilm.

Introduction

The human body is a host for one billion microorganisms. *Candida albicans* is usually found in the normal human micro-ecosystem. *Candida* inhabits several body parts of humans, such as the oral cavity, gastrointestinal tract, skin, and vagina [1,2]. The virulence of *C. albicans* is related most commonly to the formation of biofilms. A biofilm is a structured microbial community covered in an extracellular matrix (ECM) and attached to a surface. The biofilm shows phenotypic characteristics different from planktonic or motile cells [3]. In the biofilm form, there is significantly greater resistance to antimicrobial agents [4-8]. The ECM is secreted by the sessile communities and protects them. It consists mainly of polysaccharides,

such as glucans, mannan, and chitin, and it acts as a barrier and hampers the ability of antifungal molecules to reach the cells. However, it is not the only cause of biofilm resistance [5,9,10].

As the presence of the ECM can inhibit antifungal action, it is essential to develop research to find herbal remedies that can destroy the matrix material. Enzymatic treatment may be a good strategy to improve antifungal performance. In our previous work, we successfully inhibited biofilm formation using the Bgl2p ligand, glucosamine, as an inhibitory molecule. Bgl2p is a glucosyltransferase encoded by the Bgl2 gene. This glucosyltransferase is responsible for forming covalent bonds between 1,3-glucan and other cell wall components and ECM components [11]. In this work, we combined the action of the glucosamine ligand and the hydrolase mixture produced by *Achatina fulica* to support antimicrobial performance for hydrolysis of the ECM in the monomicrobial and polymicrobial models of *C. albicans*.

Therefore, the purposes of this study were (1) to prove the ability of *C. albicans* to form a mixed or polymicrobial intestinal biofilm with *Escherichia coli*;

Copyright: Masfufatun, et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

7

(2) to evaluate the interactions between *C. albicans* and *E. coli* in biofilms from rat models; and (3) to investigate the performance of glucosamine (as the Bgl2 ligand) and the combination of *A. fulica* hydro-lases against single and polymicrobial biofilms.

Materials and Methods

Ethical approval

All animal experiments were carried out in Laboratory Animals, Faculty of Veterinary, Universitas Airlangga, Surabaya, Indonesia. Before the animal experiments were conducted, the ethical clearance (Reg. 271-KE) was obtained from the Animal Care and Use Committee, Faculty of Veterinary Medicine Universitas Airlangga.

Study period and location

This study was conducted from March to November 2021. The experiment using animals was managed in Animal Laboratory, Faculty of Veterinary Medicine, Universitas Airlangga. Meanwhile, the observation of confocal laser scanning microscope (CLSM) was carried out in Central Laboratory of Biological Sciences, Universitas Brawijaya.

Microbial strains, media, and growth conditions

The strains used in this study were *C. albicans*-ATCC 10231 and *E. coli*-ATCC 25992 purchased from the Balai Besar Laboratorium Kesehatan. Overnight *C. albicans* cultures were grown in yeast extract-peptone-dextrose (YPD) medium (1% yeast extract [BD Biosciences, USA], 2% peptone [Oxoid Ltd, UK], and 2% dextrose [Conda Prudisa, Spain]) at 30°C. Overnight *E. coli* cultures were grown in Luria-Bertani (LB) medium (1% tryptone [Himedia, India], 0.5% yeast extract, and 1% NaCl [Merck, Germany]) at 37°C.

Study design

The animals used for the *in vivo* experiments were male Wistar rats weighing about 200 g. All animals were acclimatized for 1 week and given a standard *ad libitum* diet. The rats were divided into five treatment groups, each consisting of five individuals: Group 1, negative control (without biofilm induction); Group 2, monomicrobial biofilm control; Group 3, biofilm treatment with enzyme and antifungal for monomicrobial film; Group 4, polymicrobial biofilm control; and Group 5, biofilm treatment with enzyme and antifungal for polymicrobial film.

In vivo monomicrobial induction of *C. albicans* biofilm

Biofilm induction in rat models was carried out according to Masfufatun *et al.* [12] methods, with modifications. The *C. albicans* stock culture was grown on Sabouraud dextrose agar (SDA; 1% peptone, 4% dextrose, and 1.5% agar) at room temperature (25°C). A single colony was inoculated in YPD broth, with shaking at 120 rpm (37°C) overnight. The inoculum was then orally administered to rats in the biofilm groups. During the biofilm induction periods, rats were given a minimum portion of food

with 2.5 mL of spider medium (1% nutrient broth, 1% D-mannitol [Himedia], and 0.2% monopotassium phosphate [Merck]) twice per day to induce biofilm formation further. Rat models from Groups 2 and 4 were sacrificed to observe the biofilm of *C. albicans* macroscopically on the mucous membrane of the cecum after biofilm induction, 35 days after the inoculation of *C. albicans*.

In vivo polymicrobial induction of *C. albicans-E. coli* biofilms

The induction of the polymicrobial model was carried out using methods similar to those for the *in vivo* monomicrobial model, with the addition of orally administered *E. coli* inoculum in LB medium along with *C. albicans*.

Enzymatic treatments with Bgl2 ligand and antimicrobial combination

After the biofilm induction phase, Groups 3 and 5 received the combination of the enzymatic Bgl2p ligand and antimicrobial treatment for 1 week. The combination of treatment methods is described below. The *A. fulica* enzyme mixture was harvested according to Baktir *et al.* [13]. The harvested enzyme mixture (1.5 mL) and glucosamine (8.57 mg/kg body weight [BW] and fluconazole, and 6 mg/kg BW; with the addition of tetracycline for the polymicrobial model) were orally administered to the model rats of Groups 3 and 5. The rats were then sacrificed after the treatment period to macroscopically observe the *C. albicans* biofilm on the mucous membrane of the cecum after biofilm induction.

Macroscopic identification of biofilm

The intestinal model biofilm was identified by macroscopic observations of the mucous membrane of the cecum after dissecting and photographing it with a camera (Canon EOS m3).

CLSM

The sample preparation was prepared using the following protocol. Clean and sliced cecum samples were fixed in 10% formalin buffer, submerged in paraffin, and cut into sections of 5-μm thickness. The cecum tissues were deparaffinized with xylol (2 times) for 10 min each, hydrated with ethanol (absolute ethanol, 93% and 70%, consecutively), washed with phosphate-buffered saline (PBS) for 5 min, blocked with 2% bovine serum albumin (BSA) in PBS at 25°C for 1 h, and rewashed with PBS for 8 min. A 100-μg/mL aliquot of Concanavalin A (con A; bioWORLD, USA) was dropped onto the sample, which was then incubated at 25°C for 1 h. Finally, the samples were rewash with PBS, reblocked with 2% BSA in PBS, and observed under a confocal microscope (Olympus FV1000).

Statistical analysis

All data were subjected to an independent t-test to assess significant differences between two groups. Data normality was calculated using the

Shapiro-Wilk test (for samples <50). Statistical analyses were performed using Statistical Package for the Social Sciences 16.0 software (IBM Corp., NY, USA). A value of $p<0.05$ was considered to indicate a significant difference between the control and treatment groups. All values are expressed as the mean \pm standard deviation.

Results

Macroscopic and microscopic observations of monomicrobial and polymicrobial *C. albicans* biofilms in an *in vivo* rat model

The *in vivo* monomicrobial *C. albicans* biofilms in the rats of Group 2 appeared as white lesions in the mucous membrane of the cecum under macroscopic evaluation Figure-1a. The normal rat model (Group 1) showed a normal mucous membrane of the cecum without white lesions Figure-1b. Microscopic images of *C. albicans* and the matrix of monomicrobial biofilm model of the biofilm control group (Group 1) (c) and control group (Group 2) (d) are shown. The green and red intensities in the biofilm group (Figure-1c) were stronger than those in the control group (Figure-1d). Green fluorescence indicates the presence of *C. albicans* biofilm matrix, and red luminescence indicates the presence of *C. albicans* cells. Yellow fluorescence indicates the presence




Figure-1: Macroscopic and microscopic image of monomicrobial *Candida albicans* biofilm in cecum mucous membrane. Macroscopic: (a) Biofilm of the Group 2 rat model and a (b) Normal rat model (Group 1) (b). Biofilm shown in the red circle, appeared as white lesions. Microscopic: (c) Biofilm of the Group 2 rat model and (d) Normal rat model (Group 1). The green colored image displayed a matrix, colored using Concavaline A. The red colored image displayed *C. albicans* cells, colored using polyclonal antibody anti-*Candida* conjugated to Tetramethyl Rhodamine Isothiocyanate. Yellow-orange colored displayed the superimposed image of *C. albicans* and matrix confocal image.

of *C. albicans* cells in the biofilm covered with the resulting ECM.

The *in vivo* polymicrobial biofilm model of Group 4 appeared as white lesions located in the mucous membrane of the cecum, as presented in Figure-2b. The normal rat model showed a normal outer mucous membrane of the cecum without white lesions, as presented in Figure-2a. There was a considerable difference in the cecum size between the normal group and the Group 4 models. Yellow fluorescence in the biofilm group (Figure-2d) is the result of overlapping red (*E. coli*) and green (*C. albicans*) images, showing that *C. albicans* cells interacted with bacterial cells to form polymicrobial biofilms. In the control group (Figure-2c), a portion was red with moderate intensity, proving that there were still many bacteria in the digestive tract.

CLSM analysis of monomicrobial and polymicrobial *C. albicans* biofilms in an *in vivo* rat model before and after treatment

Confocal imaging was used to assess the association between *C. albicans* and *E. coli* from the same polymicrobial model sample with different fluorescences. The confocal images of *C. albicans* and *E. coli* showed an association between these two species, as *C. albicans* mainly appeared in the same location as *E. coli*, as seen from the yellow-orange superimposed image (Figure-3a). The treatment resulted in a




Figure-2: Macroscopic and microscopic image of polymicrobial *Candida albicans* biofilm in cecum mucous membrane. Macroscopic: (a) Normal rat model (Group 1) and (b) Polymicrobial biofilm of the Group 4 rat model. Biofilm shown in the red circle, appeared as white lesions. Microscopic: (c) Normal rat model (Group 1) and (d) polymicrobial biofilm of the Group 4 rat model. The green colored image displayed *C. albicans* cells, colored using polyclonal antibody anti-*Candida*-fluorescein isothiocyanate. The red colored image displayed *Escherichia coli* cells, colored using SYTO-59. Yellow-orange colored displayed the superimposed image of *C. albicans* and *E. coli* confocal image.

decrease in fluorescence intensity, as evidenced by the dimmer appearance in the confocal image of the treatment group or Group 2 (Figure-3b) and the quantitative data of the mean intensity that showed a decrease in the treatment group (Figure-4). The decreased intensity was proportional to the decreased number of cells, as the fluorescence appeared due to the specific binding of the dye with parts of the cells.

The effect of *C. albicans* cells and matrix treatment in the polymicrobial model was also observed with confocal imaging. The treatment decreased *C. albicans* cells and matrix, demonstrated by the lower intensities of the confocal image fluorescence (Figure-5b), than that of the non-treatment group (Figure-5a). The decrease in intensity was proportional to the decrease in the ECM (Figure-4).

For the monomicrobial model, the results agreed with the polymicrobial biofilm model. A dimmer confocal image (Figure-6) demonstrated that the *C. albicans* cells and matrix had reduced intensities in the treatment group. Quantitatively, the monomicrobial biofilm had a smaller number of cells and matrix (indicated by the lower intensities) than the polymicrobial biofilm of the treatment group (Figure-7).

Discussion

The biofilm lifestyle protects cells that reside within the thick layer of the ECM. The biofilm allows cells to develop up to 1000 times more resistance to antimicrobials than planktonic cells, leaving biofilm eradication a challenging task [14,15]. For *C. albicans*, the biofilms grown on surfaces show significantly increased minimum inhibitory concentrations to antifungals, including fluconazole, compared to planktonic cells [15,16].

The biofilm of *C. albicans* consists of blastospore-type cells as the basal layer and a superficial layer of ECM and hyphal-type cells. The ECM comprises carbohydrates (glucose, N-acetylglucosamine, mannose, and rhamnose), proteins, phosphorus, uronic acid, and hexosamine [9,17]. One of the carbohydrate components is β -1,3-glucan, a glucose polymer,^[16] thought to be the main ECM component in biofilms. β -1,3-Glucans are synthesized by glucan synthase, a membrane-bound protein, using uridine diphosphate glucose as a substrate and secrete the molecules to the ECM. This component in *C. albicans* biofilm cells may contribute to resistance to antifungal drugs. Some studies have reported that β -1,3-glucans can be found in the supernatant around the biofilm and in the matrix. The level of this substance increased to a high concentration during the biofilm formation phase.^[19] The other evidence is that enzymatic treatment of β -1,3-glucans using β -1,3-glucanase at high concentrations eliminates biofilms but does not significantly affect planktonic cells. Enzymatic treatment using β -1,3-glucanase at low concentrations poorly disrupts biofilms; however, in combination with fluconazole, it improves the performance of the antifungal [18-20]. A previous study by Nett *et al.* [20] also suggested that β -1,3-glucans bind to fluconazole in biofilms, decreasing the antifungal efficacy to control biofilm-associated cells. Therefore, disrupting the β -1,3-glucan component using β -1,3-glucanase is a good strategy to support antifungal effectiveness.

The roles of several genes in the biofilm formation process have been investigated. The glucan transferases Phr1p and Bgl2p, encoded by *PHR1* and *BGL2*, respectively, and exoglucanase, encoded by




Figure-3: Confocal laser scanning microscopy image of *Candida albicans* and *Escherichia coli* cells. Confocal image of *C. albicans*-*E. coli* polymicrobial biofilm model. (a) Non-treatment polymicrobial biofilm group, (b) Treatment polymicrobial biofilm group. The green colored image displayed *C. albicans* cells, colored using polyclonal antibody anti-*Candida*-fluorescein isothiocyanate. The red colored image displayed *E. coli* cells, colored using SYTO-59. Yellow-orange colored displayed the superimposed image of *C. albicans* and *E. coli* confocal image.

XOG1, are predicted to exist in the ECM and have the roles of delivering and accumulating β -1,3-glucans in the ECM. Mutant strains lacking these genes show more susceptibility to fluconazole [21]. Therefore, the competitive inhibition of enzymes responsible for biofilm construction, such as Bgl2p, can be an effective way to promote the antimicrobial potential for killing cells directly.

A. fulica is a natural source of several hydrolase enzymes. It produces the enzyme mixture through the digestive gland to help digest food. However, several nondigestible materials are digested by the microbial enzymes produced in their gastrointestinal tract. The enzyme mixtures of the digestive tract of this species are carbohydrateases, such as mannosidase, glucosidase, chitinase, β -glucanases, proteinase, and lipase [22-25].




Figure-4: Quantitative data from confocal laser scanning microscopy assessment in polymicrobial biofilm model. The mean intensities of *Escherichia coli*, *Candida albicans*, and matrix were shown in the data below the chart. All of the *E. coli*, *C. albicans*, and matrix consecutively shown the decreased intensities in the treatment group by 31.51%, 39.03%, and 19.17%. **indicates the significant differences ($p < 0.05$)

The utilization of *A. fulica* might reduce the cost of enzyme production, as this species is fairly abundant, especially in a tropical and humid area like Indonesia. The enzymes from *A. fulica* can successfully lyse the fungal cell wall of *Candida* spp. [26]. *Candida* spp. [11] walls have components similar to the ECM, with β -1,3-glucans and β -1,6-glucan being the major carbohydrate components (50%-60%). Therefore, the enzyme mixtures of *A. fulica* have great potential as an affordable and effective antibiofilm treatment, as supported by Nett *et al.* [20], wherein treatment with β -1,3-glucanase was highly effective in eradicating *C. albicans* biofilms.

Therefore, we hypothesized that the combination of treatment using *A. fulica* mixture hydrolases and the *Bgl2* ligand, the glucose substrate look-alike, glucosamine (Figure-8), could work synergistically to enhance the eradication of the ECM and increase the killing of pathogenic cells that reside within the ECM. The matrix was decreased through hydrolysis of its complex polymer components, mainly β -1,3-glucan, to oligomers or monomers. The ligand, glucosamine, binds to the specific binding site of glucan transferase, replacing its original substrate, glucose, and inhibits β -1,3-glucan delivery and the accumulation of β -1,3-glucan in the matrix, thus further disrupting biofilm formation. However, in this study, we did not compare the performance of this potential antibiofilm as a combination and as separate components. We compared the action of this combined treatment with the results of the *in vivo* application of the monomicrobial and polymicrobial biofilm model.

Biofilms consisting of single microbial species have been extensively studied; however, more recent investigations have found that polymicrobial biofilms




Figure-5: Confocal laser scanning microscopy image of *Candida albicans* cells and matrix of the polymicrobial biofilm model. (a) Non-treatment polymicrobial biofilm group. (b) Treatment polymicrobial biofilm group. The green colored image displayed matrix, colored using Concanavalin A. The red colored image displayed *C. albicans* cells, colored using polyclonal antibody anti-*Candida* conjugated to TRITC. Yellow-orange colored displayed the superimposed image of *C. albicans* and matrix confocal image.




Figure-6: Confocal laser scanning microscopy image of *Candida albicans* cells and matrix of the monomicrobial biofilm model. (a) Non-treatment monomicrobial biofilm group. (b) Treatment monomicrobial biofilm group. The green colored image displayed matrix, colored using Concanavalin A. The red colored image displayed *C. albicans* cells, colored using polyclonal antibody anti-*Candida* conjugated to TRITC. Yellow-orange colored displayed the superimposed image of *C. albicans* and matrix confocal image.




Figure-7: Quantitative data from confocal laser scanning microscopy (CLSM) assessment in polymicrobial biofilm model. Quantitative data from CLSM assessment in monomicrobial biofilm model. The mean intensities of *Candida albicans* and matrix were shown in the data below the chart. Both of the *Candida albicans* and matrix consecutively shown the decreased intensities in the treatment group by 49.83% and 53.58%. **Indicates the significant differences ($p < 0.05$).




Figure-8: The structure of glucosamine.

are the dominant form in nature. This study investigated the interaction between bacteria and fungi in a polymicrobial biofilm model composed of *E. coli* and *C. albicans*. Both species are predominant pathogens that can grow as biofilms in medical devices [2]. This study demonstrated that ECM disruption using a

ligand–enzyme combination treatment in the monomicrobial biofilm model was more effective than the treatment in the polymicrobial biofilm model. The ECM was decreased significantly by 53.58% in the monomicrobial model. However, in the polymicrobial model, the treatment decreased the matrix intensity by 19.17%; this decrease was also significant (Figures-4 and 7). The difference in the ligand–enzyme performance against the matrix material might be due to the strong and synergistic association between *E. coli* and *C. albicans*. A previous study stated that *E. coli* might facilitate the attachment of *C. albicans* to the host surface, resulting in stronger binding between the cells and the host [27,28].

Conclusion

Treatment targeting matrix hydrolysis and the inhibition of Bgl2 using a combination of antimicrobials successfully reduced the biofilm matrix of *E. coli* and *C. albicans*. In this study, we have found a mixture of glucosamine and Bgl2 enzyme as herbal remedies to destroy the matrix material of *C. albicans* biofilm in the monomicrobial model. However, it showed limited activity towards polymicrobial biofilm of *C. albicans* and *E. coli*. In the next study we will explore antibiofilm especially for *E. coli*.

Authors' Contributions

MM: Conceived and designed the study, performed the experimental work, and reviewed the manuscript. NAA and MSY: Collected microorganism isolates and reviewed the manuscript. AB and RP: Data interpretation, statistical analysis, and reviewed the manuscript. SS: Conducted the literature review and drafted the manuscript. All authors have read and approved the final manuscript.

Acknowledgments

The authors would like to thank Directorate of Research and Community Service, Directorate General Strengthening for Research and Development of the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for the research funding.

Competing Interests

The authors declare that they have no competing interests.

Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

References

1. 11riotti, M.M. and Noverr, M.C. (2011) Importance of *Candida*-bacterial polymicrobial biofilms in disease. *Trends Microbiol.*, 19(11): 557-563.
2. Samaranayake, Y.H., Bandara, H.M.H., Cheung, B.P.K., J.Y.Y., Yeung, S.K.W. and Samaranayake, L.P. (2014) Enteric gram-negative bacilli suppress *Candida* biofilms on poly urinary catheters. *APMIS*, 122(1): 47-58.
3. 74ati, M. and Nobile, C.J. (2016) *Candida albicans* biofilms: Development, regulation, and molecular mechanisms. *Microbes Infect.*, 18(5): 310-321.
4. May, F.L., Wilson, D. and Huber, B. (2013) *Candida albicans* pathogenicity mechanisms. *Virulence*, 4(2): 119-128.
5. Nobile, C.J. and Johnson, A.D. (2015) *Candida albicans* biofilms and human disease. *Annu. Rev. Microbiol.*, 69: 71-92.
6. Fu, 63Ying, Y., Wei, B., Wang, L., Xu, S., Qin, P., Wei, L., Jiang, L. (2017) Epidemiology of *Candida albicans* and *C. albicans* of neonatal candidemia at a tertiary care hospital in Western China. *BMC Infect. Dis.*, 17(1): 329.
7. Bitew, A. and Abebew, Y. (2018) Vulvovaginal candidiasis: Species distribution of *Candida* and their antifungal susceptibility pattern. *BMC Womens Health*, 18(1): 94.
8. 72H.T., Lee, T.Y., Chen, Y.C. and Charbonneau, C. (2017) Pharmacoeconomic analysis of antifungal therapy for primary treatment of invasive candidiasis caused by *Candida albicans* and non-albicans *Candida* species. *BMC Infect Dis.*, 18: 481.
9. 18Ree, C.G., Vila, T., Romo, J.A., Montelongo-Jauregui, D., 29J.G., Ramasubramanian, A. and Lopez-Ribot, J.L. (2017) The *Candida albicans* biofilm matrix: Composition, structure and function. *J. Fungi*, 3(1): 14.
10. 32nowski, R., Westler, W.M., de Lacmbouh, G.A., Marita, J.M., Bothe, J.R., Bernhardt, J., Sahraoui, A.L.H., Fontaine, J., Sanchez, H., Hatfield, R.D., Ntambi, J.M., 33tt, J.E., Mitchell, A.P. and Andes, D.R. (2014) Novel entries in a fungal biofilm matrix encyclopedia. *mBio*, 5(4): 34333-e01314.
11. Hartland, R.P., Emerson, G.W. and Sullivan, P.A. (1991) A secreted formula-glucan-branching enzyme from *Candida albicans*. *Proc. R. Soc. B Biol. Sci.*, 246(1316): 155-160.
12. Masufatun, M., Raharjo, L.H., Wiradinata, H., Tania, P.O.A., Ni'matuzahroh, N. and Bakir, A. (2021) New phenomena for clinicians, model of *Candida albicans* mobilization before and after biofilm formation in the intestinal mucosa of Wistar rats (*Rattus norvegicus*). *Int. J. One Health*, 7(2): 165-170.
13. Baktir, A., Masufatun and Hadi, S. (2015) Preclinic study of Bgl2 ligand as component of mix antibiofilm toward *Candida albicans* on mucous intestinal biofilm pre-induced *Rattus norvegicus*. *Pharm. Chem.*, 7(3): 113-118.
14. 51erry, L., Rajendran, R., Lappin, D.F., Borghi, E., 8rdoni, F., Falleni, M., Tosi, D., Smith, K., Williams, C., Jones, B., Nile, C.J. and Ramage, G. (2014) Biofilms formed by *Candida albicans* bloodstream isolates display phenotypic and transcriptional heterogeneity that are associated with resistance and pathogenicity. *BMC Microbiol.*, 14: 182.
15. 52ra, H.H., Urbina, D.G.R., Pierce, C., Ribot, J.L.L., Jiménez, M.J.A. and Yacaman, M.J. (2015) Effect of silver nanoparticles on *Candida albicans* biofilms: An ultrastructural study. *J. Nanobiotechnol.*, 13: 91.
16. Chandra, J. and Mukherjee, P.K. (2015) *Candida* biofilms: Development, architecture, and resistance. *Microbiol. Mol. Biol. Rev.*, 81(3): 1-24.
17. Lal, P., Sharma, D., Pruthi, P. and Pruthi, V. (2010) Exopolysaccharide analysis of biofilm-forming *Candida* strains. *J. Appl. Microbiol.*, 109(1): 128-136.
18. Al-Fattani, M.A. and Douglas, L.J. (2006) Biofilm matrix *Candida albicans* and *Candida tropicalis*: Chemical composition and role in drug resistance. *J. Med. Microbiol.*, 57(9 Pt 8): 999-1008.
19. Nett, J.E., Cain, M.T., Crawford, K., Andes, D.R., Middleton, W.S. and Hospital, V. (2011) Optimizing a *Candida* biofilm microtiter plate model for measurement of antifungal susceptibility by tetrazolium salt assay. *J. Clin. Microbiol.*, 49(4): 1426-1433.
20. 43Li, J., Lincoln, L., Marchillo, K., Massey, R., Holoyda, K., 3off, B., VanHandel, M. and Andes, D. (2007) Putative role of beta-1,3 glucans in *Candida albicans* biofilm resistance. *Microb. Agents Chemother.*, 51(2): 510-520.
21. 16C, H.T., Nett, J.E., Zarnowski, R., Ross, K.M., 56chez, H., Cain, M.T., Hamaker, J., Mitchell, A.P. and 3ades, D.R. (2012) A *Candida* biofilm-induced pathway for matrix glucan delivery: Implications for drug resistance. *PLoS Pathog.*, 8(8): e1002848.
22. Karasov, W.H. and Douglas, A.E. (2013) Comparative digestive physiology William. *Physiol. Behav.*, 109(2): 23-783.
23. 23aroux, S. and Colas, B. (1994) Digestive juice of *Achatina achatina* as a potential source of transglycosylating enzymes. *Int. J. Biochem.*, 26(2): 247-254.
24. 19ida, I., Shimohigashi, Y., Kihara, H. and Ohno, M. (1996) Purification and characterization of a cellulase from the giant snail *Achatina fulica*. *Biosci. Biotechnol. Biochem.*, 60(1): 122-124.
25. 20gbua, S.I.O., Anosike, E.O. and Ugochukwu, E.N. (1978) Partial Purification and Some Properties of Arylsulphatases from the Gut of the Giant African Snail, *Achatina achatina*, No. 5915. p169-173.
26. 65onye, O.U. and Okerentugba, P.O. (2001) Optimum conditions for yeast protoplast release and regeneration in *Saccharomyces cerevisiae* and *Candida tropicalis* using gut enzymes of the giant African snail *Achatina achatina*. *Lett. Appl. Microbiol.*, 32(3): 190-193.
27. Makrilia, H.C. and MacFarlane, T.W. (1983) An investigation of the factors involved in increased adherence of *C. albicans* to epithelial cells mediated by *E. coli*. *Antimicrob. Agents Chemother.*, 38(153-154): 177-185.
28. De Brucker, K., Tan, Y., Vints, K., Cremer, K., De, Braem, A., 38straeten, N., Michiels, J., Vleugels, J., Cammue, B.P.A., 22Thevissen, K. (2015) Fungal beta-1,3-glucan increases 22xacin tolerance of *Escherichia coli* in a polymicrobial *E. coli/Candida albicans* biofilm. *Antimicrob. Agents Chemother.*, 59(6): 3052-3058.

An intestinal *Candida albicans* model for monomicrobial and polymicrobial biofilms and effects of hydrolases and the Bgl2 ligand

ORIGINALITY REPORT



PRIMARY SOURCES

- | Rank | Source Description | Percentage |
|------|---|------------|
| 1 | Submitted to Columbia Union College Student Paper | 1% |
| 2 | Aida Pitarch, César Nombela, Concha Gil. "Chapter 20 Collection of Proteins Secreted from Yeast Protoplasts in Active Cell Wall Regeneration", Springer Science and Business Media LLC, 2008 Publication | 1% |
| 3 | Reichhardt, Courtney, David A Stevens, and Lynette Cegelski. "Fungal biofilm composition and opportunities in drug discovery", Future Medicinal Chemistry, 2016. Publication | 1% |
| 4 | Submitted to Charles Sturt University Student Paper | 1% |
| 5 | Alejandra Martínez, Constanza Apip, Manuel.F Meléndrez, Mariana Domínguez et al. " Dual antifungal activity against of copper metallic nanostructures and hierarchical copper oxide | 1% |

marigold - like nanostructures grown in the culture medium ", Journal of Applied Microbiology, 2020

Publication

- 6 Huidi Wang, Jie Li, Guangyan Wu, Feng Zhang, Jia Yin, Yan He. "The effect of intrinsic factors and mechanisms in shaping human gut microbiota", Medicine in Microecology, 2022 1 %
Publication
- 7 Submitted to Truckee Meadows Community College 1 %
Student Paper
- 8 Submitted to Heriot-Watt University 1 %
Student Paper
- 9 www.jove.com 1 %
Internet Source
- 10 Submitted to Universitas Mataram <1 %
Student Paper
- 11 Vital Vialás, Palani Perumal, Dolores Gutierrez, Pilar Ximénez-Embún, César Nombela, Concha Gil, W. LaJean Chaffin. "Cell surface shaving of Candida albicans biofilms, hyphae, and yeast form cells", PROTEOMICS, 2012 <1 %
Publication
- 12 Tseng, Tzu-Ling, Wei-Chung Lai, Tai-Lin Lee, Wan Hua Hsu, Yu Wen Sun, Wan Chen Li, Chun-Wen Cheng, and Jia-Ching Shieh. "A role <1 %

of *Candida albicans* CDC4 in the negative regulation of biofilm formation", Canadian Journal of Microbiology, 2014.

Publication

-
- 13 Submitted to Barry University <1 %
Student Paper
-
- 14 medicopublication.com <1 %
Internet Source
-
- 15 www.turkarchotolaryngol.net <1 %
Internet Source
-
- 16 K. Hirota, H. Yumoto, B. Sapaar, T. Matsuo, T. Ichikawa, Y. Miyake. " Pathogenic factors in biofilm-related infectious diseases ", Journal of Applied Microbiology, 2017 <1 %
Publication
-
- 17 Samaranayake, Y. H., H. M. H. N. Bandara, B. P. K. Cheung, J. Y. Y. Yau, S. K. W. Yeung, and L. P. Samaranayake. "Enteric gram-negative bacilli suppress Candida biofilms on Foley urinary catheters", Apmis, 2013. <1 %
Publication
-
- 18 pureadmin.qub.ac.uk <1 %
Internet Source
-
- 19 Youssef Bacila Sade, Camila Silva Gonçalves, Sandra Mara Naressi Scapin, Guilherme Luiz Pinheiro et al. "Identification and Characterization of a Glycoside Hydrolase <1 %

Family 9 Member from the Digestive Gland of
the Snail *Achatina fulica*", BioEnergy Research,
2021

Publication

-
- 20 ec.asm.org <1 %
Internet Source
- 21 Bernardo Ramírez-Zavala, Austin Mottola, Julia Haubenreißer, Sabrina Schneider et al. "The Snf1-activating kinase Sak1 is a key regulator of metabolic adaptation and fitness of ", *Molecular Microbiology*, 2017
Publication
- 22 Giulia Orazi, George A. O'Toole. ""It Takes a Village": Mechanisms Underlying Antimicrobial Recalcitrance of Polymicrobial Biofilms", *Journal of Bacteriology*, 2019
Publication
- 23 Guionie, O.. "Identification and primary characterization of specific proteases in the digestive juice of Archachatina ventricosa", *Comparative Biochemistry and Physiology, Part B*, 200307
Publication
- 24 Marcus Gentsch, Margret Kuschel, Susan Schlegel, Gerold Barth. " Mutations at different sites in members of the Gpr1/Fun34/YaaH protein family cause <1 %

hypersensitivity to acetic acid in as well as in
", FEMS Yeast Research, 2007

Publication

- 25 Nívea P. de Sá, Ana P. Pôssa, Pilar Perez, Jaqueline M.S. Ferreira et al. "Antifungal Activity Directed Toward the Cell Wall by 2-Cyclohexylidenhydrazo- 4-Phenyl-Thiazole Against Candida albicans", Infectious Disorders - Drug Targets, 2019
Publication
- 26 Submitted to University of Modena and Reggio Emilia <1 %
Student Paper
- 27 bsdwebstorage.blob.core.windows.net <1 %
Internet Source
- 28 vtechworks.lib.vt.edu <1 %
Internet Source
- 29 Lucia Černáková, Catilin Light, Bahare Salehi, Cristian Rogel-Castillo et al. "Chapter 400 Novel Therapies for Biofilm-Based Candida spp. Infections", Springer Science and Business Media LLC, 2019 <1 %
Publication
- 30 ijeais.org <1 %
Internet Source
- 31 www.biorxiv.org <1 %
Internet Source

| | | |
|----|--|------|
| 32 | Submitted to RMIT University Student Paper | <1 % |
| 33 | Submitted to University of Surrey Student Paper | <1 % |
| 34 | mafiadoc.com Internet Source | <1 % |
| 35 | repository.kulib.kyoto-u.ac.jp Internet Source | <1 % |
| 36 | researchoutput.ncku.edu.tw Internet Source | <1 % |
| 37 | Binkley, J., M. B. Arnaud, D. O. Inglis, M. S. Skrzypek, P. Shah, F. Wymore, G. Binkley, S. R. Miyasato, M. Simison, and G. Sherlock. "The Candida Genome Database: The new homology information page highlights protein similarity and phylogeny", Nucleic Acids Research, 2014. Publication | <1 % |
| 38 | Submitted to Trinity College Dublin Student Paper | <1 % |
| 39 | epdf.pub Internet Source | <1 % |
| 40 | onlinelibrary.wiley.com Internet Source | <1 % |
| 41 | pubs.acs.org Internet Source | |

<1 %

42 www.thieme-connect.com <1 %
Internet Source

43 Submitted to Anglia Ruskin University <1 %
Student Paper

44 JOS?? MA RUIZ-MORENO. "PHOTODYNAMIC THERAPY AND HIGH-DOSE INTRAVITREAL TRIAMCINOLONE TO TREAT EXUDATIVE AGE-RELATED MACULAR DEGENERATION", Retina, 07/2006 <1 %
Publication

45 John Wiley & Sons, Ltd.. "Current awareness on yeast", Yeast, 09/2007 <1 %
Publication

46 José Ruiz-Herrera, Lucila Ortiz-Castellanos. "Cell wall glucans of fungi. A review", The Cell Surface, 2019 <1 %
Publication

47 academic.hep.com.cn <1 %
Internet Source

48 dr.ntu.edu.sg <1 %
Internet Source

49 scholarworks.bwise.kr <1 %
Internet Source

- 50 www.abbs.info <1 %
Internet Source
-
- 51 www.springerprofessional.de <1 %
Internet Source
-
- 52 H. M. H. N. Bandara. "Chapter 4 Current and Emerging In Vitro and In Vivo Biofilm Models in Investigating Fungal-Bacterial Polymicrobial Communities", Springer Science and Business Media LLC, 2023 <1 %
Publication
-
- 53 M. M. Harriott, M. C. Noverr. "Ability of Candida albicans Mutants To Induce Staphylococcus aureus Vancomycin Resistance during Polymicrobial Biofilm Formation", Antimicrobial Agents and Chemotherapy, 2010 <1 %
Publication
-
- 54 Margarida Martins. "Presence of Extracellular DNA in the Candida albicans Biofilm Matrix and its Contribution to Biofilms", Mycopathologia, 12/13/2009 <1 %
Publication
-
- 55 Maryam Dadar, Ruchi Tiwari, Kumaragurubaran Karthik, Sandip Chakraborty, Youcef Shahali, Kuldeep Dhama. "Candida albicans - Biology, molecular characterization, pathogenicity, and advances" <1 %

in diagnosis and control – An update",
Microbial Pathogenesis, 2018

Publication

-
- 56 Submitted to University of Wisconsin,
Madison <1 %
Student Paper
-
- 57 mic.microbiologyresearch.org <1 %
Internet Source
-
- 58 uniba.sk <1 %
Internet Source
-
- 59 Gabrilska, Rebecca A, and Kendra P
Rumbaugh. "Biofilm models of polymicrobial
infection", Future Microbiology, 2015.
<1 %
Publication
-
- 60 H.-W. Jeng, A. R. Holmes, R. D. Cannon. "
Characterization of two surface mannoprotein
adhesins that bind immobilized saliva
components ", Medical Mycology, 2005
<1 %
Publication
-
- 61 J. Kang, Y. He, D. Hetzl, H. Q. Jiang, M. K. Jun,
M. S. Jun, M. Khng, N. Cirillo, M. J. McCullough.
"A Candid Assessment of the Link between
Oral *Candida* Containing Biofilms and
Oral Cancer", Advances in Microbiology, 2016
<1 %
Publication
-
- 62 L. Julia Douglas. " Adhesion of Species to
Epithelial Surfaces ", CRC Critical Reviews in
<1 %

Microbiology, 2009

Publication

| | | |
|----|--|------|
| 63 | Submitted to University of Aberdeen Student Paper | <1 % |
| 64 | encyclopedia.pub Internet Source | <1 % |
| 65 | journal.unnes.ac.id Internet Source | <1 % |
| 66 | journals.iucr.org Internet Source | <1 % |
| 67 | ouci.dntb.gov.ua Internet Source | <1 % |
| 68 | www.hindawi.com Internet Source | <1 % |
| 69 | www.koreascience.or.kr Internet Source | <1 % |
| 70 | www2.mdpi.com Internet Source | <1 % |
| 71 | A. Preethi, Jayesh R. Bellare. "Tailoring Scaffolds for Orthopedic Application With Anti-Microbial Properties: Current Scenario and Future Prospects", Frontiers in Materials, 2021 Publication | <1 % |

72

Gebremedhin Beedemariam Gebretekle,
Atalay Mulu Fentie, Girma Tekle

<1 %

Gebremariam, Eskinder Eshetu Ali et al. "Cost-effectiveness Analysis of Caspofungin and Fluconazole for Primary Treatment of Invasive Candidiasis and Candidemia in Ethiopia", Research Square Platform LLC, 2021

Publication

73

N. Vinod Kumar, D. Sreenivasulu, Y. N. Reddy.
"Prevalence of Clostridium perfringens toxin genotypes in enterotoxemia suspected sheep flocks of Andhra Pradesh", Veterinary World, 2014

<1 %

Publication

74

Shuo Zhao, Jun-Jun Huang, Xiuyun Sun, Xiaorong Huang, Shuna Fu, Liang Yang, Xue-Wei Liu, Fei He, Yinyue Deng. " (1-aryloxy-2-hydroxypropyl)-phenylpiperazine derivatives suppress virulence by interfering with morphological transition ", Microbial Biotechnology, 2018

<1 %

Publication

75

Suwannaphet, W.. "Preventive effect of grape seed extract against high-fructose diet-induced insulin resistance and oxidative stress in rats", Food and Chemical Toxicology, 201007

<1 %

Publication

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off