

Potential Hazards of Antibiotics Resistance On Escherichia Coli Isolated From Cloacal Swab In Several Layer Poultry Farms, Blitar, Indonesia

by Freshinta Jellia Wibisono

Submission date: 29-Mar-2023 12:44PM (UTC+0700)

Submission ID: 2049757990

File name: zards_of_Antibiotics_Resistance_On_Escherichia_Coli_Isolated.pdf (392.75K)

Word count: 3193

Character count: 17854

Potential Hazards of Antibiotics Resistance On Escherichia Coli Isolated From Cloacal Swab In Several Layer Poultry Farms, Blitar, Indonesia

Adiana Mutamsari Witaningrum¹, Freshinta Jellia Wibisono², Dian Ayu Permatasari¹,
Wiwiek Tyasningsih³, Mustofa Helmi Effendi^{1,4}, Fredy Kurniawan⁵

¹Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, ²Doctoral Program Science Veterinary, Faculty of Veterinary Medicine, Universitas Gajah Mada Yogyakarta, ³Department of Veterinary Microbiology, Faculty of Veterinary Medicine, Universitas Airlangga, ⁴Halal Research Center, Airlangga University, ⁵Department of Chemistry, Institute of Sepuluh Nopember, Surabaya, Indonesia

Abstract

Objective: The study was isolated Escherichia coli from cloacal swab of layer poultry farms in Blitar area to investigate their antibiotic sensitivity pattern.

Materials and Method: Fourty swab cloacal samples were collected from 8 farms where located in Blitar for 3 months. In order for MacConkey agar to be a medium for inoculation and biochemical tests to identify isolates such as IMViC and TSIA test were performed. The method of antibiotic sensitivity pattern of Escherichia coli was tested by disk diffusion.

Results: The result revealed 40 samples, 34 samples were exposed to contamination by E. coli. The pattern of antibiotic sensitivity showed high resistance against ampicillin (62%), ciprofloxacin (56%), tetracycline and trimethoprim sulfamethoxazole (53%). Sensitive antibiotics were also observed for amoxicillin clavulanic acid, cefepime, ampicillin sulbactam, amikacin and meropenem. The presence of MDR and ESBL-producing Escherichia coli isolated from cloacal swabs of layer poultry farms in Blitar were 47.1% (16/34) and 5.9% (2/34), respectively.

Conclusion: This research to find out exposed the layer poultry farms to consider critical antibiotic resistance of E. coli and regarded potential public health hazards.

Keywords: Layer poultry farms, Escherichia coli, public health hazards, MDR, ESBL.

Introduction

Antibiotic resistance is major global health in worldwide especially in poultry, such as treatment failure economic losses and source of resistant organism that may

represent a risk to human health. Antimicrobial usage in animal production raised with intensive conditions using large amount of antimicrobial to prevent and treat disease⁽¹⁾. Enterobacteria especially Escherichia coli is the pathogenic bacteria who receive antibiotic treatment in the gastrointestinal tract of animals and humans into the environment which not only devolvement of the bacterial but antimicrobial affect more than one antibiotic and become multi drug resistance⁽²⁾. Antimicrobials are commonly used in animals which human consume the animals product and antimicrobial resistance can controlled by reduce antibiotics use for animals, overuse of antibiotic and residues can lead to more drug resistance among microbe⁽³⁾.

Corresponding Author:

Mustofa Helmi Effendi

Halal Research Center, Airlangga University, Indonesia,
Surabaya, Indonesia-60115

Telp : +628175111783

e-mail: mheffendi@yahoo.com

Escherichia coli is a Gram-negative bacterium commonly found in animal and human intestinal tract. Exposure of antibiotic use in animals to inhibit microbial can cause resistance to antibiotics, antibiotics resistance appearance of Escherichia coli in poultry is main purpose to reduce transmission of resistance in the region. Observation of antibiotics resistance especially investigate the resistance appearance of microbe commonly found in poultry. The bacteria commonly detected in the environment so that it can enter the digestive tract of animals is Escherichia coli⁽⁴⁾.

Antibiotics resistance is caused by a factor already arise on the bacteria. E. coli bacteria which genes that preserve to immunity from the influence of antibiotics derived from the plasmid. E. coli was detected possess plasmids to some drug resistance genes⁽⁵⁾. Plasmids can carrying the resistance genes in bacteria sensitive to antibiotics⁽⁶⁾. This study used some antibiotics that were often associated with data about the problem of resistance in E. coli and then the antibiotic sensitivity test to determine the profile antibiotics resistance of E. coli. The purpose of this study were also to exhibit occurrence of multidrug resistant (MDR) of E. coli and extended spectrum beta-lactamase (ESBL) producing E. coli from layer poultry farms.

Materials and Method

Isolation and Identification of E. coli: The study collected 40 samples and taken from 8 layer poultry farms in Blitar, East Java, shown on table 1. Purposive sampling of layer farm was based on some specification such as low sanitation, lack of cleanliness and proper hygiene management by the farm and low maintenance⁽⁷⁾. Samples obtained from cloacal swab brought to the laboratory in Amies medium transport wrapped sterile conditions and were taken using a cool-box⁽⁸⁾. Samples were inoculated streaked onto MacConkey media agar and incubated at 37 °C for 18 ± 24 hours⁽⁹⁾. Colonies that showed lactose-fermenting was purified and continued positive presumptive test of E. coli. Identification of bacteria were performed using morphological and biochemistry. Biochemical tests include tests Indole, Methyl Red, Vagos-Pasteur, Simon Citrate (IMViC) and TSIA to determine the level of genus and continued until the sugar fermentation test to determine the species of E. coli⁽¹⁰⁾.

Antibiotic Sensitivity Test: Antibiotic sensitivity testing was done using Kirby-Bauer disc

diffusion assay on medium Mueller-Hilton agar^(11,12). Antibiotics and concentration used was ampicillin (10 µg), chloramphenicol (30 µg), gentamicin (10 µg), ciprofloxacin (10 µg), trimethoprim-sulfamethoxazole (25 µg), ceftazidime (30 µg), amoxicillin clavulanic acid (30 µg), cefepime (30 µg), ampicillin sulbactam (20 µg), cephalosporin (30 µg), amikacin (30), tetracycline (30 µg), levofloxacin (5 µg) and meropenem (10 µg). Interpretation of the antibiotic resistance use the recommendation of the Clinical and Laboratory Standards Institute was through measurement of inhibitory zone diameter formed in study⁽¹³⁾.

Confirmation ESBL using Double Disc Synergy Test (DDST): Test for ESBL in E. coli by used disk antibiotic CAZ/Ceftazidime 30µg, AMC/Amoxicillin Clavulanic Acid 30µg, CTX/Cefotaxime 30µg, ATM/Aztreonam 30µg and inoculation on Muller-Hinton agar plate, shown on figure 2. The result showed measuring inhibition of the diameter inhibitory zone formed on Clinical and Laboratory Standards Institutions⁽¹³⁾.

Results and Discussion

In this study, a total of 40 cloacal swab samples were collected from chickens in layer poultry farms in Blitar and screened for the presence of multidrug resistant (MDR) and Extended Spectrum Beta-lactamase (ESBL)-producing E. coli, shown on table 2. Total prevalence of 47.1% of E. coli was obtained with the MDR cases and ESBL cases was 5.9% in layer chicken, shown on table 3. This agrees with the findings of Kwoji et al. where a similar occurrence of E. coli from chickens was also reported⁽¹⁴⁾.


This study used fourteen antibiotics against Escherichia coli, the results of antibiotic sensitivity test of Escherichia coli showed that the antibiotic ampicillin occurrence of resistance higher at (62%), ciprofloxacin (56%), tetracycline and trimethoprim sulfamethoxazole (53%), levofloxacin (35%), gentamicin (18%), ceftazidime and cefepime (6%) cephalosporin and chloramphenicol (3%) but in this study Escherichia coli sensitive to the antibiotic amoxicillin clavulanic acid, ampicillin sulbactam, amikacin and meropenem, shown on table 2 and figure 1. Sensitivity of microbes to antibiotic can resistant depend on commonly used of antibiotics⁽¹⁵⁾. These results are ampicillin which the highest of antibiotic resistance against Escherichia coli⁽¹⁶⁾.

Table 1. Location of samples, sample size, results of isolation of E. coli and results of MDR and ESBL cases.

Location	Number of samples	Positive E.coli	Positive MDR	Positive ESBL by DDST
Farm 1	5	4	0	0
Farm 2	5	4	2	0
Farm 3	5	5	3	0
Farm 4	5	5	5	0
Farm 5	5	4	1	0
Farm 6	5	4	1	1
Farm 7	5	4	2	1
Farm 8	5	4	2	0
Total	40	34	16	2

Table 2. Antibiotic susceptibility profiles of 34 E. coli from cloacal swabs of layer poultry farms in Blitar.

Antibiotic	Resistant (%)	Intermediate (%)	Sensitive (%)
Ceftazidime (CAZ)	2 (6%)	0	32 (94%)
Amoxicillin Clavulanic Acid (AMC)	0	0	34 (100%)
Cefepime (FEP)	2(6%)	0	32 (94%)
Ampicillin Sulbactam (SAM)	0	0	34 (100%)
Cephazolin (KZ)	1(3%)	0	33 (97%)
Gentamycin (GN)	6(18%)	0	28 (82%)
Amikacin (AK)	0	0	34 (100%)
Tetracycline (TE)	18(53%)	5(15%)	11 (32%)
Ciprofloxacin (CIP)	19 (56%)	3(9%)	12 (35%)
Levofloxacin (LEV)	12 (35%)	6(18%)	16 (47%)
Meropenem (MEM)	0	0	34 (100%)
Chloramphenicol (C)	1(3%)	0	33 (97%)
Trimethoprim-sulfamethoxazole (SXT)	18(53%)	0	16 (47%)
Ampicillin (AMP)	21 (62%)	0	13 (38%)

**Figure 1. Percentage of antibiotic resistance on E. coli**

Multidrug resistant (MDR) as an organism that is resistant to three or more antimicrobial classes⁽¹⁷⁾. One method that is often used by various researchers to characterize organisms as MDR is based on in vitro antimicrobial susceptibility test results, when researchers tested resistance to multiple antimicrobial agents, classes or subclasses of antimicrobial agents⁽¹⁸⁾. The most commonly used definitions for Gram-positive⁽¹⁹⁾ and Gram-negative bacteria that are resistant to three or more antimicrobial class⁽²⁰⁾. An overview of this variability of definitions is given in a comprehensive review of MDR⁽²¹⁾, which is used as a reference by some researchers that a large number of studies do not propose specific definitions for MDR.

There were 16 multidrug resistant (MDR) and 2 extended spectrum beta lactams (ESBL) in this study, shown on table 3. The consume of antibiotics raised antibiotics resistance contain important MDR organisms in poultry. These MDR organisms can transmission to the human through direct contact or consumption and *E. coli* is the most caused high economic losses and food contamination rates which obtained antibiotic-resistant genes with the potential to spread to other populations. The abundant use of antibiotics in poultry farms has been associated with treatment failure and the development of antibiotic resistance itself. A study showed that *E. coli* from poultry in China were resistant to at least 18 different antibiotics⁽²²⁾.

Table 3. Cases of MDR and ESBL of E. coli from cloacal swabs of Layer poultry

Sample Code	Phenotype Resistance	Type of Resistance
L2A	TE, CIP, SXT	MDR
L2B	CIP, LEV, SXT	MDR
L3A	CIP, LEV, SXT	MDR
L3D	TE, SXT, AMP	MDR
L3E	TE, SXT, AMP	MDR
L4C	TE, CIP, SXT	MDR
L7E	TE, CIP, AMP	MDR
L8B	TE, SXT, AMP	MDR
L8C	TE, CIP, LEV	MDR
L5C	TE, CIP, SXT, AMP	MDR
L7B	TE, CIP, LEV, AMP	MDR
L4B	GN, CIP, LEV, SXT, AMP	MDR
L4D	GN, CIP, LEV, SXT, AMP	MDR
L4E	GN, CIP, LEV, SXT, AMP	MDR
L6B	TE, CIP, LEV, SXT, AMP	MDR
L4A	GN, CIP, LEV, C, SXT, AMP	MDR
L6A	CAZ, FEP, GN, TE, CIP, LEV, AMP	ESBL
L7D	CAZ, FEP, KZ, GN, TE, CIP, LEV, SXT, AMP	ESBL

In our research findings, 100% of ESBL-producing isolates showed multi-drug resistance to various families of antibiotics. This finding correlates with other studies in other countries such as Switzerland⁽²³⁾, Zambia⁽²⁴⁾ and in Turkey⁽²⁵⁾ almost all ESBL-producing *E. coli* isolates found in animals are multi-resistant.

The nature of multidrug resistance of these isolates may be explained by the fact that ESBL is mediated by plasmids carrying multiresistant genes by plasmids, transposons and integrons and also they are ready to be transferred to other bacteria, not necessarily

same species. Bacteria with various resistance to antibiotics are widely distributed in animals and the environment[14]. The facts supported by recent surveys from China⁽²²⁾, Thailand⁽²⁶⁾ and Indonesia⁽²⁷⁾, have illustrated an alarming trend related to resistance among ESBL-producing organisms isolated from animals and the environment. Our results support the fact that ESBL producers provide a high level of resistance to not only third generation cephalosporins but also other non beta-lactam antibiotics groups, shown on table 3.

This study also revealed that all ESBL producers

and almost isolates that MDR showed resistance to ampicillin. In contrast, better susceptibility was observed to amikacin and no resistance was observed with meropenem. Better susceptibility to amikacin and

meropenem were also noted and can be explained by the absence of routine use of amikacin as empirical therapy on poultry farms and the absence of sufficient cross resistance with the beta-lactam antibiotic group.




Figure 2. Confirmation of ESBL by Double Disc Synergy Test (DDST)

Conclusion

The high percentage of drug resistance in *E. coli* isolates were detected ampicillin, ciprofloxacin, tetracycline and trimethoprim sulfamethoxazole more than 50%. The number of MDR of *E. coli* isolates was significantly higher in healthy poultry, namely 16 isolates and found also 2 ESBL producing *E. coli* isolates. This general description of the antimicrobial resistance of these poultry bacteria creates the basis for future investigations and analyzes of resistance development in Blitar, East Java, Indonesia. In view of this, we strongly recommend assessing treatment plans in the poultry industry in Blitar to ensure prudent antimicrobial use and to minimize the potential for the spread of resistant bacteria from poultry to the environment and humans.

Acknowledgement: This study was supported in part with the Penelitian Hibah Mandat Funding from Airlangga University, Indonesia in fiscal year 2019.

Ethical Clearance: Cloacal swabs were used in this study, hence ethical clearance was not necessary. Cloacal swab samples were collected from Blitar area in East Java province, Indonesia.

Conflict of Interest: Nil.

References

1. World Health Organization. Antimicrobial Resistance: Global Report on Surveillance. World Health Organization. France. 2014.
2. Woolhouse M, Ward M, van Bunnik B, Farrar J.

- Antimicrobial resistance in humans, livestock and the wider environment. *Phil. Trans. R. Soc. B*. 2015;370: 20140083.
3. Lee, C.R., Cho, I.H., Jeong, B.C., Lee, S.H. Strategies to minimize antibiotic resistance. *Int Environ Res Public Health*. 2013;12(Sep):10(9):4274-305. doi: 10.3390/ijerph10094274. PubMed PMID: 24036486; PubMed Central PMCID: PMC3799537.
 4. Hussain HI, Iqbal Z, Seleem MN, Huang D, Sattar A, Hao H, Yuan Z. (2017). Virulence and transcriptome profile of multidrug-resistant *Escherichia coli* from chicken. *Sci Rep*. 21;7(1):8335. doi: 10.1038/s41598-017-07798-1. PubMed PMID: 28827616; PubMed Central PMCID: PMC5567091.
 5. Costa M.M., Drescher, G., Maboni, F. Weber, S.S., Schrank, A., Vainstein, M.H. Schrank, I.S and Vargas, A.C. Virulence factors, antimicrobial resistance and plasmid content of *Escherichia coli* isolated in swine commercial farms. *Arq. Bras. Med. Vet. Zootec*. 2010;62: 30-36.
 6. Pyatov, V., I. Vrtkova and A. Knoll. Detection of Aminoglycoside, Sulfonamide and Tetracycline Resistance Genes in *Escherichia coli* Isolated from Bovine Milk Sampels. Mendel University. Brno. 2014.
 7. Effendi MH, Harijani N, Yanestria SM and Hastutiek P. Identification of Shiga Toxin-Producing *Escherichia coli* in Raw Milk Samples from Dairy Cows in Surabaya, Indonesia. *Philipp. J. Ved. Med*. 2018; 55(SI): 109-114.
 8. Dahms, C, Hübner, N-O, Krow, A, Mellmann, A, Dittmann, K, Kramer, A. Occurrence of ESBL-Producing *Escherichia coli* in Livestock and Farm Workers in MecklenburgWestern Pomerania, Germany. *PLoS ONE*. 2015;10(11): e0143326.
 9. Al Humam, N.A. Special biochemical profiles of *Escherichia coli* strains isolated from humans and camels by the VITEK 2 automated system in Al-Ahsa, Saudi Arabia. *African Journal of Microbiology Research*. 2016; Vol. 10(22): 783-790.
 10. Kavitha, J.R and Devasena, T. Molecular and Bacteriological examination of cow milk in coliform mastitis. *J.Pharmacy and Biological Science*. 2013;6(2):2319- 7676.
 11. Nassar, M.SM, Hazzah, W.A., Bakr, W.M.K. Evaluation of antibiotic susceptibility test results: how guilty a laboratory could be? *J Egypt Public Health Assoc*.2019;94(1):4. doi: 10.1186/s42506-018-0006-1. Epub 2019 Jan 11. PubMed PMID: 30686832; PubMed Central PMCID: PMC6329728.
 12. Effendi, M.H., Harijani, N., Budiarto, Triningtya, N.P., Tyasningsih, W. and Plumeriastuti, H. (2019). Prevalence of Pathogenic *Escherichia Coli* Isolated from Subclinical Mastitis in East Java Province, Indonesia. *Indian Vet. J*. 2019;96(03): 22-25.
 13. Clinical and Laboratory Standarts Institute. Performances Standards for Antimicrobial Susceptibility Testing. M100 27th ed. 2017.
 14. Kwoji, I.D., Musa, J.A., Daniel, N., Mohzo, L., Bitrus, A.A., Ojo, A, A, and Ezema, K.U. Extended-spectrum beta-lactamase-producing *Escherichia coli* in chickens from small-scale (backyard) poultry farms in Maiduguri, Nigeria. *Int J One Health*. 2019;5:26-30.
 15. Vranic, S.M. and Uzunovic, A. Antimicrobial Resistance of *Escherichia Coli* Strains Isolated From Urine At Outpatient Population: A Single Laboratory Experience. *Mater Sociomed*. 2016;25(Mar):28(2):121-124. doi: 10.5455/msm.2016.28.121-124. PubMed PMID: 27147918; PubMed Central PMCID: PMC4851537.
 16. Gregova, G., Kmetova, M., Kmet, V., Venglovsky, J. and Feher, A. Antibiotic resistance of *Escherichia coli* isolated from a poultry slaughterhouse. *Annals of Agricultural and Environmental Medicine*. 2012;19(1):.75-77.
 17. Magiorakos, A.P., Srinivasan, A., Carey, R. B., Carmeli, Y., Falagas, M. E., Giske, C. G., Harbarth, S., Hindler, J. F., Kahlmeter, G., Olsson-Liljequist, B., Paterson, D. L., Rice, L. B., Stelling, J., Struelens, M. J., Voulpoulos, A., Weber, J. T., and Monnet. D. L. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*. 2012;18: 268–281. 10.1111/j.1469-0691.2011.03570.x
 18. Kall A.J., Hidron, A.I., Patel, J., Srinivasa, n A. Multidrug resistance among gram-negative pathogens that caused healthcare-associated infections reported to the National Healthcare Safety Network, 2006-2008. *Infect Control Hosp Epidemiol*. 2010;31: 528–531.
 19. Cohen, A.L., Calfee, D., Fridkin, S.K. Recommendations for metrics for multidrug-

- resistant organisms in healthcare settings: SHEA/HICPAC Position paper. *Infect Control Hosp Epidemiol.* 2008;29: 901–913.
20. Gould, I.M. The epidemiology of antibiotic resistance. *Int J Antimicrob Agents.* 2008;32 (suppl 1): 2–9.
21. Falagas, M.E., Koletsi, P.K., Bliziotis, I.A. The diversity of definitions of multidrug-resistant (MDR) and pandrug-resistant (PDR) *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. *J Med Microbiol.* 2006;55:1619–1629.
22. Gao, L., Tan, Y., Zhang, X., Hu, J., Miao, Z., Wei, L and Chai, T. Emissions of *Escherichia coli* carrying extended-spectrum β -lactamase resistance from pig farms to the surrounding environment. *Int J Environ Res Pub Health.* 2015;12: 4203-13.
23. Kraemer, J.G., Pires, J., Kueffer, M., Semaani, E., Endimiani, A. and Hilty, M. Prevalence of extended-spectrum β -lactamase-producing Enterobacteriaceae and Methicillin-Resistant *Staphylococcus aureus* in pig farms in Switzerland. *Sci Total Environ.* 2017;Vol 603-604; 401-405
24. Chishimba, K., Hangombe, B.M., Muzandu, K., Mshana, S.E., Mutee, M.I., Nakajima, C. Detection of extended-spectrum beta-lactamase-producing *Escherichia coli* in market-ready chickens in Zambia. *Int J Microbiol.* 2016;17(Apr); 5:5275724. doi: 10.1155/2016/5275724.
25. Gundogan, S., Avci, E. Prevalence and antibiotic resistance of extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* and *Klebsiella species* isolated from foods of animal origin in Turkey. *Afr J Microbiol Res.* 2013;7: 4059-64.
26. Runcharoen, C., Raven, K.E., Reuter, S., Kallonen, T., Paksanont, S., Thammachote, J. Whole-genome sequencing of ESBL producing *Escherichia coli* isolated from patients, farm waste and canals in Thailand. *Gen Med.* 2017;9:81-93.
27. Effendi, M.H., Bintari, I.G., Aksono, E.B. & Hermawan, I.P. Detection of bla_{TEM} Gene of *Klebsiella pneumoniae* Isolated from Swab of Food-Producing Animals in East Java. *Tropical Animal Science Journal.* 2018;41(3):174-178

Potential Hazards of Antibiotics Resistance On Escherichia Coli Isolated From Cloacal Swab In Several Layer Poultry Farms, Blitar, Indonesia

ORIGINALITY REPORT

20%

SIMILARITY INDEX

14%

INTERNET SOURCES

12%

PUBLICATIONS

14%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Indonesia Student Paper	3%
2	www.scicell.org Internet Source	1%
3	www.theses.fr Internet Source	1%
4	pure.ulster.ac.uk Internet Source	1%
5	www.ijlpr.com Internet Source	1%
6	Arizah Kusumawati, Apon Zaenal Mustopa, I. Wayan Teguh Wibawan, Agus Setiyono, Mirnawati Bachrum Sudarwanto. "A sequential toggle cell-SELEX DNA aptamer for targeting Staphylococcus aureus, Streptococcus agalactiae, and Escherichia coli bacteria", Journal of Genetic Engineering and Biotechnology, 2022	1%

7 Elizabeth Christenson, Lindsay Wickersham, Megan Jacob, Jill Stewart. "A watershed study assessing effects of commercial hog operations on microbial water quality in North Carolina, USA", Science of The Total Environment, 2022

Publication

8 cienciaspecuarias.inifap.gob.mx
Internet Source

9 med.uky.edu
Internet Source

10 repository-tnmgrmu.ac.in
Internet Source

11 Submitted to Regis College
Student Paper

12 shaker.umh.es
Internet Source

13 www.e-sciencecentral.org
Internet Source

14 Submitted to Baltimore City Community College
Student Paper

15 Submitted to Canterbury Christ Church University
Student Paper

16	medworm.com Internet Source	1 %
17	Mohamed Rhouma, Francis Beaudry, Ann Letellier. "Resistance to colistin: what is the fate for this antibiotic in pig production?", <i>International Journal of Antimicrobial Agents</i> , 2016 Publication	1 %
18	Aya Seleem, Maha A. Sabry, Khaled A. Abdel-Moein. " Migratory birds as a potential overseas transmitter of Shiga toxin-producing ", <i>International Journal of Veterinary Science and Medicine</i> , 2021 Publication	1 %
19	Submitted to South University Student Paper	1 %
20	Submitted to Aga Khan University Student Paper	<1 %
21	mnet.mendelu.cz Internet Source	<1 %
22	Submitted to University of Portsmouth Student Paper	<1 %
23	trialsjournal.biomedcentral.com Internet Source	<1 %
24	Submitted to Lincoln Park High School Student Paper	<1 %

25

www.colibri.udelar.edu.uy

Internet Source

<1 %

26

Kawany Gabrieli Zanetti Fazoli, Isabela Carvalho dos Santos, Isabel Cristina da Silva Caetano, Flavio Henrique Carbonera Cervantes et al. "Antibiotic Resistance in Enterobacteriaceae Family Members Isolated from Horses Used for Animal Traction", Journal of Pure and Applied Microbiology, 2020

Publication

<1 %

27

G. A. McKay. "Time-kill kinetics of oritavancin and comparator agents against Staphylococcus aureus, Enterococcus faecalis and Enterococcus faecium", Journal of Antimicrobial Chemotherapy, 03/24/2009

Publication

<1 %

28

Christina Gagliardo, Lisa Saiman. "Chapter 14 What is the Evidence Behind Recommendations for Infection Control?", Springer Science and Business Media LLC, 2012

Publication

<1 %

29

Patrick AYONGA NDEBA, Bruce WEMBULUA SHINGA, Fatimata WONE, Alain-Bruno BARDIGUYO et al. "Antimicrobial resistance of Vibrio cholerae O1 strains isolated during cholera epidemics in eastern Democratic

<1 %

Republic of Congo between January 2011 and June 2022", Research Square Platform LLC, 2023

Publication

30

W.M. Lira. "The incidence of Shiga toxin-producing Escherichia coli in cattle with mastitis in Brazil", Journal of Applied Microbiology, 10/2004

Publication

<1 %

31

feyz.kaums.ac.ir

Internet Source

<1 %

32

"Posters", FEMS Microbiology Letters, 2003

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off