

# Multidrug resistance-encoding gene in *Citrobacter freundii* isolated from healthy laying chicken in Blitar District, Indonesia

*by* Freshinta Jellia Wibisono

---

**Submission date:** 30-Mar-2023 04:51PM (UTC+0700)

**Submission ID:** 2050885293

**File name:** sistance-encoding\_gene\_in\_Citrobacter\_freundii\_isolated\_from.pdf (607.63K)

**Word count:** 4747

**Character count:** 26609

## Multidrug resistance-encoding gene in *Citrobacter freundii* isolated from healthy laying chicken in Blitar District, Indonesia

Adiana Mutamsari Witaningrum<sup>1</sup>, Freshinta Jellia Wibisono<sup>2</sup>, Dian Ayu Permatasari<sup>1</sup>, Mustofa Helmi Effendi<sup>1</sup>, and Emmanuel Nnabuike Ugbo<sup>3</sup>

1. Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Kampus C Mulyorejo, Surabaya 60115, Indonesia; 2. Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Wijaya Kusuma Surabaya, Jl. DukuhKupang XXV No.54, Surabaya 60225, Indonesia; 3. Department of Applied Microbiology, Faculty of Science, Ebonyi State University, Abakaliki, Nigeria.

**Corresponding author:** Mustofa Helmi Effendi, e-mail: [mustofa-h-e@fkh.unair.ac.id](mailto:mustofa-h-e@fkh.unair.ac.id)

**Co-authors:** AMW: [adiana\\_mutam@yahoo.co.id](mailto:adiana_mutam@yahoo.co.id), FJW: [freshinta.uwks@gmail.com](mailto:freshinta.uwks@gmail.com), DAP: [permatasariyudian@gmail.com](mailto:permatasariyudian@gmail.com), ENU: [ugbonuel2001@yahoo.com](mailto:ugbonuel2001@yahoo.com)

**Received:** 25-03-2022, **Accepted:** 11-10-2022, **Published online:** 12-12-2022

**doi:** [www.doi.org/10.14202/IJOH.2022.161-166](http://www.doi.org/10.14202/IJOH.2022.161-166) **How to cite this article:** Witaningrum AM, Wibisono FJ, Permatasari DA, Effendi MH, and Ugbo EN (2022) Multidrug resistance-encoding gene in *Citrobacter freundii* isolated from healthy laying chicken in Blitar District, Indonesia, *Int. J. One Health*, 8(2): 161-166.

### Abstract

**Background and Aim:** The increasing prevalence of resistance (MDR) of Enterobacteriaceae in Indonesia has caused concern regarding human health. *Citrobacter freundii* reportedly targets the gastrointestinal tract of animals and is a common cause of foodborne diseases associated with diarrhea, peritonitis, meningitis, brain abscess, bacteremia, and urinary tract infection. This study aimed to estimate the prevalence of MDR and the presence of Class 1 integron-encoding genes in *C. freundii* isolates obtained from cloacal swabs of healthy laying chickens in Blitar district, Indonesia.

**Materials and Methods:** One hundred and sixty-five cloacal swab samples were collected from 33 farms in Blitar over a period of 25 months. Standard microbiological techniques such as bacterial culture in MacConkey agar, Simmons citrate agar, and triple sugar iron agar and biochemical tests such as the indole test were performed to identify the isolates. The antibiotic sensitivity patterns of *C. freundii* isolates were determined by the disk diffusion method, and MDR-encoding genes (Class 1 integron) were detected by polymerase chain reaction (PCR).

**Results:** Out of 165 cloacal swab samples, 7 (4.24%) were positive for *C. freundii*. *Citrobacter freundii* was highly resistant to erythromycin (71.43%) and moderately to streptomycin, tetracycline, and trimethoprim-sulfamethoxazole (all 42.86%); however, it showed low resistance to ampicillin (28.57%). All isolates were found to exhibit MDR. Only 1 (14.29%) of the seven *C. freundii* isolates harbored a Class 1 integron gene. This study revealed that Class 1 integron-encoding genes have a low prevalence in *C. freundii* isolated from healthy laying chickens in Blitar, Indonesia.

**Conclusion:** Poultry animals can play a role in the transmission of resistance genes to humans due to the MDR of Enterobacteriaceae, including *C. freundii* in the intestines.

**Keywords:** *Citrobacter freundii*, Class 1 integron gene, human health, laying chickens, multidrug resistance.

### Introduction

*Citrobacter freundii*, a member of the Enterobacteriaceae family, is a facultative anaerobic, Gram-negative bacillus. It has flagella-driven motility and is commonly found in water, soil, food, and the intestines of animals and humans [1]. This species reportedly targets the gastrointestinal tract of animals and is a common cause of foodborne diseases, leading to diarrhea, peritonitis, meningitis, brain abscess, bacteremia, and urinary tract infection [2].

Livestock products are important for food security in Blitar, Indonesia, where the cost of meat has been a focus of the government, academia, and consumers. Poultry chicken and products such as eggs are

the most widely consumed types of meat and livestock products in Indonesia, accounting for more than 70% of total meat consumption [3]. Thus, their products are the most common sources of contamination by infectious bacteria that can cause diseases in humans and animals [4]. *Citrobacter freundii* is a bacterium for which antibiotic treatment can be applied in the gastrointestinal tract of animals and humans [5]. Antibiotics can inhibit the growth of microorganisms, although treatment with multiple antibiotics can cause multidrug resistance (MDR) [6]. Recently, MDR has been recognized as a global public health problem [7, 8]. Multidrug resistance of *C. freundii* has also increased worldwide [9, 10].

Integrons are genetic structures expressing genes usually encoding antibiotic resistance [11]. Class 1 integrons include the characteristic features of integrons while also being mobile elements. They are widely distributed in Gram-negative bacteria found clinically and in the environment and are responsible for the dissemination of different cassette-associated antibiotic-resistance genes. The Class 1 integrons

Copyright: Witaningrum, et al. This article is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

are a highly prevalent source of the spread of MDR among Gram-negative bacteria [12, 13] and are found at high frequencies in pathogens and commensals isolated from livestock, creating a worldwide crisis in the management of bacterial infections [14]. Insufficient data have been obtained on the antimicrobial resistance of *C. freundii* isolates harboring Class 1 integron genes isolated from cloacal swabs of healthy laying chickens.

Thus, this study aimed to estimate the prevalence of MDR and the presence of Class 1 integron-encoding genes in *C. freundii* isolates obtained from cloacal swabs of healthy laying chickens in Blitar, Indonesia.

## Materials and Methods

### Ethical approval

Cloacal swabs were used in this study; hence, ethical approval was not necessary. Cloacal swab samples were collected from laying chicken farms, Blitar district, Indonesia, as per standard collection procedure.

### Study period and location

This cross-sectional study was conducted from June to August 2019. One hundred and sixty-five cloacal swab samples were collected from laying chicken farms in Blitar district, Indonesia.

### Sample isolation and identification

The samples were collected from 33 farms in Blitar, Indonesia, which included 165 cloacal swabs of healthy laying chickens. Farms were selected on the basis of purposive sampling methods. The cloacal swab samples were cultured in MacConkey agar plates (Oxoid, Cheshire, UK) and incubated at 37°C for 24 h [15]. The colonies showing positivity for lactose-fermenting bacteria were subcultured to obtain pure cultures, followed by biochemical tests (Indole test, Simmons citrate agar, and TSIA) to determine *C. freundii* [16].

### Antibiotic sensitivity test

Antibiotic sensitivity testing was performed using the Kirby-Bauer disk diffusion assay on Mueller-Hinton agar medium [17]. The suspension of 3 mL of freshly prepared pure culture colonies was standardized to an equivalent of 0.5 McFarland and inoculated onto culture plates. The antibiotic disks used were as follows: Erythromycin (E, 15 µg), streptomycin (S, 10 µg), tetracycline (TE, 30 µg), trimethoprim-sulfamethoxazole (SXT, 25 µg), and ampicillin (AMP, µg). After the attachment of the antibiotic disks, the plates were then incubated at 37°C for 24 h. Interpretations of antibiotic resistance are performed in line with the recommendations in the Clinical and Laboratory Standards Institute guidelines. Multidrug resistance was classified for those isolates that were resistant to more than two of the different classes of antibiotics [18].

### Detection of Class 1 integron-encoding gene

Multidrug-resistant (MDR) *C. freundii* was further analyzed for the presence of Class 1

integron-encoding genes using molecular polymerase chain reaction (PCR) identification techniques. Bacterial DNA of *C. freundii* isolates was extracted with the QIAamp® DNA Mini Kit (Qiagen, Germany). The primers used to detect Class 1 integron-encoding genes were hep 58 (TCATGGCTTGTTATGACTGT) and hep 59 (GTAGGGCTTATTATGCACGC). The PCR conditions with denaturation temperatures for 2 minutes at 94°C; extended denaturation at 94°C for 30 seconds; annealing at 55°C for 45 seconds; extension at 72°C for 45 seconds; final extension at 72°C for 7 minutes, this reaction was carried out for 30 cycles [19]. Polymerase chain reaction results were visualized by electrophoresis using 1.5% agarose gel (Invitrogen, USA) [20, 21].

## Results

Among 165 cloacal swab samples collected from healthy laying chickens at poultry farms in Blitar, Indonesia, seven samples were positive for *C. freundii*. The prevalence of *C. freundii* in chickens was 4.24% among the 33 farms (Table-1). Five panels of different classes of antibiotics were tested against *C. freundii* isolates. The antibiotic sensitivity test indicated that *C. freundii* isolates were resistant to E at a high rate of 71.43%. Intermediate rates of resistance were recorded for S, TE, and SXT (42.86% for each), while a low rate was observed for AMP (28.57%) (Table-2 and Figure-1). This study clearly identified different levels of antibiotic resistance among *C. freundii* isolates derived from animals. The highest level of antibiotic resistance being observed for E is an indication that this antibiotic has been overused or abused in animal health care [22]. Five (71.43%) of the seven *C. freundii* isolates were discovered to be MDR (Table-3).

## Discussion

This study provides evidence that *C. freundii* of animal origin is a possible pathogen that can cause foodborne diseases. The prevalence of *C. freundii* in chickens was 4.24% among 33 farms. This is in accordance with the findings of Liu *et al.* [1], who stated that the prevalence of *C. freundii* was approximately 3%–6% among the isolates of Enterobacteriaceae in samples. This research also agreed with the previous studies recognizing *C. freundii* as an animal pathogen that causes foodborne outbreaks worldwide [23]. The identification of MDR *C. freundii*, as reported in this study, is in line with other studies on organisms that belong to the same family of Enterobacteriaceae [9]. This finding is a major public health concern. Previous

**Table-1:** Prevalence of *C. freundii* from laying chickens in Blitar, Indonesia.

Samples	165
<i>C. freundii</i> positive	7 (7/165; 4.24%)
Class 1 integron	1 (1/7; 14.29%)
<i>C. freundii</i> = <i>Citrobacter freundii</i>	

**Table-2:** Antibiotic susceptibility profile of *C. freundii* isolated from laying chickens in Blitar, Indonesia.

Antibiotic	Resistant (%)	Intermediate (%)	Sensitive (%)
Erythromycin	5 (71.43%)	2 (28.57%)	0
Streptomycin	3 (42.86%)	0	4 (57.14%)
Tetracycline	3 (42.86%)	0	4 (57.14%)
Trimethoprim-sulfamethoxazole	3 (42.86%)	0	4 (57.14%)
Ampicillin	2 (28.57%)	1 (14.29%)	4 (57.14%)

*C. freundii* = *Citrobacter freundii*



**Table-3:** Prevalence of MDR and Class 1 integron in *C. freundii* isolated from laying chickens.

Location	Code of <i>C. freundii</i>	MDR	Class 1 integron by PCR
hFarm 1	L1dn	3 MDR	1
Farm 5	L5an	4 MDR	0
Farm 10	L10cn	1 R	0
Farm 11	L11dn	4 MDR	0
Farm 16	L16an	2 R	0
Farm 26	L26dn	5 MDR	0
Farm 29	L29an	5 MDR	0
Total	7	5 (71.43%)	1 (14.29%)

*C. freundii* = *Citrobacter freundii*, MDR = Multidrug resistant, R = Resistant, PCR = Polymerase chain reaction

studies from all over the world have revealed that the prevalence of MDR bacterial pathogens is now a major public health issue [24–26]. Organisms that are resistant to three or more classes of antimicrobials are called MDR [27, 28]. *In vitro* antimicrobial susceptibility testing is commonly used to characterize organisms as MDR bacteria [29].

Using a molecular method, 1 (14.29%) of the seven *C. freundii* isolates was identified to harbor a Class 1 integron-encoding gene. The amplification of such a gene from *C. freundii* in healthy layer chickens was observed at an amplicon size of 200 bp (Figure-2) in the isolates recovered from cloacal swab samples using the primers hep 58 and hep 59 [30]. The results indicate that resistance to several different antibiotics is associated with the presence of Class 1 integron. This finding is in agreement with studies that observed MDR bacteria harboring integron genes that boost their infectivity and resistance to many classes of antibiotics [31]. Other studies have detected integrons in Gram-negative bacteria, which have also increasingly been discovered in Gram-negative bacteria [32, 33]. Class 1 integron gene was also reported as one of the fastest spreading [51](#)istance genes in Gram-negative bacteria [34, 35]. Integrons are genetic elements that can possess MDR genes; their existence can obstruct the treatment of bacterial infections [36]. Integrons are classified based on differences in the gene structure of integrases [37]. Class 1 integrons have been found more frequently in Gram-negative bacteria among the different classes of integrons [14]. This study has provided genetic evidence of the presence of an integron-encoding gene in *C. freundii*, a Gram-negative bacterium. The importance of Class 1 integrons as repositories of MDR has been reported [38], and there is increasing evidence for the presence of

**Figure-1:** Antibiotic resistance pattern of *Citrobacter freundii* isolated from laying chickens. E=Erythromycin, S=Streptomycin, TE=Tetracycline, SXT=Trimethoprim-sulfamethoxazole, AMP=Ampicillin.**Figure-2:** Class 1 integron gene identified from *Citrobacter freundii* using polymerase chain reaction from laying chickens in Blitar, Indonesia. MRK=Marker (100 bp); K+=Positive control; K-=Negative control; L1DN-L29an=Samples, identified amplicon size of Class 1 integron encoding gene = 200 bp.

49

genes encoding Class 1 integrons in poor sanitation. The presence of Class 1 integron-encoding gene in an amplicon obtained using hep 58–59 primers suggested the possibility of MDR gene, which was confirmed by studying the isolated strains. These results indicate that Class 1 integron genes are present in poultry animals and their products. *Citrobacter freundii* has mostly been collected from poultry [39, 40]. This bacterium is potentially pathogenic to animals and humans, being cytotoxic and aggregative [2]. Aggregative adherence to epithelial cells has been recognized as a putative virulence factor contributing to bacterial pathogenicity [2]. The presence of integrons in MDR properties of *C. freundii* will add to the difficulties in treating infections by this bacterium [41–43].

Recent studies on environmental pollution have confirmed that MDR bacteria can be transmitted to humans and poultry animals through various

distribution routes [44]. The MDR bacteria involvement in Enterobacteriaceae is well established as an important part of global public health policy. This idea that poultry is a good indicator as a source for MDR bacteria from environmental pollution and argues the importance of the One Health approach because these poultry animals can significantly contribute indirectly to the transmission of resistant genes to other segments of environments [45]. Immediate efforts are required to deal with the emergence of MDR bacteria, including antimicrobial resistance in humans and poultry. Antimicrobial resistance from animals has become a public health issue, as evidence has shown the transmission of antimicrobial-resistant bacteria, including Gram-positive bacteria [46–48], or their resistance genes, between animals and humans [49]. Poultry animals can play a role in the transmission of resistance genes to humans due to the MDR of Enterobacteriaceae, including *C. freundii* in the intestines.

### Conclusion

The prevalence of *C. freundii* in healthy laying chickens was observed to be low, with an incidence of 37.3%. *Citrobacter freundii* was discovered, show resistance to multiple classes of antibiotics. A Class 1 integron-encoding gene was identified in *C. freundii* of animal origin. The finding of MDR *C. freundii* with a plasmid containing a characteristic Class 1 integron supported the concept that there is an environmental reservoir of resistance. Exploration of the molecular genetic features of *C. freundii* harboring a Class 1 integron gene in poultry may also facilitate epidemiological monitoring and source tracking of resistance determinants and their environments. This would deepen our understanding of the ecological cycle of antibiotic resistance. This may enable practical interventions to reduce the traffic of resistance genes through the use of appropriate and effective antibiotics in laying hens, good farm biosecurity practices, and proper hygiene in farms. As such, this study provides a basis for improving public health.

### Authors' Contributions

MHE and FJW: Conceptualization. MHE, AMW, and DAP: Data curation. FJW and AMW: Formal analysis. MHE and FJW: Funding acquisition. AMW and DAP: Investigation. MHE and AMW: Methodology. MHE and AMW: Project administration. MHE, AMW, and DAP: Resources. MHE and FJW: Supervision. MHE and ENU: Validation. FJW and AMW: Visualization. MHE, FJW, and ENU: Writing original draft. MHE and ENU: Writing – review and editing. All authors have read and approved the final manuscript.

### Acknowledgments

This study was partly funded by the Direktorat Riset dan Pengabdian Masyarakat, Deputi Bidang

Penguatan Riset dan Pengembangan, Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi, Indonesia in the fiscal year 2022 (Grant number: 761/UN3.15/PT/2022).

### Competing Interests

The authors declare that they have no competing interests.

### Publisher's Note

Veterinary World (Publisher of International Journal of One Health) remains neutral with regard to jurisdictional claims in published institutional affiliation.

### References

- Liu, L.H., Wang, N.Y., Wu, A.Y.J., Lin, C.C., Lee, C.M. and Liu, C.P. (2018) *Citrobacter freundii* bacteremia: Risk factors of mortality and prevalence of resistance genes. *J. Biol. Microbiol. Immunol. Infect.*, 51(4): 565–572.
- Bai, L., Xia, S., Lan, R., Liu, L., Ye, C. and Wang, Y. (2012) Isolation and characterization of cytotoxic, aggregative. *Citrobacter freundii*. *PLoS One*, 7: e33054.
- Effendi, M.H., Tyasningsih, W., Yurianti, Y.A., Rahmahani, J., Harijani, N. and Plumeriastuti, H. (2021) Prevalence of multidrug resistance (MDR) and extended-spectrum beta-lactamase (ESBL) of *Escherichia coli* isolated from cloacal swabs of broilers in several wet markets in Surabaya, Indonesia. *Biodiversitas*, 22(1): 304–310.
- Ariza-Proto, M.A., Pando-Sandoval, A., Garcia-Clemente, M., Fernández, R. and Casan, P. (2015) Community-acquired pneumonia and empyema caused by *Citrobacter koseri* in an immunocompetent patient. *Case Rep. Pulmonol.*, 2015: 670373.
- Hidayatullah, A.R., Effendi, M.H., Plumeriastuti, H., Wibisono, F.M., Hartadi, E.B. and Sofiana, E.D. (2020) A review of the opportunistic pathogen *Citrobacter freundii* in piglets post-weaning: Public health importance. *Syst. Rev. Pharm.*, 11(9): 767–773.
- Rahmahani, J., Salamah, Mufasirin, Tyasningsih, W. and Effendi, M.H. (2020) Antimicrobial resistance profile of *Escherichia coli* from cloacal swab of domestic chicken in Surabaya Traditional Market. *Biochem. Cell. Arch.*, 20(1): 2993–2997.
- Sharieta, R., Effendi, M.H., Plumeriastuti, H. (2021) Genetic identification of shiga toxin encoding gene from cases of multidrug resistance (MDR) *Escherichia coli* isolated from raw milk. *Trop. Anim. Sci. J.*, 44(1): 10–15.
- Widodo, A., Effendi, M.H. and Khairullah, A.R. (2020) Extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* from livestock. *Syst. Rev. Pharm.*, 11(7): 382–392.
- Liu, L., Chen, D., Liu, L., Lan, R., Hao, S., Jin, W., Sun, H., Wang, Y., Liang, Y. and Xu, J. (2018) Genetic diversity, multidrug resistance, and virulence of *Citrobacter freundii* from diarrheal patients and healthy individuals. *Front. Cell. Infect. Microbiol.*, 8: 233.
- Pepperell, C., Kus, J.V., Gardam, M.A., Humar, A. and Burrows, L.L. (2002) Low-virulence *Citrobacter* species encode resistance to multiple antimicrobials. *Antimicrob. Agents Chemother.*, 46(11): 3555–3560.
- Gillings, M.R. (2014) Integrons: Past, present, and future. *Microbiol. Mol. Biol. Rev.*, 78(2): 257–277.
- Kargar, M., Mohammadalipour, J., Doost, A., Lorzadeh, S. and Japoni-Nejad, A. (2014) High prevalence of class 1 to 3 integrons among multidrug-resistant diarrheagenic *Escherichia coli* in Southwest of Iran. *Osong Public Health Res. Perspect.*, 5(4): 193–198.
- Permatasari, D.A., Witaningrum, A.M., Wibisono, F.J. and

- Effendi, M.H. (2020) Detection and prevalence of multi-drug-resistant *Klebsiella pneumoniae* strains isolated from poultry farms in Blitar, Indonesia. *Biodiversitas*, 21(10): 242–4647.
14. Gillings, M., Boucher, Y., Labbate, M., Holmes, A., Krishnan, S., Holley, M. and Stokes HW. (2008) The evolution of class 1 integrons and the rise of antibiotic resistance. *J. Bacteriol. Res.*, 190(14): 5095–5100.
  15. Al Humam, N.A. (2016) Special biochemical profiles of *Escherichia coli* strains isolated from humans and camels by the VITEK 2 automated system in Al-Ahsa, Saudi Arabia. *Afr. J. Microbiol. Res.* 10(22): 783–790.
  16. Janda, J.M., Abbott, S.L., Cheung, W.K.W. and Hanson, D.F. (1994) Biochemical identification of citrobacteria in the clinical laboratory. *J. Clin. Microbiol.*, 32 (8): 1850–1854.
  17. Nassar, M.S.M., Hazzah, W.A. and Bakr, W.M.K. (2019) Evaluation of antibiotic susceptibility test results: How guilty a laboratory could be? *J. Egypt Public Health Assoc.*, 24(1): 4.
  18. Clinical and Laboratory Standards Institute (2017) Performances Standards for Antimicrobial Susceptibility Testing. M100 27<sup>th</sup> ed. Clinical and Laboratory Standards Institute, United States
  19. Singh, T., Das, S., Ramachandran, V.G., Wani, S., Shah, D., Maroof, K.A. and Sharma, A. (2017) Distribution of integrons and phylogenetic groups among 45 enteropathogenic *Escherichia coli* isolates from children <5 years of age in Delhi, India. *Front. Microbiol.*, 8: 561.
  20. Yanestria, S.M., Rahmaniari, R.P., Wibisono, F.J. and Effendi, M.H. (2019) Detection of *invA* gene of *Salmonella* from milkfish (*Chanos chanos*) at Sidoarjo wet fish market, Indonesia, using polymerase chain reaction technique. *Vet. World*, 12(1): 170–175.
  21. Ansharieta, R., Effendi, M.H., Ramandinianto, S.C. and Plumeriastuti, H. (2021) Molecular identification of *bla*<sub>CTX-M</sub> and *bla*<sub>TEM</sub> genes encoding extended-spectrum  $\beta$ -lactamase (ESBL) producing *Escherichia coli* isolated from raw cow's milk in East Java, Indonesia. *Biodiversitas*, 22(4): 1600–1605.
  22. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A. and Witaningrum, A.M. (2020) The presence of extended-spectrum beta-lactamase (ESBL) Producing *Escherichia coli* on layer chicken farms in Blitar Area, Indonesia. *Biodiversitas*, 21(6): 2667–2671.
  23. Aminharati, F., Ehrampoush, M.H., Dal, M.M., Yaseri, M., Tafti, A.A. and Rajabi, Z. (2019) *Citrobacter freundii* foodborne disease outbreak related to environmental conditions in Yazd province, Iran. *Iran J. Public Health*, 9(6): 1099–1105.
  24. Van Duin, D. and Paterson, D.L. (2016) Multidrug-resistant bacteria in the community: Trends and lessons learned. *Infect. Dis. Clin. North Am.*, 30(2): 377–390.
  25. Aslam, B., Wang, W., Arshad, M.I., Khurshid, M., Muzammil, S., Rasool, M.H., Nisar, M.A., Alvi, R.F., Aslam, M.A., Qamar, M.U., Salamat, M. and Baloch, Z. (2018) Antibiotic resistance: A rundown of a global crisis. *Infect. Drug Resist.*, 11: 1645–1658.
  26. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A. and Witaningrum, A.M. (2020) Short communication: Pattern of antibiotic resistance on extended-spectrum beta-lactamases genes producing *Escherichia coli* on laying hens in Blitar, Indonesia. *Biodiversitas*, 21(10): 4631–4635.
  27. Magiorakos, A.P., Srinivasan, A., Carey, R.B., Carmeli, Y., Falagas, M.E., Giske, C.G., Harbarth, S., Hindler, J.F., Kahlmeter, G., Lilijequist, B.O., Paterson, D.L., Rice, L.B., Stelling, J., Struelens, M., Vatopoulos, A., Weber, J.T. and Monnet, D.L. (2012) Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol. Infect.*, 18(3): 268–281.
  28. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A. and Witaningrum, A.M. (2021) Molecular identification of CTX gene of extended-spectrum beta-lactamases (ESBL) producing *Escherichia coli* on layer chicken in Blitar, Indonesia. *J. Anim. Plant Sci.*, 31(4): 954–959.
  29. Kallen, A.J., Hidron, A.I., Patel, J. and Srinivasan, A. (2010) Multidrug resistance among gram-negative pathogens that caused healthcare-associated infections reported to the national healthcare safety network, 2006–2008. *Infect. Control Hosp. Epidemiol.*, 31(5): 528–531.
  30. Nagachinta, S. and Chen, J. (2009) Integron-mediated antibiotic resistance in Shiga toxin-producing *Escherichia coli*. *Food Prot.*, 72(1): 21–27.
  31. Akrami, F., Rajabnia, M. and Pourmajaf, A. (2019) Resistance integrons; A Mini review. *Casp. J. Intern. Med.*, 1(4): 370–376.
  32. Deng, Y., Bao, X., Ji, L., Chen, L., Liu, J., Miao, J., Chen, D., Bian, H., Li, Y. and Yu, G. (2015) Resistance integrons: Class 1, 2 and 3 integrons. *Ann. Clin. Microbiol.*, 14: 45.
  33. Pormohammad, A., Pouriran, R., Azimi, H. and Goudarzi, M. (2019) Prevalence of integron classes in Gram-negative clinical isolated bacteria in Iran: A systematic review and meta-analysis. *Iran. J. Basic Med. Sci.*, 22(2): 118–127.
  34. Deng, Y., Wu, Y., Jiang, L., Tan, A., Zhang, R. and Luo, L. (2016) Multidrug resistance mediated by Class 1 Integrons in *Aeromonas* isolated from farmed freshwater animals. *Front. Microbiol.*, 7: 935.
  35. Jones-Dias, D., Manageiro, V., Ferreira, E., Barreiro, P., Vieira, L., Moura, I.B. and Caniça, M. (2016) Architecture of Class 1, 2, and 3 integrons from gram-negative bacteria recovered among fruits and vegetables. *Front. Microbiol.*, 7: 1400.
  36. Mostafa, M., Siadat, S.D., Shahcheraghi, F., Vaziri, F., Japoni-Nejad, A., Yousefi, J., Rajaei, B., Mood, E.H., Zadeh, N.E., Moshiri, A., Siamdoust, S.A. and Rahbar, M. (2015) Variability in gene cassette patterns of class 1 and 2 integrons associated with multi drug resistance patterns in *Staphylococcus aureus* clinical isolates in Tehran-Iran. *BMC Microbiol.*, 15(1): 152.
  37. Cury, J., Jové, T., Touchon, M., Néron, B. and Rocha, E.P. (2016) Identification and analysis of integrons and cassette arrays in bacterial genomes. *Nucleic Acids Res.*, 44(10): 4539–4550.
  38. Yim, G., Kwong, W., Davies, J. and Miao, V. (2013) Complex integrons containing *qnrB4-ampC (blaDHA-1)* in plasmids of multidrug-resistant *Citrobacter freundii* from wastewater. *Can. J. Microbiol.* 59(2): 110–116.
  39. Zhou, W., Chen, Q., Shen, K., Zhu, X., Zhou, J., Lu, W., Sun, Z., Liu, H., Li, K., Xu, T., Bao, Q. and Lu, J. (2019) *In vitro* susceptibility and florfenicol resistance in *Citrobacter* isolates and whole-genome analysis of multidrug-resistant *Citrobacter freundii*. *Int. J. Genomics.*, 2019: 7191935.
  40. Krauland, M.G., Marsh, J.W., Paterson, D.L. and Harrison, L.H. (2009) Integron-mediated multidrug resistance in a global collection of nontyphoidal *Salmonella enterica* isolates. *Emerg. Infect. Dis.*, 15(3): 388–396.
  41. Ansharieta, R., Effendi, M.H. and Plumeriastuti, H. (2020) Detection of multidrug-resistant (MDR) *Escherichia coli* isolated from raw milk in east Java Province, Indonesia. *Indian J. Forensic Med. Toxicol.*, 14(4): 3403–3407.
  42. Witaningrum, A.M., Wibisono, F.J., Permatasari, D.A. and Effendi, M.H. (2021) Detection of Class 1 integron encoding gene in multidrug resistance (MDR) of *Citrobacter freundii* isolated from healthy broiler chicken. *Trop. Anim. Sci. J.*, 44(3): 363–368.
  43. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A. and Witaningrum, A.M. (2021) Cases of multidrug resistance (MDR) and extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* from broiler chicken in Blitar, Indonesia. *Biochem. Cell. Arch.*, 21(Suppl 1): 1923–1929.

- 43
44. Orh, M.K., Kwaga, J.K.P., Hendriksen, R.S., Okol, E.C. and Thakur, S. (2021) Genetic relatedness of multidrug-resistant *Escherichia coli* isolated from humans, chickens and poultry environments. *Antimicrob. Resist. Infect. Control*, 10: 58.
45. Sudatip, D, Chasiri, K., Kritiyakan, A., Phanprasit, W., Thinphovong, C., Tiengrim, S., Thamlikitul, V., Abdallah, R., Baron, SA., Rolain, J.M., Morand, S., Hilty M. and Oppliger, A. (2021) A one health approach to assessing occupational exposure to antimicrobial resistance in Thailand: The FamResist project. *PLoS One*, 16(1): e0245250.
46. Effendi, M.H., Oktavianto, A. and Hastutiek, P. (2018) Tetracycline resistance gene in *Streptococcus agalactiae* isolated from bovine subclinical mastitis in Surabaya, Indonesia. *Philipp. J. Vet. Med.*, 55(SI): 115–120.
47. Rahmaniar, R.P., Yunita, M.N., Effendi, M.H. and Yanest S.M. (2020) Encoding gene for methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from Nasal Swab of Dogs. *Indian Vet. J.*, 97(2): 37–40.
48. Iline, V., Effendi, M.H., Rahmaniar, R.P., Yanestria, S.M. and Harijani, N. (2020) Profile of antibiotic-resistant and presence of methicillin-resistant *Staphylococcus aureus* from nasal swab of dogs from several animal clinics in Surabaya, Indonesia. *Int. J. One Health*, 6(1): 90–94.
49. Riwu, K.H.P., Effendi, M.H. and Rantam, F.A. (2020) A review of extended-spectrum  $\beta$ -lactamase (ESBL) producing *Klebsiella pneumoniae* and multidrug-resistant (MDR) on companion animals. *Syst Rev. Pharm.*, 11(7): 270–277.

\*\*\*\*\*

# Multidrug resistance-encoding gene in *Citrobacter freundii* isolated from healthy laying chicken in Blitar District, Indonesia

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

13%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to Liverpool Hope Student Paper	1%
2	researchers.mq.edu.au Internet Source	1%
3	zombiedoc.com Internet Source	1%
4	R. Ansharieta, M. H. Effendi, H. Plumeriastuti. "Genetic Identification of Shiga Toxin Encoding Gene from Cases of Multidrug Resistance (MDR) Escherichia coli Isolated from Raw Milk", Tropical Animal Science Journal, 2021 Publication	1%
5	cdstest.net Internet Source	1%
6	C. Yelnik. "CMI editorial report 2012 : Editorial", Clinical Microbiology and Infection, 03/2012 Publication	1%



7	Submitted to Cornell University Student Paper	1 %
8	pesquisa.bvsalud.org Internet Source	1 %
9	www.ijcmas.com Internet Source	1 %
10	www.indianjournals.com Internet Source	1 %
11	ir.ymlib.yonsei.ac.kr Internet Source	1 %
12	Sejalben S. Chaudhari, Harshadkumar C. Chauhan, Kishan K. Sharma, Sandip S. Patel et al. "Antibiotic susceptibility pattern of canine coagulase positive and coagulase negative Staphylococcus spp. in a hot and dry region of India", Topics in Companion Animal Medicine, 2022 Publication	<1 %
13	Submitted to Georgia Virtual Technical College Student Paper	<1 %
14	annals.org Internet Source	<1 %
15	Submitted to Petroleum Research & Development Center Student Paper	<1 %

16 Ronald Rodriguez, Nick Campbell-Kruger, Jesus Gonzalez Camba, John Berude, Rachel Fetterman, Sarah Stanley. " MarR-Dependent Transcriptional Regulation of Induces Ethionamide Resistance in Mycobacterium abscessus ", Antimicrobial Agents and Chemotherapy, 2023  
Publication

---

17 Submitted to Massey University  
Student Paper

---

18 Submitted to University of the Sunshine Coast  
Student Paper

---

19 m.beyotime.com  
Internet Source

---

20 Submitted to Institut Seni Indonesia Surakarta  
Student Paper

---

21 Submitted to Royal Veterinary College  
Student Paper

---

22 archive.aessweb.com  
Internet Source

---

23 udsspace.uds.edu.gh  
Internet Source

---

24 www.thepharmajournal.com  
Internet Source

---

- 25 E.O. Nwankwo, N. Ekwunife, K.C. Mofolorunsho. "Nosocomial pathogens associated with the mobile phones of healthcare workers in a hospital in Anyigba, Kogi state, Nigeria", *Journal of Epidemiology and Global Health*, 2014  
Publication <1 %
- 
- 26 [brage.bibsys.no](http://brage.bibsys.no)  
Internet Source <1 %
- 
- 27 [www.e-jyms.org](http://www.e-jyms.org)  
Internet Source <1 %
- 
- 28 Özge Kahraman-Ilıkkın. "Comparison of *Propionibacterium* genomes: CRISPR-Cas systems, phage/plasmid diversity, and insertion sequences", *Archives of Microbiology*, 2022  
Publication <1 %
- 
- 29 M. Gillings, Y. Boucher, M. Labbate, A. Holmes, S. Krishnan, M. Holley, H. W. Stokes. "The Evolution of Class 1 Integrons and the Rise of Antibiotic Resistance", *Journal of Bacteriology*, 2008  
Publication <1 %
- 
- 30 Ashraf M Ahmed, Tadashi Shimamoto. " A plasmid-encoded class 1 integron carrying , a putative phosphoserine phosphatase gene <1 %

and from enterotoxigenic O159 isolated in Japan ", FEMS Microbiology Letters, 2004

Publication

31

L. Awawdeh, C. Turni, JL. Mollinger, J. Henning, RN. Cobbold, DJ. Trott, J.S. Gibson. "

Antimicrobial susceptibility, plasmid replicon typing, phylogenetic grouping, and virulence potential of avian pathogenic and faecal isolated from meat chickens in Australia ", Avian Pathology, 2022

Publication

<1 %

32

[jabonline.a2hosted.com](http://jabonline.a2hosted.com)

Internet Source

<1 %

33

[pubannotation.org](http://pubannotation.org)

Internet Source

<1 %

34

Submitted to sirgeorgemonouxcollege

Student Paper

<1 %

35

[www.iosrjournals.org](http://www.iosrjournals.org)

Internet Source

<1 %

36

Mohan Amarasiri, Daisuke Sano, Satoru Suzuki. "Understanding human health risks caused by antibiotic resistant bacteria (ARB) and antibiotic resistance genes (ARG) in water environments: Current knowledge and questions to be answered", Critical Reviews in Environmental Science and Technology, 2019

Publication

<1 %

37

V. L. Kung, E. A. Ozer, A. R. Hauser. "The Accessory Genome of *Pseudomonas aeruginosa*", *Microbiology and Molecular Biology Reviews*, 2010

Publication

<1 %

38

Farzaneh AMINHARATI, Mohammad Hassan EHRAMPOUSH, Mohammad Mehdi SOLTAN DALLAL, Mehdi YASERI et al. "Citrobacter freundii Foodborne Disease Outbreaks Related to Environmental Conditions in Yazd Province, Iran", *Iranian Journal of Public Health*, 2020

Publication

<1 %

39

Submitted to Sankalchand Patel University

Student Paper

<1 %

40

Submitted to Teerthanker Mahaveer University

Student Paper

<1 %

41

[stud.epsilon.slu.se](http://stud.epsilon.slu.se)

Internet Source

<1 %

42

Shrijana Bista, Bindeshwar Yadav, Gopiram Syangtan, Jivan Shakya, Reshma Tuladhar, Dev Raj Joshi, Binod Lekhak. " Emergence of IS-linked oxacillinase genes among carbapenem resistant isolates in a tertiary cardiac center, Nepal ", Cold Spring Harbor Laboratory, 2023

Publication

<1 %

43

[ci.lib.ncsu.edu](http://ci.lib.ncsu.edu)

Internet Source

&lt;1 %

44

[www.kjvr.org](http://www.kjvr.org)

Internet Source

&lt;1 %

45

Submitted to Jaypee University of Information  
Technology

Student Paper

&lt;1 %

46

Renáta Kolínská, Petra Španělová, Michal  
Dřevínek, Jaroslav Hrabák, Helena Žemličková.  
"Species identification of strains belonging to  
genus *Citrobacter* using the biochemical  
method and MALDI-TOF mass spectrometry",  
*Folia Microbiologica*, 2014

Publication

&lt;1 %

47

Verónica Elizabeth Álvarez, María Paula  
Quiroga, Gabriel Alejandro Castro, Marcelo  
Hernán Cassini, Daniela Centrón. "Chapter 7  
Molecular Ecology of Class 1 Integrons in  
Patagonia as Model System for  
Understanding the Rise of Antibiotic  
Resistance Isolates Around the World",  
Springer Science and Business Media LLC,  
2016

Publication

&lt;1 %

48

Fereshteh Shahcheraghi. "Molecular  
characterization of class 1 integrons in MDR  
*Pseudomonas aeruginosa* isolated from clinical

&lt;1 %

settings in Iran, Tehran", FEMS Immunology & Medical Microbiology, 12/2009

Publication

---

49

H W Stokes. "Gene cassettes encoding resistance to quaternary ammonium compounds: a role in the origin of clinical class 1 integrons?", The ISME Journal, 10/16/2008

Publication

---

50

[media.suub.uni-bremen.de](http://media.suub.uni-bremen.de)

Internet Source

---

51

[s3.eu-central-1.amazonaws.com](http://s3.eu-central-1.amazonaws.com)

Internet Source

---

52

Katakweba, Abdul, Karen Skou Møller, Justice Muumba, Amandus Muhairwa, Peter Damborg, Jesper Tjørnhøj Rosenkrantz, Uswege M. Minga, Madundo Mtambo, and John Elmerdahl Olsen. "Antimicrobial resistance in faecal samples from buffalo, wildebeest and zebra grazing together with and without cattle in Tanzania", Journal of Applied Microbiology, 2014.

Publication

---

53

Chang-Ho Kang, Yujin Shin, SeokCheol Jang, Yeojung Jung, Jae-Seong So. "Antimicrobial susceptibility of *Vibrio alginolyticus* isolated from oyster in Korea", Environmental Science and Pollution Research, 2016

<1 %

<1 %

<1 %

<1 %

<1 %

## Publication

---

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off