

www.berkalahayati.org

Accredited by Ministry of Research and Higher Education Number 32a/E/KPT/2017

SINTA 2

Berkala PENELITIAN HAYATI

Journal of Biological Researches

ISSN : 08526834
e-ISSN : 2337-389X

PBI
East Java

Volume 28 No. 1

December 2022

Doc: Visualization of 3D models mir-135b's siRNA © Valeska and Parikesit

THE EAST JAVA BIOLOGICAL SOCIETY

A H U - 0 0 1 1 5 6 4 . A H . 0 1 . 0 7 . Y E A R O F 2 0 1 7

The East Java Biological Society is a non-profit organization made up of biology graduates with a variety of interests. The East Java Biological Society is affiliated with several East Java universities, including collaborations for the publication of proceedings and seminars.

For further information about membership in The East Java Biological Society, please email perkumpulanbiologijatim@gmail.com.

The East Java Biological Society

Chairman	:	Prof. Dr. Bambang Irawan, M. Sc
Secretary	:	Irfan Mustafa, S.Si, M.Si, Ph.D
Treasurer	:	Didik Wahyudi, S.Si., M.Si

Editorial Board (Berkala Penelitian Hayati)

Editor in Chief	:	Prof. Sutiman Bambang Sumitro, SU., D.Sc.
Managing Editor	:	Irfan Mustafa, S.Si, M.Si, Ph.D Hadiatullah, S.Si, M.Si Wulida Khoirunnisa, S.Si, M.Si
Member	:	Dr. Akira Kikuchi Dr. Bagyo Yanuwiadi Prof. Bambang Irawan, M. Sc Prof. Dr. Endang Semiarti, M.S., M.Sc. Prof. Fatchiyah, Ph. D Dr. Hitoshi Sawada Prof. Intan Ahmad, Ph. D Prof. Widodo, S.Si., M.Si., Ph.D.Med.Sc.

AUTHOR GUIDELINES

Berkala Penelitian Hayati is a peer-reviewed, open-access journal published by Indonesia Biological Association, East Java, Indonesia. Botany, zoology, ecology, microbiology, physiology, nanobiology, coastal biology, hydrobiology, neurobiology, genetics, developmental biology, biochemistry and molecular biology, biophysics, and life science are the biological fields covered by the journal. Authors are encouraged to submit complete unpublished and original works that are not under review in other journals. Manuscripts should be written and will be published in English. Berkala Penelitian Hayati is currently distributed in several reputable libraries in Indonesia. Authors should refer to the instructions below to prepare their manuscripts. One of the editors and at least two reviewers will review the submissions. Authors may suggest names of potential reviewers. Berkala Penelitian Hayati uses anonymous reviewing, in which the identities of authors and reviewers are kept confidential.

Manuscript Submission

Authors need to register with the journal prior to submitting to <http://berkalahayati.org> or send by email berkalahayati@yahoo.com. (manuscript template can be downloaded on the website)

Type of Manuscript

Original Article presents an original and significant major advance in bioscience, having a wide research scheme and deep discussion of the findings. The article's structure consists of Abstract, Introduction, Methods, Results, Discussion, and References (see Manuscript Format in details). The article should be typed in paper length of 15 pages or 3000–4500 words.

Review Article is expected to present an original and review of significant advances in bioscience with a comprehensive research scheme and deep discussion of the findings. The article's structure consists of Abstract, Introduction, Contents, and References (see Manuscript Format in details). The article should be typed in the paper length of 15 pages or 3000–4500 words.

Short Communication is based on the urgency, research scope and/or depth of discussion, Berkala Penelitian Hayati will publish an article as a short communication. The short communication (~1500 words) receives the same in-depth review as the full article.

Manuscript Formatting Guide

- Please provide the manuscript with no authors identity.
- The manuscript is written in 12 point Times New Roman font double space in A4 paper. Margins on all four sides are 2 cm with left alignment and do not split it into double-columns (should be single-column).
- Page numbers are located at the right bottom.
- Use Line numbers. To add the line numbers in the Microsoft Word program, go to the "Layout" menu in the "Page Setup" main menu. Click Line Numbers, make a tick in "Add line numbering" and choose "Continuous" option.
- Figures and Tables are uploaded separately in another file. Provide figures in jpeg/jpg and compile in pdf format, while tables are provided and collected in Microsoft Excel.

Manuscript sections are written in this order i.e.:

Title, Abstract, Keywords, Introduction, Methods, Results, Discussion, Acknowledgement, References

Article Processing Charge

The author whose works is accepted for publication is subjected to pay 1,400,000 IDR (100 USD) per published article up to 7 pages.

Open-Access

Berkala Penelitian Hayati is an open-access journal: all articles will be permanently and immediately free for everyone to download. After acceptance, the articles will be published under a noncommercial licence that everyone free to share (copy and redistribute) the material in any medium or format by the Creative Commons Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0).

Originality, Acknowledgement and Plagiarism of Sources

Author should submit only entirely original works and experiments, also appropriately cite or quote the work and/or words of others. Publications that have been influential in determining the nature of the reported work should be cited.

Privacy Statement

The names and e-mail addresses entered in this journal site will be used exclusively for the stated purposes of this journal and will not be made available for any other purpose or to any other part.

T A B L E O F C O N T E N T

Solikin Solikin <i>Infestation and host specificity of mistletoe parasitic plants in Purwodadi Botanic Garden</i>	1-9
Budhi Setiawan, Nugroho Eko Wirawan Budianto, Masfufatun Masfufatun <i>Potential improvement in the safety and quality of traditional fermented soybean products: A narrative review</i>	10-19
Nyan Hayu Widiastuti, Bambang Retnoaji <i>Life-stage dependent histological characterization of liver HIF-1α expression in the Indonesian shortfin eel (<i>Anguilla bicolor bicolor</i> McClelland, 1884)</i>	20-25
Miftahul Jannah, Qurrota A`yun, Nida Afifah, Eko Prasetya, Muhammad Rifqi Hariri <i>Usnea in West Java: a potential source of bioactive secondary metabolites</i>	26-31
Hendra Susanto <i>Physiological alteration and the expression of fibrogenesis-proinflammatory genes in hepatocellular carcinoma patients with chronic hepatitis C and onset metabolic syndrome: a preliminary report at Saiful Anwar General Hospital Malang</i>	32-38
Tintrim Rahayu, Gatra Ervi Jayanti, Dita Agisimanto <i>Indole-3-butyric acid immediately induced adventitious root of <i>Dendrodium milla</i> nayla x <i>Dendrobium striaenopsis</i> planted on coco-hust and wood charcoal</i>	39-43

TABLE OF CONTENT

Pebraniafrahul I Siregar, Sri Pujiyanto, Arina Tri Lunggani <i>Inhibitory activity of endophytic fungi against alpha-amylase isolated from raru (Cotylelobium melanoxylon)</i>	44-50
Joko Pilianto, Xiayu Hu, Hadiatullah Hadiatullah, Tshenolo Tsheko, Asim Munawar, A.S. Abou El-Ela, Wen-Wu Zhou, Zeng-Rong Zhu <i>Consumer perception of insect-based foods in the community of Zhejiang University</i>	51-56
Arum Linangkung, Adeodatus Yuda Handaya, Hendra Susanto, Nurcahya Setyawan, Imam Sofii, Agus Barmawi <i>Expressions of ZEB1 of patients with colorectal cancer in Indonesia: Correlations with clinicopathological characteristics and liver metastasis profile</i>	57-61
Margareta Deidre Valeska, Arli Aditya Parikesit <i>Determination of 3D structure and molecular interaction for mir-135b and its silencer as Triple Negative Breast Cancer (TNBC) biomarkers</i>	62-66
I.B.G Darmayasa, I.W Suwanda, I.K Widnyana <i>Isolation and identification of lontar-degrading fungus in Bali island, Indonesia</i>	67-71
I Putu Agus Hendra Wibawa, Sutomo, I Nyoman Lugrayasa <i>Antioxidant assay of Averrhoa bilimbi L flower extract, chemical compound and its utilization potential</i>	72-77

Potential improvement in the safety and quality of traditional fermented soybean products: A narrative review

Budhi Setiawan¹, Nugroho Eko Wirawan Budiarto¹, Masfufatun²

¹Department of Pharmacology, Medical Faculty, University of Wijaya Kusuma Surabaya, Dukuh Kupang XXV/54 Surabaya 60225, Indonesia

²Department of Biochemistry, Medical Faculty, University of Wijaya Kusuma Surabaya, Dukuh Kupang XXV/54 Surabaya 60225, Indonesia

Abstract

Soybeans are one of the most significant crops in the world due to their nutritionally valuable. It has been converted to produce a variety of fermented soybean products. During conventional soybean fermentation, microorganisms are involved, biochemical reactions occur, and bioactive components are produced. With the rise of people's living standards, customers are paying more attention not only to the flavor and nutrition of fermented soybean meals, but also to their safety and quality. Mycotoxins, biogenic amine production, and high salt content are among the public health concerns associated with fermented soybean foods. This paper reviews the prevalent concerns about the safety and quality of fermented soybean foods, as well as potential improvements. Attempts and methods have been proposed to ensure the safety of the fermentation process and food quality. Official regulations, the employment of suitable microbes, the use of high-quality cultivars, and the administration of chemicals are all viable options for improving safety and quality. We conclude that implementing international food standards, guidelines, and codes of practice such as the Codex Alimentarius for fermented soybean products and the application of scientific novel methods (e.g., starter combination, high-pressure processing, or low-dose gamma irradiation, additive usage, low salt fermentation technique) are the potential solutions to mitigate the issues and improve the safety and quality of the products.

Keywords: Fermentation, soybean, safety, quality, food, nutrition

Received: September 25, 2021 Revised: April 09, 2022 Accepted: April 11, 2022

Introduction

Soybean (*Glycine max* [L.] Merrill) is a low-cost major protein source for the human diet that can be utilized as a meat substitute (Xiang et al., 2019). Soy-based foods are known as non-fermented (soymilk, okara, tofu, yuba, soy nuts, etc.) and fermented (miso, tempe, sufu, natto, etc.) (Liu, 2008). Several studies have reported that fermenting soybeans results in numerous beneficial properties. The fermentation procedure can shorten cooking time, improve the nutritional quality of soybeans, boost bioactive components such as phytoestrogens, and make soybeans more digestible (Chen et al., 2016). Some fermented soybean products (FSPs) are often used as condiments to stimulate the appetite and improve food flavor (Xiang et al., 2019).

It is widely known that anti-nutritional substances included in whole soybeans have a deleterious impact on nutrient and micronutrient absorption. Whole soybeans contain anti-nutritional factors such as trypsin inhibitor, lectins, phytic acid, and oligosaccharides, raising concern for their consumption (Thakur et al., 2019). Briefly, it has a negative effect on nutrient absorption and utilization, as well as vitamin degradation (Thakur et al., 2019). This impact can trigger nutrient deficiencies, which can lead to a variety of health-related issues, particularly in animals. The trypsin inhibitor activity in unfermented soybean has a negative impact on the

growth performance and pancreas weight of broiler chickens (Hoffman et al., 2019; Hemetsberger et al., 2021).

Studies have shown that microorganisms used in conventional fermentation are capable of producing toxins that contaminate FSPs (Chen et al., 2016). For example, *Rhizopus microspores*, a popular species employed as a soybean fermentation starter, can produce two types of mycotoxins, rhizoxins and rhizonins. These mycotoxins are present in tempe (an Indonesian fermented soybean cake) and fermented bean curd (a fermented soybean paste) (Rohm et al., 2010). *Aspergillus oryzae*, which is often employed in traditional soybean fermentation, also produces toxic secondary metabolites, such as cyclopiazonic acid, aspergillomarasmine, 3-nitropropionic acid, kojic acid, maltoryzine, and violacetin (Blumenthal, 2004).

Biogenic amines (BA), harmful substances generated during the fermentation process, are frequently identified in substantial concentrations in fermented soybean products from Korea, China, Japan, and Indonesia (Park et al., 2019). Certain fermented soybean products, such as fermented bean curd, soy sauce, douchi, and dajiang, are typically stored without being sterilized, but with a high salt content. This technique prevents spoilage and extends the shelf life of the food (Chen et al., 2016; Liu et al., 2020). However, an excessive intake of salt may result in an expansion in circulating volumes and lead to an increase in blood pressure in humans (Kommenov et al., 2019). As such, this narrative review will evaluate the issues associated with the safety and quality of commonly fermented soybean products and offer some potential solutions.

* Corresponding Author:
Budhi Setiawan
Department of Pharmacology, Medical Faculty, University of
Wijaya Kusuma Surabaya, Dukuh Kupang XXV/54 Surabaya
60225, Indonesia
Phone: +6287751644522 Fax: -
E-mail: budhisetiawan@uwks.ac.id

Inconsistency of quality improvement

Traditionally fermented soybean foods are typically prepared using traditional fermentation process. There are numerous methods for producing indigenous soy-based foods (Figure 1). Asian countries offer a variety of fermented soybean preparation products based on traditional customs (Table 1). Standard quality procedures are not followed properly throughout the typical fermentation process, resulting in varied outcomes for fermented soybean products. Numerous elements contribute to the consistency of quality for particular fermented soybean food. For instance, the quality of tempe (Indonesia’s fermented soybean cake)

might be influenced by soybeans, process water, yeast, fermentation time, and fermentation temperature (Novita & Abidin, 2020). In daily practice, fermentation is a complicated process, and the final product's quality is highly dependent on a variety of factors, including geographical location, the environment, weather, seasons, etc. (Liu et al., 2020). For instance, there is a possibility in Indonesia that the small-scale industry of fermented soybean business may fall short of achieving the government's national food quality and safety criteria due to environmental factors and raw materials (Anggriawan, 2018; Kadar et al., 2021).




Figure 1. Different methods of soybean fermentation process

Table 1. Countries’ distribution and variety of fermented soybean products

Country	Local name	Microorganism	References
India	Bekang	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. pumilus</i> , <i>Bacillus cereus</i> , <i>B. amyloliquefaciens</i>	Chettri & Tamang, (2014); Singh et al., (2014a, 2014b)
	Tungrymbai	<i>B. subtilis</i> , <i>B. pumilus</i> , <i>B. licheniformis</i> , <i>B. amyloliquefaciens</i> , <i>Lactobacillus brevis</i> ,	
Japan	Hawaijar	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. amyloliquefaciens</i> , <i>B. cereus</i> ,	Tamang et al., (2016)
	Miso	<i>Bacillus subtilis</i> , <i>Pediococcus acidilactici</i> , <i>Leuconostoc mesenteroides</i> , <i>Micrococcus halobius</i> , <i>Aspergillus oryzae</i> , <i>Zygosaccharomyces rouxii</i> and <i>Torulopsis sp</i>	
Indonesia	Natto	<i>Bacillus subtilis</i> , <i>Aspergillus oryzae</i>	Liu, (2012)
	Kecap	<i>Rhiz. oligosporus</i> , <i>Rhiz. oryzae</i> , <i>Asp. oryzae</i> , <i>Ped. halophilus</i> , <i>Staphylococcus sp.</i> , <i>Candida sp.</i> , <i>Debaromyces sp.</i> , <i>Sterigmatomyces sp.</i>	Alexandraki et al., (2013)
	Tauco	<i>Rhiz. oryzae</i> , <i>Rhiz. oligosporus</i> , <i>Asp. oryzae</i> , <i>Zygosaccharomyces soyae</i> , <i>Bacillus sp.</i> , <i>Ent. hermanniensis</i> , <i>Lb. agilis</i> , <i>Lb. brevis</i> , <i>Lb. buchneri</i> , <i>Lb. crispatus</i> , <i>Lb. curvatus</i> , <i>Lb. delbrueckii</i> , <i>Lb. farciminis</i> , <i>Lb. fermentum</i> , <i>Lb. pantheris</i> , <i>Lb. salivarius</i> , <i>Lb. vaccinostercus</i> , <i>Lc. lactis</i> , <i>Lactococcus sp.</i> , <i>Leuc. camosum</i> , <i>Leuc. citreum</i> , <i>Leuc. fallax</i> , <i>Leuc. lactis</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. pseudomesenteroides</i> , <i>Ped. acidilactici</i> , <i>Strep. bovis</i> , <i>Strep. macedonicus</i> , <i>W. cibaria</i> , <i>W. confusa</i> , <i>W. paramesenteroides</i> , <i>W. soli</i>	Kindossi et al., (2012); Winarno, (1973)
Korea	Chungkookjang	<i>Bacillus amyloliquefaciens</i> , <i>B. licheniformis</i> , <i>Bacillus megaterium</i> , and <i>B.subtilis</i> , <i>Leuconostoc</i> , <i>Lactobacillus</i> , <i>Pseudomonas</i> , <i>Pantoea</i> and <i>Weissella</i> genera	Hong et al. (2012)
	Doenjang	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. pumilus</i> , <i>Mucor plumbeus</i> ,	Chang et al., (2013); Kim et al.,

		<i>Aspergillus oryzae</i> , <i>Debaryomyces hansenii</i> , <i>Leuconostoc mesenteroides</i> , <i>Tor. halophilus</i> , <i>Lactobacillus sp.</i>	(2016)
	Gochujang	<i>B. velegensis</i> , <i>B. amyloliquefacious</i> , <i>B. subtilis</i> , <i>B. licheniformis</i> , <i>Oceanobacillus spp.</i> <i>Zygosaccharomyces</i> ,	Nam et al., (2012); Jang et al., (2019); Sanni et al., (1991)
	Meju	<i>Candida lactis</i> , <i>Zygorouxii</i> , <i>Aspergillus</i> , <i>Penicillium</i> , <i>Rhizopus Bacillus cereus</i> , <i>B. circulans</i> , <i>B. licheniformis</i> , <i>B. megaterium</i> , <i>B. mesentricus</i> , <i>B. subtilis</i> , <i>B. pumilus</i> , <i>Aspergillus spp.</i> , <i>Botrytis cinerea</i> <i>Rhizopus oryzae</i> , <i>Rhodotorula flava</i> , <i>Zygosaccharomyces japonicus</i> , <i>Lactobacillus sp.</i> , <i>Ped. acidilactici</i>	
	Kanjang	<i>Asp. oryzae</i> , <i>B. subtilis</i> , <i>B. pumillus</i> , <i>B. citreus</i> , <i>Sarcina mazima</i> , <i>Sacch. rouxii</i>	Shin, (2012)
China	Furu	<i>B. pumilus</i> , <i>B. megaterium</i> , <i>B. stearothermophilus</i> , <i>B. firmus</i> , <i>Staphylococcus hominis</i>	Lin et al., (2016)
	Yandou	<i>B. subtilis</i>	Qin, (2013); Ouoba, (2005); Rabie, (2009); Sanni, (2002)
Taiwan	Jiang-sun	<i>Lb. plantarum</i> , <i>Ent. faecium</i> , <i>Lc. lactis subsp. lactis</i>	Chen, (2010)
China, Taiwan	Douchi	<i>B. amyloliquefaciens</i> , <i>B. subtilis</i> ,	Yang et al., (2019a, 2019b)
	Meitauza	<i>B. subtilis</i> , <i>Asp. oryzae</i> , <i>Rhiz. oligosporus</i> , <i>Mu. meitauza</i> , <i>Actinomucor elegans</i>	Zhu et al., (2008)
India, Nepal, Bhutan	Kinema	<i>B. subtilis</i> , <i>B. licheniformis</i> , <i>B. cereus</i> , <i>B. circulans</i> , <i>B. thuringiensis</i> , <i>B. sphaericus</i>	Chettri et al., (2016); Chettri & Tamang, (2015); Singh et al., (2014a, 2014b)
Japan, Korea, China	Shoyu	<i>Asp. oryzae</i> or <i>Asp. sojae</i> , <i>Z. rouxii</i> , <i>C. versatilis</i>	Sugawara, (2010).
Indonesia (Origin), Netherlands, Japan, USA	Tempe	<i>Bacillus pumilus</i> and <i>B. brevis</i> , <i>Rhizopus spp.</i> , <i>Lactobacillus casei</i>	Frias et al., (2017)

Regulatory implications

Despite fermented soybean products are generally considered safe, there are still health risks associated with improperly fermented soybean products, which are produced under inadequate hygienic standards and uncontrolled conditions, especially in household industry settings. It has been claimed that traditional fermented soybean products marketed in Northeast India are contaminated with pathogens (*Bacillus cereus*, *Clostridium botulinum*, *Escherichia coli*, *Proteus mirabilis*) (Keisam et al., 2019). To improve hygienic standards and safeguard community health, various food safety assurance systems should be implemented, such as education and information, starting culture utilization, high-quality raw materials, and ideal fermentation conditions (Tamang et al., 2020). Standardization of the manufacturing process is necessary to provide consistent quality, particularly in small-scale industry settings (Anal, 2019).

There are a variety of certification schemes available, including Good Manufacturing Practices (GMP), Good Agricultural Practices (GAP), Good Hygienic Practices (GHP), Hazard Analysis Critical Control Points (HACCP), and International Organization for Standardization (ISO) standards. One or more of these certifications should be implemented and supported by government policies in order to eliminate public health concerns associated with fermented soy products, particularly in small scale industrial settings. Having Codex Alimentarius registration for fermented soybean products, such as kimchi, gochujang, and doenjang, will improve the quality standard, particularly in international trade and commerce (Lee et al., 2012). Food safety and quality audits are also commonly used to ensure the compliance and effectiveness of food safety initiatives. Audits are commonly used to evaluate management systems, check the state of premises and products, and certify that they meet specified food safety and quality requirements (Kotsanopoulos & Arvanitoyannis, 2017).

There are specific fermented soybean product standards that can be followed, such as the Codex Alimentarius regional standard for fermented soybean paste (Asia) CODEX STAN 298R-2009 or Indonesian national standard SNI 3144:2009 for tempe. However, the goal of food safety and quality audits should be one of continuous development rather than one-time evaluation results or policing (Bradford-Knox, 2017).

Anti-nutritional factor

Numerous processing techniques, such as soaking, pressure cooking, blanching, germination, fermentation, autoclaving, genetic editing, and other methods, all assist in reducing the anti-nutrient effects (Thakur et al., 2019; ten Brink et al., 1990). It is feasible to reduce the antinutrient content of meals by utilizing any of these varied strategies alone or in combination (ten Brink et al., 1990). It has been suggested that autoclaving may be a more effective method than other processing methods for reducing levels of several anti-nutritional components (Santiya et al., 2020). However, further study is still required to develop strategies for removing heat-stable anti-nutritional components found in a variety of foods without impairing their nutritional content. Phytic acid, tannins, alkaloids, saponins, and non-protein amino acids are examples of heat-stable antinutritional factors, whereas lectins, cynogenic glycosides, and protease inhibitors are examples of heat-labile antinutritional factors. These substances are found in a variety of grain legumes, including soybeans (Thakur et al., 2019). When present in trace amounts, several anti-nutritional agents and their metabolites may have beneficial health benefits such as decreased blood glucose and lipid levels, as well as a decreased risk of cancer. In contrast, a lack of understanding and information about their properties can lead to health issues (Gemedé & Ratta, 2014).

Innovative fermentation techniques with specific microbes have been used to lower the level of anti-nutritional elements in fermented soybean products, such

as soybean meal. Fermentation of soybean meal protein with *Bacillus siamensis* isolate JL8 resulted in a significant reduction of major antinutritional factors (ANFs) in soybean meal (SBM), specifically glycinin, beta-conglycinin, and trypsin inhibitor, which were reduced by 86.0%, 70.3%, and 95.01%, respectively (Zheng *et al.*, 2017). After 24 hours, a comparable study using solid-state *Bacillus subtilis* fermentation was able to lower the quantities of beta-conglycinin subunits, glycinin subunits, and trypsin inhibitors in soybean meal by 70%, 50%, and 58%, respectively (Seo & Cho, 2016). Additionally, soybean meal fermentation at 35 °C with an appropriate proportion of *Bacillus subtilis*, *Bacillus lactis*, and *Saccharomyces cerevisiae* might increase the levels of free amino acids and short peptides while decreasing the activity of protease inhibitors (Zhang *et al.*, 2017).

Mycotoxin contamination

The common microorganisms involved in FSPs, especially in the traditional fermentation process, have been described (Table 1). Mycotoxins are secondary metabolites of molds that may be harmful to humans or animals when consumed and may also have unfavorable effects on crops (Zain, 2011). When mycotoxins are inhaled, they can induce acute disease or persistent illness episodes known as mycotoxicosis. Due to mycotoxin's extreme toxicity, exposure should be limited to a minimum. Different countries have set their own maximum levels of aflatoxins in food for humans, which range from 5 to 50 g/kg (Wu, 2004) and 4 to 5 g/kg for beans (Wu, 2004) and 4 to 5 µg/kg for beans (Adejumo & Adejoro, 2014). The term "mycotoxin" originates from the Greek word "*mykes*", which means fungus, and the Latin word "*toxicum*" which means poison or toxin (Kirovska & Velickova, 2021). Mycotoxin is produced during fungal growth and is detected in the hyphae and spores of organisms. Numerous mycotoxins have been detected in fermented food products, including aflatoxins, ochratoxins, deoxynivalenol (DON), trichothecenes, zearalenone, and citrinin (Omotayo *et al.*, 2019; Sivamaruthi *et al.*, 2019). Although aflatoxin contamination is more prevalent in crops such as peanuts and maize, a prior study discovered that soybean seeds contained more than 20 aflatoxins (Roy *et al.*, 2001). Some trichothecene mycotoxins, such as DON and T-2 toxins (a trichothecene produced by *Fusarium* spp.) can cause direct damage to mucosal tissues by disrupting the intestinal epithelial barrier, hence facilitating translocation of intestinal commensal microbiota, and pathogens. As a result, microorganism infection and inflammation may be increased (Nazhand *et al.*, 2020). Aflatoxin (AF) is a prevalent mycotoxin produced by *Aspergillus flavus*, *A. parasiticus* and *A. nomius*. Aflatoxins are formed chemically from difuranocoumarins with bifuran-based coumarin nuclei groups and lactone rings (Aflatoxin Gs) or pentanone rings (Aflatoxin Bs and Aflatoxin Ms). During the biosynthesis of AF in plants by *A. flavus* and *A. parasiticus*, the main substrate of hexanoyl is converted to polyketide via a polyketide synthase and two fatty acid synthases (Nazhand *et al.*, 2020).

It has been reported that 97.5% of tested samples of 40 domestic and imported (from Japan) soy sauces sold in China were contaminated with DON. Domestic soy sauces had a mean incidence rate of DON contamination of 97.1%, with concentrations ranging from 4.5 to 1245.6 g/l, whereas those imported from Japan had a mean incidence rate of 100%, with concentrations ranging from 30.5-238.3 g/l (Zhao *et al.*, 2013). Meju prepared without additives may result in the isolation of *Aspergillus ruber*, which releases aflatoxin B₁ and ochratoxin A (Shukla *et al.*, 2017). *Aspergillus oryzae* and *Aspergillus sojae* are frequently employed in the fermentation of soybeans to produce soy sauce and miso. These strains are closely related to the aflatoxigenic fungi *Aspergillus flavus* and *Aspergillus parasiticus*. While it has been demonstrated that these fermented soybean products do not create aflatoxin, they may contain homologs of numerous aflatoxin production pathway genes (Shukla *et al.*, 2017). Aflatoxin is frequently found in doenjang, which is made through inoculation with a natural starter. The aflatoxigenic *Aspergillus flavus* is identified as the fungal community present throughout conventional manufacturing (Zain, 2011).

Mycotoxin mitigation

Official regulation and policy are critical in reducing mycotoxin levels in foods in many countries. In 2003, the FAO reported that at least 99 nations regulated mycotoxins in food and/or feed. The number of countries has increased by around 30% compared to 1995. These countries collectively account for around 87% of the world's population ("Worldwide Regulations for Mycotoxins in Food and Feed in 2003" n.d.). Additionally, novel approaches for reducing mycotoxin levels in fermented soybean products should be studied. For instance, fermentation of Meju with plant extracts (*Nelumbo nucifera* leaves, *Ginkgo biloba* leaves, and *Allium sativum* cloves) has a substantial effect on the fungal microflora, resulting in an enhancement in the product's quality. This outcome could be explained by the extracts' diminished activity against toxin-producing fungal infections (Shukla *et al.*, 2017).

Biogenic amines

Biogenic amines (Bas) are low molecular weight, non-volatile nitrogenous bases with an aliphatic, aromatic, or heterocyclic structure that have been identified in fermented foods, such as miso, chunjang, jajang, and sufu (Komprda *et al.*, 2007; Pachlová *et al.*, 2012; Renes *et al.* 2014; Alvarez & Moreno-Arribas 2014; Bai *et al.*, 2013; Lee *et al.*, 2019; Shukla *et al.*, 2010; Guan *et al.*, 2013; Byun & Mah, 2012). The biogenic amine is named for the amino acid from which it is produced. For instance, histamine is created when histidine is decarboxylated, while tyramine is formed when tyrosine is decarboxylated (Santos, 1996). Biogenic amines are formed in foods as a result of the presence of bacteria capable of decarboxylation of specific amino acids (Renes *et al.*, 2014; Bover-Cid *et al.*, 2003; Buňková *et al.*, 2013). The bacteria with amino acid-decarboxylase activity are *Pseudomonas*,

Clostridium, *Bacillus*, *Photobacterium*, *Enterobacteriaceae* family (*Escherichia*, *Klebsiella*, *Citrobacter*, *Proteus*, *Shigella*, and *Salmonella*), *Micrococcaceae* family (*Staphylococcus*, and *Micrococcus*), *Enterococcus*, *Lactobacillus*, *Carnobacterium*, *Pediococcus*, *Lactococcus*, and *Leuconostoc* (Santos, 1996; Stadnik & Dolatowski, 2010).

BA is generally not harmful to human health in small doses since it can be detoxified by amine oxidase enzymes found in the intestine. The recommended BA values include a maximum of 100 mg/kg histamine in food and 2 mg histamine per liter of alcoholic beverage, a maximum of 100 to 800 mg/kg tyramine in food, and a hazardous dose of 30 mg/kg β -phenylethylamine in food (ten Brink, 1990). Enzymes, such as monoamine oxidase (MAO), diamine oxidase (DAO), and histamine N-methyltransferase (HNMT), can metabolize dietary BA in a healthy human being (Fogel et al., 2007). However, BA can turn into toxic metabolites that might threaten human health when consumed in excess or when humans' detoxifying capacity is limited or damaged (Anderson, 2008; Ruiz-Capillas & Herrero, 2019). Because accumulation of biogenic amines is possible in the presence of uncontrolled microbial enzymatic reactions (Halász et al., 1994), the presence of BA in food is always undesirable, as it can cause headaches, respiratory distress, palpitation, hypo or hypertension, and allergic reactions when absorbed at high concentrations (Restuccia et al., 2015).

Biogenic amine mitigation

The formation of BA in fermented soybean products is a complex process that is influenced by a number of parameters, including the BA-producing strain or species, salt concentration, and high temperatures (Bai et al. 2013; Chun et al. 2020). Salt concentrations may have an effect on the production of BA. This occurs as a result of elevated salt levels, which restrict the growth of microorganisms that make biogenic amino acids, resulting in a decrease in the production of decarboxylase enzymes (Chun et al., 2020; Chin & Koehler, 1986). For example, during fermentation, the amounts of key biogenic amines such as cadaverine and putrescine that were detected from doenjang were higher at 9% salt than at 12%, 15%, and 18% salt (Chun et al. 2020). Thermal breakdown of BA has been demonstrated to occur under high-temperature treatment, such as boiling in Chunjang (black soybean paste). After 10 min of frying at 200 °C, tryptamine and β -phenylethylamine levels are approximately 40% lower. While other BAs, such as cadaverine, histamine, tyramine, spermidine, and spermine, exhibit declines of approximately 10% to 20% (Bai et al., 2013).

Currently, the majority of efforts to develop worldwide regulatory standards for biogenic amines are concentrated on fish and seafood products rather than fermented soybean foods (Park, Lee & Mah, 2019). The suggested toxic limit of BA content in fermented soybean products is 1000 mg/kg (Santos 1996), while the indicated harmful limit of β -phenylethylamine is 30 mg/kg, 100 mg/kg for histamine, and 100 mg/kg for

tyramine (ten Brink et al., 1990). Attempts have been made to minimize the formation of BA in foods through sanitary handling and processing, the use of biogenic amine-degrading starter cultures (e.g. *B. subtilis* and *B. amyloliquefaciens* strains), the addition of some probiotic bacterial strains alone or in combination with the starter cultures, high-pressure processing, and low-dose gamma irradiation (Oh et al., 2014; Kim et al., 2005a; Mendes et al., 2005; Zhang et al., 2013). High pressures of 400 MPa for 5 min or 400 MPa for 5 min + 50 MPa for 72 h were able to considerably reduce BA levels to levels comparable to or less than control (Novella-Rodríguez et al., 2002). Gamma-irradiation of raw materials at dosages of 5, 10, and 15 kGy considerably decreased the amounts of histamine, putrescine, tryptamine, and spermidine by approximately 20–50% in the final fermented soybean products (Kim et al., 2003).

To reduce the concentration of BA in fermented soybean foods, one typical method is to employ a starting culture containing bacteria with lower decarboxylase activity (Alvarez & Moreno-Arribas, 2014). The application of *Bacillus subtilis* and *Bacillus amyloliquefaciens* strains can reduce histamine levels in vitro by 27–46% and 70%, and tyramine contents by 42–59% and 71%, respectively (Kim et al., 2012). Additionally, the use of *Lactobacillus plantarum* as a starter culture can reduce histamine and total BA content by 58–100% and 27%, respectively (Lee et al., 2016; Kung et al., 2017). Apart from using BA-deficient bacteria for fermentation, the use of additives has been recommended as an alternative strategy. When ethanol is added to the dressing mixture (red kojic rice, alcoholic beverage, salt, sugar, bean paste, and spices) during sufu fermentation, the BA content reduction decreases. This could be because ethanol inhibits the decomposition of water-soluble proteins during the ripening fermentation process (Qiu et al., 2018). Another addition, such as nicotinic acid, may inhibit the activity of *Enterococcus faecium* isolated from cheonggukjang's tyrosine decarboxylase. It implies that nicotinic acid could be employed as a tyrosine decarboxylase inhibitor to reduce tyramine content in vitro and in situ (Kang et al., 2018). Additionally, the application of gamma irradiation may help lower not just salt consumption but also biogenic amine levels during soybean fermentation (Kim et al., 2005). Though numerous strategies for reducing BA concentration in vitro and in situ have been published, practical implementation is still necessary to ensure the safety of fermented food items (Park et al., 2019).

High salt content

Numerous traditional fermented soybean products are manufactured utilizing salt-intensive processes. Sufu, aving, miso, dajiang, doenjang, and soy sauce are examples of traditional salt-fermented soybean foods; natto, tempe, and kinema are examples of non-salt-fermented soybean products. In general, sufu has a salt content of 5 to 11%, doenjang has a salt concentration of 9 to 18%, soy sauce has a salt concentration of more than 18%, douchi has a salt concentration of 10 to 15%, and dajiang has a salt concentration of more than 10% (Han

et al., 2004; Jeong et al., 2014; Chun et al., 2019; Liu et al., 2020). Due to the fact that salt is used to regulate the content of water, it has an effect on microbial growth and fermentation pace. Additionally, salt enhances the sensory quality of fermented soybean products and prolongs their shelf life (Han et al., 2003; Jeon et al., 2016; Shim et al., 2016; Liu et al., 2020).

However, excessive salt consumption can be detrimental to human health and the majority of epidemiological research has demonstrated that salt consumption is dose-dependently linked with blood pressure in participants with normotension and hypertension (Mohan et al., 2009; Graudal et al., 2019). Hypertension is a significant risk factor for heart disease and stroke (He et al., 2009; Komnenov et al., 2019). According to a previous study, hypertension is responsible for 62% of strokes and 49% of coronary heart disease (He et al., 2010). Increased sodium intake has also been linked to a number of adverse health effects, including worsening chronic kidney disease (Borrelli et al., 2021), an increased risk of gastric cancer (Tsugane et al., 2004; Peleteiro et al. 2011; D'Elia et al., 2014), decreased bone density (Teucher et al., 2008; Caudarella et al., 2009) and possible obesity (He et al. 2009; Grimes et al., 2013; Ma et al., 2015). According to the guidelines issued by the World Health Organization, adults should consume less than 2000 mg of sodium or 5 g of salt every day (Härtl 2013). Several organizations, including the European Society of Hypertension, the European Society of Cardiology, the American Heart Association, and the American Academy of Physician Assistants, recommend reducing salt consumption (Mancia et al., 2013; Welton et al., 2018). Salt-fermented soybean products are a substantial source of sodium in various nations, including Japan and Korea, and their consumption is positively connected with daily salt consumption (Anderson et al., 2010; Jeong et al., 2014).

A major difficulty in reducing the use of NaCl content in meals has been observed to be a loss in consumer approval (Liem et al., 2011). Efforts have been made to lower sodium concentration while maintaining the physicochemical features and sensory quality of fermented soybean products (Park et al. 2015; Lin et al., 2021). Several novel methods have been developed to produce low salt, or no salt fermented soybean products, such as natto, tempe, and kinema (Wang et al., 2007). In terms of physical strategies, the combination of high pressure and ultrasonic technology appears promising; chemical strategies include the right use of salt substitutes, flavor enhancers, and quality improvers. Biological techniques include functional microbial augmentation and microbial enzymatic technology. Cell fusion techniques and genetic engineering technology could be used in the future to improve the quality and safety of low-sodium fermented meals (Lin et al., 2021).

Potential quality improvement through soybean cultivar selection

Food-grade soybeans are the raw material for fermented and non-fermented soy food products, including tofu, soymilk, miso, natto, soy sprouts, and tempeh. Food-grade soybeans contain greater protein

concentrations than feed-grade soybeans, ranging from 40 to 45% (Jegadeesan et al., 2020). Aside from high protein content, most food-grade cultivars share common characteristics such as low oil content, high seed quality with no mottling or cracking, and rapid water uptake, because most soy food products require a soaking step, thus rapid water uptake is essential (Mullin et al., 2001; Zhang et al., 2010; Anderson et al., 2019). Aside from these characteristics, advancements in soybean breeding and cropping systems provide not only improved chemical composition but also yield preservation through weed, pathogen, insect pest, and abiotic stress resistance (Anderson et al., 2019). The use of cadmium (Cd) safe soybean cultivars for fermentation may help to minimize unwanted metal deposition in the human body caused by Cd-contaminated soil (Zhi et al., 2020; 2015). Accumulation of Cd can occur in soybeans when they are grown on Cd-polluted soil (Arao et al. 2003; Sugiyama et al., 2011). A specific soybean cultivar may be more suitable for various fermented soybean products; for example, the Saedanbaek cultivar delivers higher quality standards than other cultivars (Jinpong, Daepung 2, Pyeongwon, Cheonga, and Saekolkong) during the soy paste fermentation process (Shin et al., 2019). This cultivar has demonstrated the ability to produce greater amylase and protease activities, viscous material amounts, and amino-type nitrogen content (Shin et al., 2019).

Conclusion

Fermented soybean products (FSPs) have piqued the interest of scientists and consumers due to their nutritional values, bioactive components, and flavor profiles and texture. Despite the fact that FSPs are considered safe foods, with the establishment of soybean fermented product standards, people have grown more conscious of the possibility of food-borne illness. Improving the safety and quality of fermented soybean products is critical in order to avoid future public health issues. Antinutritional factors, pathogenic microorganisms and mycotoxins, the formation of biogenic amines, the use of excessive salt concentrations during the fermentation process, and other safety dangers are still major concerns in fermented soybean production. The following characteristics are critical and should be considered in order to promote nutritional quality and safe fermented soybean foods. First and foremost, food regulation policies and standards must be enforced, particularly on industrial soybean fermentation products. Second, using appropriate microorganism strains in a controlled setting will result in a superior conventional soybean fermentation process.

Acknowledgement

The Institute for Research and Community Service of Wijaya Kusuma University, Surabaya, Indonesia, has supported this paper through an internal grant scheme (No. 22/LPPM/UWKS/2021).

References

- Adejumo, T.O., Adejoro, D.O. (2014). Incidence of aflatoxins, fumonisins, trichothecenes and ochratoxins in Nigerian foods and possible intervention strategies. *Food Science and Quality Management*, 31, 127-47.
- Alexandraki, V., Tsakalidou, E., Papadimitriou, K., Holzapfel, W.H. (2013). Status and Trends of the Conservation and Sustainable Use of Microorganisms in Food Processes. *Commission on Genetic Resources for Food and Agriculture*, FAO Background Study Paper No. 65.
- Alvarez, M.A., Moreno-Arribas, M.V. (2014). The Problem of Biogenic Amines in Fermented Foods and the Use of Potential Biogenic Amine-Degrading Microorganisms as a Solution. *Trends in Food Science & Technology*, 39(2), 146-55.
- Anal, A.K. (2019). Quality Ingredients and Safety Concerns for Traditional Fermented Foods and Beverages from Asia: A Review. *Fermentation*, 5(1), 8.
- Anderson, A.K. (2008). Biogenic and Volatile Amine-Related Qualities of Three Popular Fish Species Sold at Kuwait Fish Markets. *Food chemistry*, 107(2), 761-767.
- Anderson, C.A., Appel, L.J., Okuda, N., Brown, I.J., Chan, Q., Zhao, L., Ueshima, H., Kesteloot, H., Miura, K., Curb, J.D., Yoshita, K. (2010). Dietary sources of sodium in China, Japan, the United Kingdom, and the United States, women and men aged 40 to 59 years: the INTERMAP study. *Journal of the American Dietetic Association*, 110(5), 736-745.
- Anderson, E.J., Ali, M.L., Beavis, W.D., Chen, P., Clemente, T.E., Diers, B.W., Graef, G.L., Grassini, P., Hyten, D.L., McHale, L.K., Nelson, R.L. (2019). Soybean [*Glycine Max* (L.) Merr.] Breeding: History, Improvement, Production and Future Opportunities. In *Advances in Plant Breeding Strategies: Legumes: Volume 7*, edited by Jameel M. Al-Khayri, Shri Mohan Jain, and Dennis V. Johnson, 431-516. Cham: Springer International Publishing.
- Anggriawan, R. (2018). Microbiological and Food Safety Aspects of Tempeh Production in Indonesia [Dissertation]. Göttingen, Germany: Georg-August-University Göttingen, Germany
- Arao, T., Ae, N., Sugiyama, M., Takahashi, M. (2003). Genotypic differences in cadmium uptake and distribution in soybeans. *Plant and Soil*, 251(2), 247-53.
- Bai, X., Byun, B.Y., Mah, J.H. (2013). Formation and Destruction of Biogenic Amines in Chunjang (a Black Soybean Paste) and Jajang (a Black Soybean Sauce). *Food Chemistry*, 141(2), 1026-31.
- Blumenthal, C.Z. (2004). Production of Toxic Metabolites in *Aspergillus Niger*, *Aspergillus Oryzae*, and *Trichoderma Reesei*: Justification of Mycotoxin Testing in Food Grade Enzyme Preparations Derived from the Three Fungi. *Regulatory Toxicology and Pharmacology*, 39(2), 214-28.
- Bover-Cid, S., Hernández-Jover, T., Miguélez-Arrizado, M.J., Vidal-Carou, M.C. (2003). Contribution of Contaminant Enterobacteria and Lactic Acid Bacteria to Biogenic Amine Accumulation in Spontaneous Fermentation of Pork Sausages. *European Food Research and Technology*, 216(6), 477-82.
- Bradford-Knox R. (2017). Approaches to and the Management of the Audit Process in the Food Industry. *British Food Journal*, 119 (4), 759-70.
- Borrelli, S., Provenzano, M., Gagliardi, I., Ashour, M., Liberti, M.E., De Nicola, L., Conte, G., Garofalo, C., Andreucci, M. (2020). Sodium intake and chronic kidney disease. *International Journal of Molecular Sciences*, 21(13), 4744.
- Buňková, L., Adamcová, G., Hudcová, K., Velichová, H., Pachlová, V., Lorencová, E., Buňka, F. (2013). Monitoring of Biogenic Amines in Cheeses Manufactured at Small-Scale Farms and in Fermented Dairy Products in the Czech Republic. *Food Chemistry*, 141(1), 548-51.
- Byun, B.Y., Mah, J.H. (2012). Occurrence of Biogenic Amines in Miso, Japanese Traditional Fermented Soybean Paste. *Journal of Food Science*, 77(12), T216-23.
- Caudarella, R., Vescini, F., Rizzoli, E., Francucci, C.M. (2009). Salt intake, hypertension, and osteoporosis. *Journal of endocrinological investigation*, 32(4 Suppl), 15-20.
- Chang, M.I., Kim, J.Y., Kim, U.S., Baek, S.H. (2013). Antioxidant, tyrosinase inhibitory, and anti-proliferative activities of gochujang added with cheonggukjang powder made from sword bean. *Korean Journal of Food Science and Technology*, 45(2), 221-226.
- Chen, F., Zhang, X., and Li, Z. (2016). Chapter 15 – Safety of Fermented Products Based on Soybean. In *Regulating Safety of Traditional and Ethnic Foods*, edited by Vishweshwariah Prakash, Olga Martín-Belloso, Larry Keener, Siân Astley, Susanne Braun, Helena McMahon, and Huub Lelieveld, 311-18. San Diego: Academic Press.
- Chen, Y.S., Wu, H.C., Liu, C.H., Chen, H.C., Yanagida, F. (2010). Isolation and characterization of lactic acid bacteria from jiang-sun (fermented bamboo shoots), a traditional fermented food in Taiwan. *Journal of the Science of Food and Agriculture*, 90(12), 1977-1982.
- Chettri, R., Tamang, J.P. (2014). Functional properties of tungrymbai and bekaang, naturally fermented soybean foods of North East India. *International Journal of Fermented Foods*, 3(1), 87-103.
- Chettri, R., Tamang, J.P. (2015). Bacillus species isolated from tungrymbai and bekaang, naturally fermented soybean foods of India. *International Journal of Food Microbiology* 197:72-76.
- Chettri, R., Bhutia, M.O., Tamang, J.P. (2016). Poly-γ-glutamic acid (PGA)-producing bacillus species isolated from kinema, Indian fermented soybean food. *Front Microbiol Frontiers*, 7, 971.
- Chin, K.D., Koehler, P.E. (1986). Effect of Salt Concentration and Incubation Temperature on Formation of Histamine, Phenethylamine, Tryptamine and Tyramine During Miso Fermentation. *Journal of Food Protection*, 49(6), 423-27.
- Chun, B.H., Kim, K.H., Jeong, S.E., Jeon, C.O. (2020). The Effect of Salt Concentrations on the Fermentation of Doenjang, a Traditional Korean Fermented Soybean Paste. *Food Microbiology*, 86(April), 103329.
- D'Elia, L., Galletti, F., Strazzullo, P. (2014). Dietary salt intake and risk of gastric cancer. *Advances in Nutrition and Cancer*, 83-95.
- Fogel, W.A., Lewinski, A. (2007). Histamine in Food: Is There Anything to Worry About? *Biochemical Society Transactions*, 35(Pt 2), 349-52.
- Frias, J., Penas, E., Martinez-Villaluenga, C. (2017). Chapter 16 Fermented pulses in nutrition and health promotion. In: *Fermented foods in health and disease prevention*, 385-416
- Gemedé, H.F., Ratta, N. (2014). Antinutritional Factors in Plant Foods: Potential Health Benefits and Adverse Effects. *International Journal of Nutrition and Food Sciences* 3(4), 284.
- Guan, R.F., Liu, Z.F., Zhang, J.J., Wei, Y.X., Wahab, S., Liu, D.H., Ye, X.Q. (2013). Investigation of Biogenic Amines in Sufu (Furu): A Chinese Traditional Fermented Soybean Food Product. *Food Control*, 31(2), 345-52.
- Graudal, N., Hubeck-Graudal, T., Jürgens, G., Taylor, R.S. (2019). Dose-response relation between dietary sodium and blood pressure: a meta-regression analysis of 133 randomized controlled trials. *The American journal of clinical nutrition*, 109(5), 1273-1278.
- Grimes, C.A., Wright, J.D., Liu, K., Nowson, C.A., Loria, C.M. (2013). Dietary sodium intake is associated with total fluid and sugar-sweetened beverage consumption in US children and adolescents aged 2-18 y: NHANES 2005-2008. *The American Journal of Clinical Nutrition*, 98(1), 189-96.
- Halász, A., Barath, A., Simon-Sarkadi, L., Holzapfel, W. (1994). Biogenic Amines and Their Production by Microorganisms in Food. *Trends in Food Science & Technology*, 5(2), 42-49.
- Han, B.Z., Wang, J.H., Rombouts, F.M., Nout, M.R. 2003. Effect of NaCl on Textural Changes and Protein and Lipid Degradation during the Ripening Stage of Sufu, a Chinese Fermented Soybean Food. *Journal of the Science of Food and Agriculture*, 83(9), 899-904.
- Han, B.Z., Cao, C.F., Rombouts, F.M., Nout, M.R. (2004). Microbial changes during the production of Sufu-a Chinese fermented soybean food. *Food Control*, 15(4), 265-270.
- Härtl, G. (2013). WHO issues new guidance on dietary salt and potassium. *Central European Journal of Public Health* 21:16.
- Hemetsberger, F., Hauser, T., Domig, K.J., Kneifel, W., Schedle, K. (2021). Interaction of Soybean Varieties and Heat Treatments and Its Effect on Growth Performance and Nutrient Digestibility in Broiler Chickens. *Animals*, 11(9), 2668.
- Hoffmann, Thurner, S., Ankerst, D., Damme, K., Windisch, W., Brügger, D. (2019). Chickens' Growth Performance and Pancreas Development Exposed to Soy Cake Varying in

- Trypsin Inhibitor Activity, Heat-Degraded Lysine Concentration, and Protein Solubility in Potassium Hydroxide. *Poultry Science*, 98(6), 2489-2499.
- He, F.J., MacGregor, G.A. (2009). A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *Journal of Human Hypertension* 23(6), 363-384.
- He, F.J., MacGregor, G.A. (2010). Reducing population salt intake worldwide: from evidence to implementation. *Progress in Cardiovascular Diseases*, 52(5): 363-382.
- Hong, S.W., Choi, J.Y., Chung, K.S. (2012). Culture-based and denaturing gradient gel electrophoresis analysis of the bacterial community from Chungkookjang, a traditional Korean fermented soybean food. *Journal of Food Science*, 77, 572-578.
- Jang, M., Jeong, D.W., Lee, J.H. (2019). Identification of the predominant Bacillus, Enterococcus, and Staphylococcus species in meju, a spontaneously fermented soybean product. *Microbiology and Biotechnology Letters*, 47(3), 359-63.
- Jegadeesan, S., Yu, K. (2020). Food Grade Soybean Breeding, Current Status and Future Directions. *Legume Crops: Prospects, Production and Uses*, 27, 51.
- Jeon, H.H., Jung, J.Y., Chun, B.H., Kim, M.D., Baek, S.Y., Moon, J.Y., Yeo, S.H., Jeon, C.O. (2016). Screening and Characterization of Potential Bacillus Starter Cultures for Fermenting Low-Salt Soybean Paste (Doenjang). *Journal of Microbiology and Biotechnology*, 26(4), 666-74.
- Jeong, D.W., Kim, H.R., Jung, G., Han, S., Kim, C.T., Lee, J.H. (2014). Bacterial community migration in the ripening of doenjang, a traditional Korean fermented soybean food. *Journal of microbiology and biotechnology*, 24(5), 648-660.
- Jeong, Y.S., Lim, H.J., Kim, S.B., Kim, H.J., Son, S.M. (2014). Blood pressure and dietary related risk factors associated with high sodium intake assessed with 24-hour urine analysis for Korean adults. *Korean Journal of Community Nutrition*, 19(6), 537-549.
- Kadar, A.D., Astawan, M., Putri, S.P., Fukusaki, E. (2020). Metabolomics-Based Study of the Effect of Raw Materials to the End Product of Tempe – An Indonesian Fermented Soybean. *Metabolites*, 10(9), 367.
- Kang, H.R., Kim, H.S., Mah, J.H., Kim, Y.W., Hwang, H.J. (2018). Tyramine Reduction by Tyrosine Decarboxylase Inhibitor in Enterococcus Faecium for Tyramine Controlled Cheonggukjang. *Food Science and Biotechnology*, 27(1), 87-93.
- Keisam, S., Tuikhar, N., Ahmed, G., Jeyaram, K. (2019). Toxicogenic and pathogenic potential of enteric bacterial pathogens prevalent in the traditional fermented foods marketed in the Northeast region of India. *International Journal of Food Microbiology*, 296, 21-30.
- Kim, J.H., Ahn, H.J., Kim, D.H., Jo, C., Yook, H.S., Park, H.J., Byun, M.W. (2003). Irradiation effects on biogenic amines in Korean fermented soybean paste during fermentation. *Journal of Food Science*, 68(1), 80-4.
- Kim, J.H., Ahn, H.J., Lee, J.W., Park, H.J., Ryu, G.H., Kang, I.J., Byun, M.W. (2005). Effects of Gamma Irradiation on the Biogenic Amines in Pepperoni with Different Packaging Conditions. *Food Chemistry*, 89(2), 199-205.
- Kim, J.H., Kim, D.H., Ahn, H.J., Park, H.J., Byun, M.W. (2005). Reduction of the Biogenic Amine Contents in Low Salt-Fermented Soybean Paste by Gamma Irradiation. *Food Control*, 16(1): 43-49.
- Kim, M.J., Kwak, H.S., Jung, H.Y., Kim, S.S. (2016). Microbial communities related to sensory attributes in Korean fermented soy bean paste (doenjang). *International Food Research Journal*, 89 (Pt 1), 724-732.
- Kim, Y.S., Cho, S.H., Jeong, D.Y., Uhm, T.B. (2012). Isolation of Biogenic Amines-Degrading Strains of Bacillus subtilis and Bacillus amyloliquefaciens from Traditionally Fermented Soybean Products. *Korean Journal of Microbiology*, 48(3), 220-24.
- Kindossi, J.M., Anihouvi, V.B., Vieira-Dalodé, G., Akissoé, N.H., Jacobs, A., Dlamini, N., ... Hounhouigan, DJ. (2012). Production, consumption, and quality attributes of Lanhouin, a fish-based condiment from West Africa. *Food Chain*, 2(1), 117-130.
- Kirovska, I., Velickova, N. (2021). The Importance of Qualitative and Quantitative Biological Methods for Evaluation and Screening of Mycotoxins. *Albanian Journal of Medical and Health Sciences*, 55(February), 1-6.
- Kommenov, D., Levanovich, P.E., Rossi, N.F. (2019). Hypertension Associated with Fructose and High Salt: Renal and Sympathetic Mechanisms. *Nutrients*, 11(3), 569.
- Komprda, T., Smělá, D., Novická, K.A., Kalhotka, L., Šustová, K., Pechová, P. (2007). Content and Distribution of Biogenic Amines in Dutch-Type Hard Cheese. *Food Chemistry*, 102(1), 129-37.
- Kotsanopoulos KV, Arvanitoyannis IS. (2017). The Role of Auditing, Food Safety, and Food Quality Standards in the Food Industry: A Review. *Comprehensive Reviews in Food Science and Food Safety*, 16(5), 760-75.
- Kung, H.F., Lee, Y.C., Huang, Y.L., Huang, Y.R., Su, Y.C., Tsai, Y.H. (2017). Degradation of Histamine by Lactobacillus Plantarum Isolated from Miso Products. *Journal of Food Protection*, 80 (10), 1682-88.
- Lee, G.I., Lee, H.M., Lee, C.H. (2012). Food Safety Issues in Industrialization of Traditional Korean Foods. *Food Control*, 24(1), 1-5.
- Lee, J.H., Jin, Y.H., Park, Y.K., Yun, S.J., Mah, J.H. (2019). Formation of Biogenic Amines in Pa (Green Onion) Kimchi and Gat (Mustard Leaf) Kimchi. *Foods*, 8(3), 109.
- Lee, Y.C., Kung HF, Huang YL, Wu CH, Huang YR, Tsai YH. 2016. Reduction of Biogenic Amines during Miso Fermentation by Lactobacillus Plantarum as a Starter Culture. *Journal of Food Protection*, 79(9), 1556-61.
- Lin, X., Tang, Y., Hu, Y., Lu, Y., Sun, Q., Lv, Y., Zhang, Q., Wu, C., Zhu, M., He, Q. and Chi, Y., (2021). Sodium reduction in traditional fermented foods: Challenges, strategies, and perspectives. *Journal of Agricultural and Food Chemistry*, 69(29): 8065-8080.
- Liu, K. (2008). 14 – Food Use of Whole Soybeans. In *Soybeans*, edited by Lawrence A. Johnson, Pamela J. White, and Richard Galloway, 441-81. AOCS Press.
- Liu, K. (2012). *Soybeans: chemistry, technology, and utilization*. Springer.
- Liu, L., Chen, X., Hao, L., Zhang, G., Jin, Z., Li, C., Yang, Y., Rao, J., Chen, B. (2020). Traditional Fermented Soybean Products: Processing, Flavor Formation, Nutritional and Biological Activities. *Critical Reviews in Food Science and Nutrition*, 0 (0), 1-19.
- Ma, Y., He, F.J., MacGregor, G.A. (2015). High salt intake: independent risk factor for obesity? *Hypertension*, 66(4), 843-9.
- Mancia, G., Fagard, R., Narkiewicz, K., Redon, J., Zanchetti, A., Böhm, M., Christiaens, T., Cifkova, R., De Backer, G., Dominiczak, A., Galderisi, M. (2013). 2013 ESH/ESC Guidelines for the management of arterial hypertension. *Arterial Hypertension*, 17(2), 69-168.
- Mendes, R., Silva, H.A., Nunes, M.L., Empis, J.M. (2005). Effect of Low-Dose Irradiation and Refrigeration on the Microflora, Sensory Characteristics and Biogenic Amines of Atlantic Horse Mackerel (*Trachurus Trachurus*). *European Food Research and Technology*, 221(3), 329-35.
- Mohan, S., Campbell, N.R. (2009). Salt and high blood pressure. *Clinical Science*, 117(1), 1-11.
- Mullin, W.J., Xu, W. (2001). Study of soybean seed coat components and their relationship to water absorption. *Journal of Agricultural and Food Chemistry*, 49(11), 5331-5.
- Nazhand, A., Durazzo, A., Lucarini, M., Souto, E.B., Santini, A. (2020). Characteristics, Occurrence, Detection and Detoxification of Aflatoxins in Foods and Feeds. *Foods*, 9(5), 644.
- Novella-Rodríguez, S., Veciana-Nogués, M.T., Saldo, J., Vidal-Carou, M.C. (2002). Effects of high hydrostatic pressure treatments on biogenic amine contents in goat cheeses during ripening. *Journal of agricultural and food chemistry*, 50(25), 7288-92.
- Novita, N.R., Abidin, Z. (2020). Supporting Factors of Tempeh Products Quality in Sukawening Village, Dramaga District, Bogor Regency. *Jurnal Pusat Inovasi Masyarakat*, (PIM) 2(6), 925-930-925-930.
- Oh, H., Ryu, M., Heo, J., Jeon, S., Kim, Y.S., Jeong, D., Uhm, T.B. (2014). Characterization of Biogenic Amine-reducing Pediococcus pentosaceus Isolated from Traditionally Fermented Soybean Products. *Korean Journal of Microbiology*, 50(4), 319-26.

- Omotayo, O.P., Omotayo, A.O., Mwanza, M., Babalola, O.O. (2019). Prevalence of Mycotoxins and Their Consequences on Human Health. *Toxicological Research*, 35(1), 1–7.
- Ouoba, L.I.I., Diawara, B., Annan, N.T., Poll, L., Jakobson, M. (2005). Volatile compounds of 'soumbala', a fermented African locust bean, *Parkia biglobosa*, food condiment. *Journal of Applied Microbiology*, 99,1413-1421.
- Pachlová, V., Buňka, F., Flasarová, R., Válková, P., Buňková, L. (2012). The Effect of Elevated Temperature on Ripening of Dutch Type Cheese. *Food Chemistry*, 132(4), 1846–54.
- Park, H.S., Cho, H.Y., Shin, J.K. (2015). A study of sodium reduction effect in foods using fermented soy sauce. *Korean Journal of Food Science and Technology*, 47(4), 468-473.
- Park, Y.K., Lee, J.H., Mah, J.H. (2019). Occurrence and Reduction of Biogenic Amines in Traditional Asian Fermented Soybean Foods: A Review. *Food Chemistry*, 278 (April), 1–9.
- Peleteiro, B., Lopes, C., Figueiredo, C., Lunet, N. (2011). Salt intake and gastric cancer risk according to *Helicobacter pylori* infection, smoking, tumour site and histological type. *British journal of cancer*, 104(1), 198-207.
- Qin, H., Yang, H., Qiao, Z., Gao, S., Liu, Z. (2013). Identification and characterization of a *Bacillus subtilis* strain HB-1 isolated from Yandou, a fermented soybean food in China. *Food Control*, 31(1), 22-7.
- Qiu, S., Wang, Y., Cheng, Y., Liu, Y., Yadav, M.P., Yin, L. (2018). Reduction of Biogenic Amines in Sufu by Ethanol Addition during Ripening Stage. *Food Chemistry*, 239 (January), 1244–52.
- Rabie, M., Simin-Sarkadi, L., Siliha, H., El-seedy, S., El Badawy, A-A. (2009). Changes in free amino acids and biogenic amines of Egyptian salted-fermented fish (Feseekh) during ripening and storage. *Food Chemistry*, 115, 635-638.
- Renes, E., Diezhandino, I., Fernandez, D., Ferrazza, R.E., Tornadijo, M.E., Fresno, J.M. (2014). Effect of Autochthonous Starter Cultures on the Biogenic Amine Content of Ewe's Milk Cheese throughout Ripening. *Food Microbiology*, 44 (December), 271–77.
- Restuccia, D., Spizzirri, U.G., Puoci, F., Picci, N. (2015). Determination of Biogenic Amine Profiles in Conventional and Organic Cocoa-Based Products. *Food Additives & Contaminants: Part A*, 32(7), 1156–63.
- Rohm, B., Scherlach, K., Möbius, N., Partida-Martinez, L.P., Hertweck, C. (2010). Toxin Production by Bacterial Endosymbionts of a *Rhizopus Microsporus* Strain Used for Tempe/Sufu Processing. *International Journal of Food Microbiology*, 136(3), 368–71.
- Roy, K.W., Baird, R.E., Abney, T.S. (2001). A review of soybean (*Glycine max*) seed, pod, and flower mycofloras in North America, with methods and a key for identification of selected fungi. *Mycopathologia*, 150(1), 15-27.
- Ruiz-Capillas, C., Herrero, A.M. (2019). Impact of Biogenic Amines on Food Quality and Safety. *Foods*, 8(2), 62.
- Sanni, A.I., Onilude, A., Fadahunsi, I., Ogunbanwo, S., Afolabi, R. 2002. Selection of starter cultures for the production of ugba, a fermented soup condiment. *European Journal of Food Research and Technology*, 215:176-180.
- Samtiya, M., Aluko, R.E., Dhewa, T. (2020). Plant Food Anti-Nutritional Factors and Their Reduction Strategies: An Overview. *Food Production, Processing and Nutrition*, 2(1), 6.
- Santos, M.S. (1996). Biogenic Amines: Their Importance in Foods. *International Journal of Food Microbiology*, 29(2–3), 213–31.
- Seo, S.H., Cho, S.J. (2016). Changes in Allergenic and Antinutritional Protein Profiles of Soybean Meal during Solid-State Fermentation with *Bacillus Subtilis*. *LWT* 70 (July), 208–12.
- Shim, J.M., Lee, K.W., Yao, Z., Kim, H.J., Kim, J.H. (2016). Properties of Doenjang (Soybean Paste) Prepared with Different Types of Salts. *Journal of Microbiology and Biotechnology*, 26(9), 1533–41.
- Shin, D.H., Kwon, D.Y., Kim, Y.S., Jeong, D.Y. (2012). Science and Technology of Korean Gochujang. Seoul: Public Health Edu
- Shin, D.S., Park, C.H., Choi, I.D., Lee, S.K., Park, J.Y., Kim, N.G., Choi, H.S. (2019). Quality Properties of Soy-paste Soybean Cultivar for Fermented Soybean Products. *The Korean Journal of Food and Nutrition*, 32(2), 114–21.
- Singh, B., Karan, R., Singh, V. (2014a). Microbial quality and safety of Axone- Akhuni, a fermented soybean food of Nagaland. *Notore Med*, 2014,13
- Singh, T.A., Devi, K.R., Ahmed, G., Jeyaram, K. (2014b). Microbial and endogenous origin of fibrinolytic activity in traditional fermented foods of Northeast India. *International Food Research Journal*, 55:356–362.
- Shukla, S., Park, H.K., Kim, J.K., Kim, M. (2010). Determination of Biogenic Amines in Korean Traditional Fermented Soybean Paste (Doenjang). *Food and Chemical Toxicology*, 48(5), 1191–95.
- Shukla, S., Park, J., Park, J.H., Kim, M.K., Park, S., Dubey, A., Jeon, J., Khan, Y., Kim, M. (2017). Evaluation of Fungal Microflora for Aflatoxin Producing Possibility in Novel Quality Meju Fermented with Single and/or Multiple Additions of *Nelumbo Nucifera*, *Ginkgo Biloba*, and *Allium Sativum* Extracts. *Journal of Food Safety*, 37(4), e12368.
- Sivamaruthi, B.S., Kesika, P., Chaiyasut, C. (2019). Toxins in Fermented Foods: Prevalence and Preventions - A Mini Review. *Toxins*, 11(1), 4.
- Stadnik, J. J., Dolatowski, Z. (2010). Biogenic Amines in Meat and Fermented Meat Products. *Acta Scientiarum Polonorum Technologia Alimentaria*, 9(3).
- Sugawara, E. (2010). Fermented soybean pastes miso and shoyu with reference to aroma, in *Fermented Foods and Beverages of the World*, eds J. P. Tamang and K. Kailasapathy, (New York, NY: CRC Press, Taylor and Francis Group), 225–245.
- Sugiyama, M., Ae, N., Hajika, M. (2011). Developing of a simple method for screening soybean seedling cadmium accumulation to select soybean genotypes with low seed cadmium. *Plant and Soil*, 341, 413-422.
- Tamang, J.P., Shin, D.H., Jung, S.J., Chae, S.W. (2016). Functional properties of microorganisms in fermented foods. *Frontiers in Microbiology*, 7, 578.
- Tamang, J.P., Cotter, P.D., Endo, A., Han, N.S., Kort, R., Liu, S.Q., Mayo, B., Westerik, N., Hutkins, R. (2020). Fermented Foods in a Global Age: East Meets West. *Comprehensive Reviews in Food Science and Food Safety*, 19(1), 184–217.
- Ten Brink, B., Damink, C., Joosten, H.M., In't Veld, J.H. (1990). Occurrence and formation of biologically active amines in foods. *International journal of food microbiology*, 11(1), 73–84.
- Thakur, A., Sharma, V., Thakur, A. (2019). An Overview of Anti-Nutritional Factors in Food. *International Journal of Chemical Studies*, 7(1), 2472-2479.
- Wang, L.J., Li, D., Zou, L., Dong Chen, X., Cheng, Y.Q., Yamaki, K., Li, L.T. (2007). Antioxidative Activity of Douchi (A Chinese Traditional Salt-Fermented Soybean Food) Extracts During Its Processing. *International Journal of Food Properties*, 10(2), 385–96.
- Wang, L.J., Yin, L.J., Li, D., Zou, L., Saito, M., Tatsumi, E. (2007). Influences of Processing and NaCl Supplementation on Isoflavone Contents and Composition during Douchi Manufacturing. *Food Chemistry*, 101(3), 1247–53.
- Whelton, P.K., Carey, R.M., Aronow, W.S., Casey, D.E., Collins, K.J., Dennison Himmelfarb, C., DePalma, S.M., Gidding, S., Jamerson, K.A., Jones, D.W., MacLaughlin, E.J. (2018). 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/AphA/ASH/ASPC/NM A/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Journal of the American College of Cardiology*, 71(19), e127-e248.
- Winarno, F.G., Fardiaz, S., Daulay, D. (1973). Indonesian fermented foods. Indonesia: Department of Agricultural Product Technology, Bogor Agricultural University.
- Worldwide Regulations for Mycotoxins in Food and Feed in (2003). N.d. Accessed July 5, 2021. <http://www.fao.org/3/y5499e/y5499e07.htm>.
- Wu, F. (2004). Mycotoxin risk assessment for the purpose of setting international regulatory standards, 4049-4055.
- Xiang, H., Sun-Waterhouse, D., Waterhouse, G.L., Cui, C., Ruan, Z. (2019). Fermentation-Enabled Wellness Foods: A Fresh Perspective. *Food Science and Human Wellness*, 8(3), 203–43.
- Yang, H., Yang, L., Zhang, J., Li, H., Tu, Z., Wang, X. (2019a). Exploring functional core bacteria in fermentation of a traditional Chinese food, *Aspergillus-type douchi*. *PLoS One*, 14(12), e0226965.

- Yang, J., Wu, X.B., Chen, H.L., Sun-Waterhouse, D., Zhong, H.B., Cui, C. (2019b). A value-added approach to improve the nutritional quality of soybean meal by product: enhancing its antioxidant activity through fermentation by *Bacillus amyloliquefaciens* SWJS22. *Food Chemistry*, 272, 396–403.
- Zain, M.E. (2011) Impact of Mycotoxins on Humans and Animals. *Journal of Saudi Chemical Society*, 15(2), 129–44.
- Zhang, B., Chen, P., Florez-Palacios, S.L., Shi, A., Hou, A., Ishibashi, T. (2010). Seed quality attributes of food-grade soybeans from the US and Asia. *Euphytica*, 173(3), 387-96.
- Zhang, Q., Lin, S., Nie, X. (2013). Reduction of Biogenic Amine Accumulation in Silver Carp Sausage by an Amine-Negative *Lactobacillus Plantarum*. *Food Control*, 32(2), 496–500.
- Zhang, Y.T., Yu, B., Lu, Y.H., Wang, J., Liang, J.B., Tufarelli, V., Laudadio, V., Liao, X.D. (2017). Optimization of the Fermentation Conditions to Reduce Anti-Nutritive Factors in Soybean Meal. *Journal of Food Processing and Preservation*, 41(5), e13114.
- Zhao, H., Wang, Y., Zou, Y., Zhao, M. (2013). Natural Occurrence of Deoxynivalenol in Soy Sauces Consumed in China. *Food Control*, 29(1), 71–75.
- Zheng, L., Li, D., Li, Z.L., Kang, L.N., Jiang, Y.Y., Liu, X.Y., Chi, Y.P., Li, Y.Q., Wang, J.H. (2017). Effects of *Bacillus* Fermentation on the Protein Microstructure and Anti-Nutritional Factors of Soybean Meal. *Letters in Applied Microbiology*, 65(6), 520–26.
- Zhi, Y., He, K., Sun, T., Zhu, Y., Zhou, Q. (2015). Assessment of Potential Soybean Cadmium Excluder Cultivars at Different Concentrations of Cd in Soils. *Journal of Environmental Sciences*, 35(September), 108–14.
- Zhi, Y., Sun, T., Zhou, Q., Leng, X. (2020). Screening of Safe Soybean Cultivars for Cadmium Contaminated Fields. *Scientific Reports*, 10(1), 12965.
- Zhu, Y.P., Cheng, Y.Q., Wang, L.J., Fan, J.F., Li L.T. (2008). Enhanced antioxidative activity of Chinese traditionally fermented okara (meitauza) prepared with various microorganism. *International Journal of Food Properties*, 11(3), 519-29.