

# Sorghum Flakes: Studies of the Type and Concentration of Sorghum Flour

*by* Cek Turnitin4

---

**Submission date:** 30-Oct-2023 08:59AM (UTC+0700)

**Submission ID:** 2211221742

**File name:** 42-\_Sorghum\_Flakes\_-\_Endang\_Noerhartati.pdf (614.27K)

**Word count:** 3016

**Character count:** 16219

## Sorghum Flakes: Studies of the Type and Concentration of Sorghum Flour

Endang Noerhartati<sup>1\*</sup> dan Diana Puspitasari<sup>2</sup>

<sup>1,2</sup>Study Program of Agroindustrial Technology - Faculty of Engineering-Wijaya Kusuma Surabaya  
University, East-Java, Indonesia

Jl. Dukuh Kupang XXV/54 Surabaya

\*Email: [endang.noerhartati@gmail.com](mailto:endang.noerhartati@gmail.com) ; 08563046119

### ABSTRACT

This study aimed at finding the best treatment combination of sorghum flakes using Randomized Block Design (RBD) 2 factors with 3 repeated measures. Factor I: types of flour (T): T<sub>1</sub>: red sorghum flour (Sorghum bicolor) and T<sub>2</sub>: white sorghum flour (KD4). Factor II: Concentration (K): K<sub>1</sub>: 25%; K<sub>2</sub>: 50%; K<sub>3</sub>: 75% and K<sub>4</sub>: 100%. Observations: organoleptic test on parameters of taste, color, flavour and crispness, moisture content and starch. Data analysis used organoleptic test with Friedman test, while the water content and starch used analysis of variance test (ANOVA) at the 5% level followed by Duncan's Test. The results: 1) Flake sorghum was one of the alternative forms of diversification products based on sorghum flour, 2) a combination of the best treatment for sorghum flakes T<sub>1</sub>K<sub>2</sub> (treatment type sorghum flour (T): T<sub>1</sub> = Red Flour/Sorghum bicolor (T<sub>1</sub>) and the concentration (K) : K<sub>2</sub> = 50%), the results of Organoleptic parameters were: taste of 4,68 (really like) , color of 4,62 (really like), flavor of 4,58 (really like), crispness 4,68 (really like), water content 2,19% and starch 20,8% and 3) a combination of the best treatments for sorghum flakes second: T<sub>2</sub>K<sub>2</sub> (types of sorghum flour treatment (T): white flour / KD4 (T<sub>2</sub>) and concentration (C): K<sub>2</sub> = 50%), the results of organoleptic parameter were taste of 4,62 (really like), color of 4,65 (really like), flavor of 4,52 (really like), crispness of 4,62 (really like), water content of 2,17% and starch of 34,42%.

Keywords: sorghum flour, flakes, organoleptic testing, moisture content and starch

### I. INTRODUCTION

Sorghum (Sorghum bicolor) as the world's food source is the fifth rank after wheat, rice, corn and barley. Sorghum has the potential of relatively higher yield than rice, wheat and maize. Sorghum yields can exceed 6-7 tonnes/ha. Sorghum development in Indonesia is now considered indispensable either as a material for industrial or alternative food materials, its production is used as food, beverages, animal feed and other industrial interests. The pictures of sorghum gardens are presented in Figure 1.



Figure 1. Sorghum gardens in Keyongan Village - Babat Subdistrict Lamongan District

One of sorghum results is sorghum flour. A product that can be made from sorghum flour is sorghum flakes. Flake is one of food products that uses cereal foods such as rice, wheat or corn and tubers for examples potatoes, cassava, sweet potatoes. Flake is classified into the types of ready made which is processed and modified depending on the types and forms. To understand the existence of the development of sorghum plant towards the environment (implications of its existence), if it is developed maximally will be creating a chain of the development of national sorghum cluster which support the economy of Indonesia, and open up the opportunities of sorghum plant development, which is directed in line with the effort to increase the empty land productivity (marginal land, idle land, and other non-productive land). That effort is a step to safe the environment. The objective of the research at: introducing the alternative food sources, namely sorghum flour; increasing the potential of sorghum into flake product with added value; and obtaining the best treatment for the combination of sorghum flakes.

## II. MATERIALS AND METHODS

The materials used in this research were in the form of red sorghum flour (*Sorghum bicolor*), white sorghum flour (KD 4), wheat flour, sugar, margarine, and egg yolk. Tools used in this research were knife, basin, cutting board. oven, grinding, mixer, flour grinding machine, and flakes mold.

This research used Randomized Block Design (RBD) of 2 factors with 3 repeated measures. Factor I: types of sorghum flour (T): T<sub>1</sub>: red sorghum flour (*Sorghum bicolor*) and T<sub>2</sub>: white sorghum flour (KD 4). Factor II: Concentration of sorghum flour (K): K<sub>1</sub>: 25%; K<sub>2</sub>: 50%; K<sub>3</sub>: 75% and K<sub>4</sub>: 100%. Flow diagram of sorghum flakes process can be seen in Figure 2. Analysis Method: Chemical Analysis conducting on

sorghum flakes products was water content and starch Analysis, and organoleptic test was conducted by using scoring method. In this method, panelists were asked to rate the level of their favorite tastes, colours, flavours, and crisps of sorghum flakes products by providing value or score as below: 5 = really like, 4 = like, 3 = neutral, 2 = somewhat like, and 1 = dislike. Panelists on this research were around 30 persons, which were selected by random, so they were expected to represent overall consumers.




Figure 2. Flow Diagram of Sorghum Flakes Process

Data processing for water and carbohydrate contents of sorghum flakes were conducted by using variant analysis, if there were differences then Duncan Test was conducted at the 5% confidence level, whereas for the organoleptic test with Friedman test. Alternative selection aimed at finding of the best alternative process. It was based on the concept of the acquisition of the expected value. The decision concept of expected value is to select a decision which has the maximum payoff (profit or usefulness). According to Siagian (1987), mathematical equation for payoff of the expected value is as below:

$$E_{pj} = \sum_{i=1}^n P(x_i) \cdot f(x_i, d_j)$$

Note:  $E_{pj}$  = Expected value of payoff  
 $P(x_i)$  = The probability of each  $x_i$  ground state  
 $x_i$  = Different ground state  
 $d_j$  = Decisions which are taken into account  
 $f(x_i, d_j)$  = Acquisition of the ground state and  $d_j$  decision

### III. RESULTS AND DISCUSSION

Observation results of organoleptic test for sorghum flakes with taste, color, flavor, and crispness parameters are presented in Figure 3, 4, 5, and 6, while results of Friedman test for sorghum flakes is presented on Table 1.




Figure 3. Calculation of results from organoleptic test percentage of taste parameter

Table 1. The result of the Friedman Test for sorghum flakes products

Parameter	Taste	Color	Flavour	Crispness
N	60	60	60	60
Chi-Square	12,476	19,893	7,470	3,403
df	7	7	7	7
Asymp. Sig.	0,086	0,006	0,382	0,845

a. Friedman Test




Figure 4. Calculation of results from organoleptic test percentage of color parameter

Based on the result of organoleptic test for sorghum flakes were: taste, color, flavor, and crispness parameters, showed that the highest score for taste parameter was in the treatment of T1K2 (T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K2= 50%), with percentage of 68.3% score 5 (really like), the second was in treatment of T2K2 (T1= white flour / KD4 (T2) and sorghum flour concentration K2= 50%), with percentage of 63.3% score 5 (really like) and the treatment T1K3 (T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K3= 75%);




Figure 5. Calculation of results from organoleptic test percentage of flavour parameter



Figure 6. Calculation of results from organoleptic test percentage of crispness parameter

The highest score for color parameter was in the treatment of T2K2 (T1= white flour / KD4 (T2) and sorghum flour concentration K2= 50%), with percentage of 66.6.% score 5 (really like), the second was in the treatment of T1K (T1= T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K2= 50%), with percentage of 63.3.% score 5 (really like); the highest score for flavour parameter was in the treatment of T1K2 (T1= T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K2= 50%), with percentage of 63.3.% score 5 (very favorable), the second was in the treatment of T1K1 and T1K3 (T1= T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K1= 25% and K3= 75%), with with percentage of 60.0% score 5 (really like); the highest score for crispness parameter was in the treatment of T1K2 (T1= T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K2= 50%), with percentage of 68.3% score 5 (really like), the second was in the treatment of T2K1 (T1= white flour / KD4 (T2) and sorghum flour concentration K2= 50% and K3= 75%), both of them have percentage of 65% score 5 (really like).

On Table 1, it can be seen the result of Friedman Test for sorghum flakes showed that color parameter was significantly different (Sig 0,006 < 0,05), while taste parameter <sup>2</sup>(Sig 0,086 > 0,05), flavor (Sig 0,382 > 0,05) and crispness (Sig 0,845 > 0,05) were not significantly different. This means that for color parameter of sorghum flakes was influenced by both factors, the type and sorghum flour concentration. Color of sorghum flakes was very influenced by color of sorghum flour and the more flour was used, then the color was getting darker. The result of water content test was presented in Figure 7, showed that water content of sorghum flakes was around 2.17-2.22%.



The result of variant analysis showed that water content of sorghum flakes was around Sig 0,144 > 0,05, meant that water content of sorghum flakes was not significantly different. This was in accordance with the result of organoleptic test for crispness, average percentage of more than 63% score 5 (really like).




Figure 7. Calculation of results from water content of sorghum flakes

The result of starch test was presented in Figure 8, showed that starch of sorghum flakes was around 18.13-35.08. The result of variant analysis showed that starch of sorghum flakes was around Sig 0,00 < 0,05, this meant that starch of sorghum flakes was significantly different and there was an interaction Sig 0,00 < 0,05. Data analysis for starch of sorghum flakes was continued by Duncan Test at the confidence level of 5% such as presented on Table 2, showed that there was difference between starch of the both sorghum flour, in which red sorghum flour highest starch lowest than white flour and the more flour was used.




Figure 8. Calculation of results from starch



Table 2. Calculation of results from an interaction of treatment combinations for starch

		Starch (%)								
	Interaction Type and Concentration Flour	N	Subset							
			1	2	3	4	5	6	7	8
Duncan <sup>a,b</sup>	Interaction T1 and K1	3	18.1300 a							
	Interaction T1 and K2	3		20.8700 b						
	Interaction T1 and K3	3			24.4600 c					
	Interaction T1 and K4	3				24.8967 d				
	Interaction T2 and K1	3					31.8100 e			
	Interaction T2 and K2	3						34.4167 f		
	Interaction T2 and K3	3							34.5067 g	
	Interaction T2 and K4	3								35.4100 h
	Sig.		1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000

Means for groups in homogeneous subsets are displayed.  
 Based on observed means.  
 The error term is Mean Square(Error) = ,002.  
 a. Uses Harmonic Mean Sample Size = 3,000.  
 b. Alpha = ,05.  
 c. Mean in columns with noted by different alphabets show a significant level (Duncan Test at 5%)

And then starch of sorghum flakes was increasing Calculation of results for probability of sorghum flakes was presented in Figure 9. Probability or opportunity was confidence level of a person against an uncertain event. Probability analysis was conducted to determine the probability of each ground state. Ground state of quality includes taste, color, flavour, crispness, water content, and starch. Probability value showed the interest rate of a ground state, the greater the probability value of a ground state, then the most important that ground state. In the interest of sorghum flakes, taste parameter (24.5%) was considered as the most important parameter when compared to the other parameter, i.e. color (20.25%), crispness (19%), water content (15.25%), starch (12.75%), and flavor (8.25%) respectively.




Figure 9. Calculation of results for probability of sorghum flakes based on the highest order.

The existing alternative process was compared to determine the optimal process. This alternative selection was conducted by calculating the expected value which was obtained from each alternative process. Calculation of results for the expected value of each alternative process was presented in Figure 10.




Figure 10. Calculation of results for the expected value of sorghum flakes.

Based on calculation of results for the expected value, then the alternativetreatment selected was the treatment of T1K2 (T1= red flour / *Sorghum bicolor* (T1) and sorghum flour concentration K2= 50%), with calculation of results for the expected value was about 7.73, the second was the treatment of T2K1 (T2= white flour /

KD4 (T2) and sorghum flour concentration K2= 50%), with calculation of results for the expected value was about 7.2. This means that treatment was based on the best quality if compared to the other treatment. Picture of materials and products of sorghum flakes was presented in Figure 11.




Figure 11. Materials and Products of Sorghum Flakes

### The Best Treatment Combinations of Sorghum Flakes




Figure 12. The best treatment combinations of sorghum flakes treatment of sorghum flour type (T): T1= red flour / *Sorghum bicolor* (T1) *Sorghum bicolor* (T1) and white flour / KD4 (T2); treatment of sorghum flour concentration (K): K2= 50%.

#### IV. CONCLUSIONS

Based on research findings, it can be concluded that: Sorghum flakes is an alternative product diversification based on sorghum flour. The best treatment combinations for sorghum flakes is the treatment of T<sub>1</sub>K<sub>2</sub> (treatment of sorghum flour type (T): T1= red flour / *Sorghum bicolor* (T1) and treatment of sorghum flour concentration (K): K2= 50%), and the treatment of T<sub>2</sub>K<sub>2</sub> (treatment of sorghum flour type (T): white flour / KD4 (T2) and treatment of sorghum flour concentration (K): K2= 50%).

#### REFERENCES

1. [epkes](#). 1996. List of Food Composition. Publisher Bhratara. Jakarta
2. Kentz, N.L. 1970. Technology of Cereal. Pergamon Press. Sydney.
3. Metz, S.A. 1970. Cereal Technology. The AVI Publishing Company. USA.
4. National Sorghum Producers. 2007. Grain Sorghum Production Handbook. University of Arkansas Division of Agriculture. North Inerstate 27 lubbock. Texas ([www.sorghumgrowers.com](http://www.sorghumgrowers.com))
5. National Sorghum Producers. 2007. Sorghum Grower. University of Arkansas Division of Agriculture. North Inerstate 27 lubbock. Texas ([www.sorghumgrowers.com](http://www.sorghumgrowers.com))
6. Noerhartati, E. 2009. Modul of Industrial Materials Second Edition. Study Program: TIP, Faculty of Engineering, UWKS, January 2009

7. \_\_\_\_\_, 2010. Various Industrial Products Made from Raw Corn and Sorghum. Business Meeting Papers And Retrieval Technology ". Department of Agriculture Government of EastJava province July 27 to 28, 2010 - Batu Malang
8. \_\_\_\_\_, 2010. Assorted Cookies Made BakuTepung Sorghum (Sorghum bicolor). The paper "The Unyil Coverage TV TRANS 7". Cooperation of Study Program of Agro Industrial Technology, Faculty of Engineering-University of Wijaya Kusuma Surabaya Si Unyil TV Coverage TRANS 7 (2 September 2010)
9. \_\_\_\_\_, 2012. Post Harvest Technology and Processing of Commodity Wheat and sorghum. Alternative Food Development Coordination Meeting "Agriculture Department Government ofEast Java. Surabaya. (11 April 2012)
10. \_\_\_\_\_, 2012. Various Food Products Processing Technology of Wheat and Sorghum. Increased Activity Commodity Production and Productivity in East Java Other Cereal Year 2012 Department of Agriculture Government of East Java. Malang. (June 19, 2012)
11. \_\_\_\_\_, 2014. Alternative Food Ingredients Sorghum (Sorghum sp): Seeds, Flour, Assorted Cookies and liquid sugar sorghum stems. Media Sciences. Sidoarjo
12. \_\_\_\_\_, 2014. Business Plan Agroindustri Sorghum. Adoption Facilitation Partnership activities and Alternatives Food Technology. Agriculture Office East Java Provincial Government. Surabaya
13. \_\_\_\_\_, 2014. Business Incubator Pastry Sorghum. Sorghum Development Facilitation activities. Agriculture Office East Java Provincial Government. Lamongan.
14. Illaningtyas, F., Sri I, Sri P, IIM A, and Fajarwati. 2013. Effects of Supplementation of Sorghum Protein Isolates Against Biological and Chemical Properties Appearance Confectionery Sorghum. **Proceedings of the National Seminar. PATPI 2013. Jember. It 51-66**
15. Sirappa, M.P.2003. Sorghum Development Prospects As Alternative Commodities Food, Feed and Industrial. Journal of Agricultural Research, 22 (4) Stonestreet, N. J. M. S Alawi and SR. Bean. 2010. Sorghum Proteins: The Concentration, Isolation, modification and food applications of infidels. J. Food Science Vol. 75 (5): R90-104
16. Revitriani, M, Endang RW, and Diana P. 2013 Flour Purse Concentration Study On Making Wet Foodle. J. REKA Agro Industry. I (1) 2013: 28-36
17. Taylor. J.R. Schober T.J and Bean, S.R. 2006. Novel Food and Non-Food Uses for Sorghum and Millets. J. Cereal Scier, 44(3): 252-271
18. Yousif,A., D.Nhepera and S.Johson. 2012. Influence of Sorghum Floor Addition on Flat Bread In Vitro Starch Digestibility, Antioxidant, Capacity and Consumer Acceptability Food. Ford. Chem. 134 (2012): 880-887

# Sorghum Flakes: Studies of the Type and Concentration of Sorghum Flour

---

## ORIGINALITY REPORT

---

11%

SIMILARITY INDEX

7%

INTERNET SOURCES

8%

PUBLICATIONS

5%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Syam, Netty, Tatik Wardiyati, Muhammad Dawam Maghfoer, Eko Handayanto, Bahtiar Ibrahim, and Aminah Muchdar. "Effect of Accumulator Plants on Growth and Nickel Accumulation of Soybean on Metal-contaminated Soil", Agriculture and Agricultural Science Procedia, 2016. Publication **3%**
  - 2** [jurnal.unej.ac.id](http://jurnal.unej.ac.id) Internet Source **2%**
  - 3** Submitted to University of Sunderland Student Paper **1%**
  - 4** [acikbilim.yok.gov.tr](http://acikbilim.yok.gov.tr) Internet Source **1%**
  - 5** Submitted to Curtin University of Technology Student Paper **1%**
  - 6** Submitted to Higher Education Commission Pakistan Student Paper **1%**
-

7

[journal.uwks.ac.id](http://journal.uwks.ac.id)

Internet Source

<1 %

8

Submitted to Asian Institute of Technology

Student Paper

<1 %

9

[digitalcommons.uri.edu](http://digitalcommons.uri.edu)

Internet Source

<1 %

10

Submitted to Hopkinton High School

Student Paper

<1 %

11

[pustaka.setjen.pertanian.go.id](http://pustaka.setjen.pertanian.go.id)

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off