

Original Research Article

The Relationship between Maternal Education Level and Family Income on the Incidence of Stunting in Toddlers at the Madiun City Health Center**Nurwasilah^{1*}, Sukma Sahadewa², Andiani³**Faculty of Medicine, Wijaya Kusuma University Surabaya¹Department of IKM, Faculty of Medicine, Wijaya Kusuma University Surabaya²Department of IKM, Faculty of Medicine, Wijaya Kusuma University Surabaya³

Jl. Dukuh Kupang XXV No. 54, Dukuh Kupang, Kec. Dukuh, Surabaya City, East Java

Correspondence author email: wasilahnur761@gmail.com

Phone: 082157418818

Abstract

Stunting is a condition that occurs in toddlers who experience growth failure due to chronic malnutrition based on a z-score (TB/U) value of less than -2 SD (Standard Deviation). *Stunting* can be caused by a low level of maternal knowledge. Family income is also a risk factor for *stunting*. The purpose of this study was to analyze the relationship between maternal education level and family income on the incidence of *stunting* in toddlers at the Madiun City Health Center. This study used an analytical observational method with a *case control* design. The population in this study was all mothers who had toddlers in Madiun City. Respondents in this study consisted of 34 people, obtained from questionnaires with the total sampling method. The results of *Chi-Square* analysis showed that there was no relationship between maternal education level ($p = 0.290$), no relationship between family income level ($p = 0.628$), maternal education level and family income level ($p = 0.545$) greater than p value = 0.05; showed that there was no relationship between maternal education level and family income level with the incidence of stunting in toddlers at the Madiun City Health Center. There is no relationship between maternal education level and family income with the incidence of stunting in toddlers at the Madiun City Health Center.

Keywords: *Stunting*, maternal education, family income

Received: _____ Revised: _____ Accepted: _____

INTRODUCTION

Stunting is a condition that occurs in toddlers who experience growth failure due to chronic malnutrition based on a z-score (TB/U) value of less than -2 SD (Standard Deviation). Event *stunting* It can be caused by decreased cognitive function, decreased concentration, decreased learning power and school achievement, memory impairment, and impaired motor development of children, and has a long-term impact on decreased productivity that can hinder economic growth and cause intergenerational poverty. (Darmini et al., 2022; Rahmawati, 2020).

Number event *stunting* Currently it is still a nutritional problem in toddlers. The prevalence in the world is around 21.9% or 149 million toddlers experience *stunting* (Setyo et al., 2022). In Indonesia, the prevalence of children aged less than five years (toddlers) who suffer *stunting* is the second highest in Southeast Asia with an incidence rate of 31.8% in 2020. Incidence rate *stunting* The highest in Indonesia is in Timor Leste at 48.8%. In Southeast Asia, Laos is in position after Indonesia with a prevalence of 30.2%.

Cambodia then ranks fourth by incidence figures *stunting* under-five by 29.9%. The Philippines follows with prevalence rates *stunting* by 28.7%. Meanwhile, Singapore is the prevalence rate of children sufferers *stunting* the lowest in Southeast Asia with a prevalence rate of only 2.8% (Asian Development Bank, 2021).

According to the results of the Indonesian Nutritional Status Study (SSGI) by the Ministry of Health, the incidence rate of toddlers *stunting* in 2021 it was 24.4%. These are *stunting* experienced by almost a quarter of Indonesian toddlers last year. In 2021, there were 23.5% of toddlers who experienced *stunting* in East Java Province (Ministry of Health of the Republic of Indonesia, 2021). Current prevalence *stunting* in Madiun City at 12.4%. Even though it is far below the 14% figure that is the national target in 2024, the Madiun City Government continues to reduce the incidence rate *stunting* with a target of reaching 0%. (Madiun City Health Office, 2019).

Some influencing factors *stunting* Among them are poverty, lack of awareness about health, poor nutrition and also improper parenting causing growth and development failure in toddlers. *Stunting* can also be caused by the factor of low maternal knowledge level (Agustin & Rahmawati, 2021). The level of education in a person can be affected when a person receives information. People with better education will be easier to receive information than people with less level of education. This information is used as a provision for mothers to take care of their children in everyday life. The maternal's education level usually affects the maternal's knowledge about toddler nutrition. Where the higher the education, it will be easy to absorb information about health, for example nutritional knowledge. The maternal's level of knowledge plays a significant role with the incidence *stunting*. (Sari & Zelharsandy, 2022; Husnaniyah et al., 2020).

Level Education of the population in Madiun City with elementary school graduates and below the percentage is 12.87%, junior high school is 14.05%, high school is 19.93%, the highest percentage is SMK graduates at 26.31%, Diploma I/II/III at 5.23%, while residents who graduated from higher education are S1/Diploma IV at 21.60% (Central Bureau of Statistics Madiun, 2022). Based on these data, it can be seen that the most education completed by residents in Madiun City in 2022 is SMK, which is 26.31%, while the least is Diploma I/II/III, which is 5.23% (Central Bureau of Statistics Madiun, 2022).

Family income factors are also risk factors of *stunting*. This is because parents are able to meet all the primary and secondary needs of their children, an adequate family income will support their growth and development (Soetjiginisih, 2012). Families with incomes less than the Regional Minimum Wage are 6 times more likely to experience *stunting* (Agustin & Rahmawati, 2021). According to the Central Statistics Agency (BPS), the economy of Madiun City grew by 4.73% in 2021. The economic growth of Madiun City in 2021 which reached 4.73 percent, increased 7% from 2020 which was minus 3.39 percent.

The purpose of this study was to analyze the relationship between maternal education level and family income on incidence *stunting* in toddlers at the Madiun City Health Center.

METHODS

This study used an analytical observational method with a *case control* design. In this study to determine the relationship between maternal education level and family income on the incidence of *stunting* in toddlers at the Madiun City Health Center. The population in this study was all mothers who had toddlers in Madiun City. Respondents in this study consisted of 34 people, obtained from questionnaires with the total sampling method.

RESULTS

1. Univariate Analysis Results

Figure 1. Distribution of Stunting Incidence

Respondents in this study consisted of 34 people (100%) who suffered from stunting as much as 50% (17 respondents) and those who did not suffer from stunting as much as 50% (17 respondents). The distribution of respondents based on the degree of Stunting Incidence can be seen in table 1 below.

Table 1: Distribution of respondent on stunting incidence

Stunting Events	No. of respondents	%
Stunting	17	50
No stunting	17	50
Total	34	100

Source: Research 2023

Figure 2. Distribution of Toddlers Age

The ages of respondents in this study were mostly aged ≤ 10 months is 67.6% (23 respondents), while the remaining 32.4% (11 respondents) are aged > 10 months. The distribution of respondents by sex can be seen in table 2 below.

Table 2: Frequency Distribution of Respondents by Age

Age	No. of respondents	%
≤ 10 months	23	67,6
> 10	11	32,4
Total	34	100

Source: Research 2023

Figure 3. Distribution of Toddlers Gender

Most of the respondents in this study had a female gender, which was 58.8% (20 respondents), while the remaining 41.2% (14 respondents) had a male gender. The distribution of respondents by gender be seen in table 3 below.

Table 3: Frequency distribution of respondents by gender

Gender	No. of respondents	Percent
Woman	20	58,8
Man	14	41,2
Total	34	100

Source: Research 2023

Figure 4. Distribution of Maternal’s Occupation

In this study, respondents who fell into the working category were 35.3% (12 respondents) and respondents who were included in the non-working category as many as 64.7% (22 respondents). The distribution of respondents based on maternal’s occupation can be seen in table 4 below.

Table 4: Frequency Distribution of Respondents by Maternal’s Occupation

Category	No. of respondents	Percent
Work	12	35,3
Does not work	22	64,7
Total	34	100

Source: Research 2023

Figure 5. Distribution of Maternal Education Level

In this study, respondents who had a low level of education were 38.2% (13 respondents) and respondents who had higher education as many as 61.8% (21 respondents). The distribution of respondents based on maternal education level can be seen in table 5 below.

Table 5 : Frequency distribution of respondents based on maternal education level

Category	No. of respondents	Percent
Lower education	13	38,2
Higher education	21	61,8
Total	34	100

Source: Research 2023

Figure 6. Distribution of Family Income Level

In this study, respondents who had income < MSEs in Madiun City were 14.7% (5 respondents) and respondents who had income ≥ UMK Kota Madiun as much as 85.3% (29 respondents). The distribution of respondents based on family income level can be seen in table 6 below.

Table 6 : Distribution of respondents' frequency by family income level

Category	No. of respondents	Percent
< UMK Kota	5	14,7
≥ UMK City	29	85,3
Total	34	100

Source: Research 2023

Figure 7. Distribution of Maternal Education Level and Family Income Level to Stunting Groups

In this study, respondents who had risk factors based on maternal education level and income were 8.8% (3 respondents), respondents who did not have risk factors based on maternal education level and income were 91.2% (31 respondents). The distribution of respondents based on parental employment status can be seen in table 7 below.

Table 7: Distribution of respondents' frequency based on maternal's education level, family income level against stunting groups

Category	No. of respondents	Percent
High risk (<3)	3	8,8
Low risk (≥3)	31	91,2
Total	34	100

Source : Research 2023

2. Results of Bivariate Analysis

Figure 8. The relationship between the maternal's education level and the incidence of stunting that occurs in toddlers at the Madiun City Health Center.

Table 8: Results of Hypothesis Test Based on Maternal Education Level on Stunting Incidence

Maternal's Education Level	Stunting				P-Value
	Case		Control		
	N	%	N	%	
Lower Education	8	47%	5	29%	0,290
Higher Education	9	53%	12	71%	

Source : Research 2023

From the results of the analysis *Chi-Square* indicates that the significance value of 0.290 is greater than the value of p value = 0.05; showed that there was no relationship between the maternal's education level and the incidence of stunting in toddlers at the Madiun City Health Center.

Figure 9. The relationship between family income level and the incidence of stunting that occurs in toddlers at the Madiun City Health Center.

Table 9: Results of Hypothesis Test Based on Family Income Level on Stunting Incidence

Family Income Level	Stunting				P-Value
	Case		Control		
	N	%	N	%	
< MSEs	2	12%	3	18%	0,628
≥ UMK	15	88%	14	82%	

Research Source : 2023

From the results of the analysis *Chi-Square* indicates that the significance value of 0.628 is greater than the value of p value = 0.05; showed that there was no relationship between family income level and the incidence of stunting in toddlers at the Madiun City Health Center.

Figure 10. The relationship between the maternal's education level and family income level with the incidence of stunting that occurs in toddlers at the Madiun City Health Center.

Table V.10: Hypothesis Test Results Based on Maternal Education Level and Family Income Level

Risk Factors (Maternal's Education Level and Family Income Level)	Stunting				P-Value
	Case		Control		
	N	%	N	%	
High Risk	2	12%	1	6%	0,545
Low Risk	15	88%	16	94%	

Source: Research 2023

From the results of the analysis *Chi-Square* indicates that the significance value of 0.545 is greater than the value of p value = 0.05; showed that there was no relationship between maternal education level and family income level with the incidence of stunting in toddlers at the Madiun City Health Center.

DISCUSSION

Figure 4 and 6 showed that respondents who had income < MSEs in Madiun City as much as 14.7% and respondents who had income in MSEs ≥ in Madiun City as much as 85.3%. This shows that the majority of toddler families in the Madiun City Health Center have MSE income in Madiun City. ≥

Both high and low income levels impact a family's ability to afford nutritious food. A family with sufficient money will be able to meet all the primary and secondary needs of its children. Families with good financial condition also have easier access to health services. Children from low-income families typically eat less, both in terms of quantity and variety. People with high socioeconomic levels tend to choose and buy a variety of healthy foods (Sutarto *et al.*, 2020).

Low family income has a direct impact on stunting because it forces families to choose less nutritious foods to meet their nutritional needs. In addition, due to poor food quality, more and more children are malnourished, which causes nutritional problems in toddlers (Yanti, 2021). Stunting is more prevalent in children from low-income families than children from well-off or high-income families (Habimana, 2019).

The results of Wati's research (2018) show that most toddlers with low nutritional status have low-income families. Families with low incomes will usually find it difficult to meet their nutritional needs. Families with middle to lower economic status have the possibility to consume food with low nutrition in toddlers, and this has an impact on the nutritional status of toddlers.

The results found that out of 100% of toddlers, there are 50% of toddlers who are stunted and 50% of toddlers who are not stunted. Some toddlers who are not stunted occur because their parents have provided them with all the essential nutrients for their growth and development, including minerals, vitamins, fats, carbohydrates, and proteins. While stunted toddlers are caused by mothers who do not pay enough attention to the food and health needs of their children, this problem can also be caused by infectious infections that worsen children's health and interfere with their growth (Fakhma and Dhewi, 2020).

Figure 5 showed that a highly educated mother may have a better outcome for her child's health and nutrition. The maternal's level of education can ease the mother's ability to understand and absorb nutritional information. Education is needed so that there are more mothers who are able to solve nutrition and nutrition problems in the family and can act as soon as possible if they are not fulfilled (Sutarto *et al.*, 2020).

The function of maternal education is to increase children's insight into themselves and their environment. The length of education taken determines the level of education of the mother. Highly educated mothers are easier to digest health information, especially in educating toddlers every day. Toddler development can be influenced by a number of factors, especially those related to parenting and education. Children who have mothers who are poorly educated may find it difficult to get information, so children tend to experience delays in growth, due to parenting style (Nurmalasari *et al.*, 2020).

The level of education, especially the mother, affects the health of her family, including the nutritional condition of family members. Maternal education also affects parenting for children because mothers are the main food managers at home, so it has a great influence on the nutritional status of all family members (Noviyanti, Rachmawati, and Sutajo, 2020).

Stunting is a picture of chronic undernutrition status at the time most important for growth and development in infancy. Stunting is caused by various causes, not only malnutrition that experiences children under five and pregnant women (Ministry of Health of the Republic of Indonesia, Directorate General of Community Nutrition, 2018).

Figure 8 found that there was no relationship between the maternal's education level and the incidence of stunting in toddlers at the Madiun City Health Center, as evidenced by a significance value of

0.290 greater than 0.05. These results show that the level of maternal education is not a factor that causes stunting in toddlers at the Madiun City Health Center. This result is in accordance with research by Suharmianti and Agus (2018) which found no relationship between maternal education level and stunting status. Mothers who have a low level of education may or may not have knowledge about nutrition. Since a high level of education does not mean the mother has sufficient knowledge about healthy nutrition, a high level of maternal education does not guarantee that her child will not suffer from malnutrition. Mothers with low education diligently participate in posyandu and toddler nutrition counseling in order to gain the necessary knowledge and be able to take good care of their children (Suharmianti and Agus, 2018).

Based on research conducted by Satoto in 1997, this could be because the height/age indicator reflects previous nutritional history and is less sensitive to changes in nutritional inputs, where in this case the mother plays a role in the distribution of nutritional inputs. Height can only increase or stay at a certain moment, unlike weight, which can increase, decrease, or remain constant. Under normal circumstances, height grows with age. In addition, there is no relationship between maternal education and the incidence of stunting because the maternal's education level is not the only factor causing the problem of malnutrition because there are many other factors that can influence the occurrence of malnutrition problems.

Based on the data obtained by researchers, there are mothers who have a low level of education but have toddlers who are not stunted, and vice versa mothers who have a high level of education but have toddlers who are stunted due to one factor, namely the sex of the toddler itself. Based on other data obtained, it is known that mothers who have a low level of education but have toddlers who are not stunted are more female than male. Conversely, mothers who have a high level of education but have stunted toddlers are more male than female.

Baby girls have less muscle and more fat than baby boys. Muscle burns more energy than fat so men and women of the same height, weight, and age have different body compositions, which means their energy and nutrient needs will also be different (Almatsier, 2004).

In general, male toddlers are more active than female toddlers. Male toddlers often engage in more outdoor activities, such as running around, which makes them more likely to come into contact with dirty environments and expend more energy when their intake is restricted (Christin Angelina F., Agung Aji Perdana, and Humairoh. 2019).

There is a relationship between nutritional status and sex because gender determines the amount of nutrition a person needs. The difference in body composition between women and men affects the large gap in nutritional needs so that it requires even greater intake needs (Febriani et al., 2018).

Figure 9 showed that there was no relationship between family income level and the incidence of stunting in toddlers at the Madiun City Health Center, as evidenced by a significance value of 0.628 greater than 0.05. The results of this study are the same as research conducted by Dakhi in 2019 which stated that family income is not a risk factor for stunting. This can be because the income collected is used for various other purposes, not only to buy food. Because high income does not necessarily indicate that the money set aside is sufficient for food needs and is not a guarantee that toddlers will have a good nutritional status (Dakhi, 2019).

Based on the data obtained by researchers, there are families with low education levels but have toddlers who are not stunted, and vice versa there are families with high income levels but have toddlers who are stunted due to one of the factors, namely the maternal's occupation. Based on other data obtained, it is known that families with low income levels but have toddlers who are not stunted mostly have mothers who do not work. In contrast, families with high income levels but stunted toddlers mostly have working mothers.

A number of articles discussing the characteristics of mothers' employment status state that 75% of working mothers have children who are stunted. The majority of mothers under five in mountainous areas are farmers, so more children are stunted. This is related to non-exclusive breastfeeding because wives who have farmer husbands will help to work in the garden, especially at the beginning of the growing season. As mothers work more in the garden, mothers will leave their children to the care of others while they work. Therefore, before the baby is 6 months old the majority of mothers stop breastfeeding their babies (Leo et al, . 2018).

As many as 66% of unemployed mothers have normal children or are not stunted because mothers are always involved in the childcare process and are related to good parenting. Working mothers have a positive impact on income growth, but on the other hand have a negative impact on child development and maintenance (Syahida, 2019).

Figure 10 showed that there was no relationship between maternal education level and family income with the incidence of stunting in toddlers at the Madiun City Health Center, as evidenced by a significance value of 0.545 greater than 0.05. As previously explained, the results of this study are in accordance with research conducted by Erfince and Minarni (2020) which found no relationship between maternal education level and stunting incidence and research conducted by Grace, et al (2019) which stated that there was no relationship between income level and stunting incidence.

There is no relationship between the maternal's education level and stunting nutritional status because maternal education is not a guarantee that mothers have more knowledge related to nutrition. From direct observation, mothers who have a low level of education tend not to work, thus giving more time in the morning to go to the posyandu every day to get additional food and nutrition and health counseling (Erfince & Minarni, 2020).

There is no relationship between family income level and stunting because families with income below MSEs can manage nutritious and healthy food with easy and affordable products so that during pregnancy mothers can meet nutritional needs and prevent stunting. High family income is not necessarily allocated enough for food needs so it is not a guarantee that the mother will have a good nutritional status. The study did not find any relationship between income and stunting prevention behavior during pregnancy because families with incomes below MSEs can still meet their family's food needs (Niswa, Joni, & Florentina, 2019).

Based on the data obtained by researchers, there are toddlers who have high risk factors (mothers who have low levels of education and income) but have toddlers who are not stunted, and vice versa toddlers who have either (high education level with low income levels or vice versa) or all high risk factors (mothers who have high levels of education and income) but have toddlers who are stunted due to several factors i.e. the sex of the toddler and the maternal's occupation.

Based on other data obtained, it is known that toddlers who have high risk factors (mothers who have low levels of education and income) but have toddlers who are not stunted are more female than men and mothers who are not working than working mothers. Conversely, toddlers who have either (higher education level with low income level or vice versa) or all high risk factors (mothers who have high education and income levels) but have stunted toddlers are more male than female and working mothers than non-working mothers.

Baby girls have less muscle and more fat than baby boys. Muscle burns more energy than fat so men and women of the same height, weight, and age have different body compositions, which means their energy and nutrient needs will also be different (Almatsier, 2004).

In general, male toddlers are more active than female toddlers. Male toddlers often engage in more outdoor activities, such as running around, which makes them more likely to come into contact with dirty environments and expend more energy when their intake is restricted (Christin Angelina F., Agung Aji Perdana, and Humairoh. 2019).

There is a relationship between nutritional status and sex because gender determines the amount of nutrition a person needs. The difference in body composition between women and men affects the large gap in nutritional needs so that it requires even greater intake needs (Febriani et al., 2018).

Working mothers are associated with non-exclusive breastfeeding because wives who have farmer husbands will help to work in the garden, especially at the beginning of the growing season. As mothers work more in the garden, mothers will leave their children to the care of others while they work. Therefore, before the baby is 6 months old the majority of mothers stop breastfeeding their babies (Leo et al., 2018).

As many as 66% of unemployed mothers have normal children or are not stunted because mothers are always involved in the childcare process and are related to good parenting. Working mothers have a positive impact on income growth, but on the other hand have a negative impact on child development and maintenance (Syahida, 2019).

CONCLUSION

Based on the results of the research and discussion above, several things can be concluded as follows:

1. The results found that 50% of toddlers who suffer from stunting and those who do not suffer from stunting as much as 50%.
2. At the Madiun City Health Center, most toddlers have a female gender.
3. Most toddlers at Puskesmas Kota Madiun have working mothers.
4. At the Madiun City Health Center, most toddlers have mothers with higher education.
5. Most of the families under five in Puskesmas Kota Madiun have income from \geq MSEs in Madiun City.
6. From the results of the study, it was found that there was no relationship between the level of maternal education and the incidence of stunting in toddlers at the Madiun City Health Center.

7. From the results of the study, it was found that there was no relationship between the level of family income and the incidence of stunting in toddlers at the Madiun City Health Center.
8. From the results of the study, it was found that there was no relationship between the level of maternal education and family income with the incidence of stunting in toddlers at the Madiun City Health Center.

REFERENCE

- Agustin, L., & Rahmawati, D. 2021. The Relationship of Family Income with the Incidence of Stunting. Indonesian Journal of Midwifery (IJM), 4(1), 30. <https://doi.org/10.35473/ijm.v4i1.715>
- Akmal, Y., Hikmah, H., Subekti, I., & Hardono, I. H. 2019. Strategy for decreasing the rate of stunting through early childhood health and nutrition training for tutors/parents of early childhood education. Journal of Obsession: Journal of Early Childhood Education, 4(1), 454. <https://doi.org/10.31004/obsesi.v4i1.302>
- Anasari, T., & Suryandari, A. E. 2022. The Relationship between History of Hypertension and Birth Spacing with the Incidence of Stunting. Journal of Bina Cipta Husada, 18(1), 107-117.
- Angelina F., Christin., Prime, Agung Aji., & Humairoh. 2019. Factors of *Stunting* in Lampung Province. World Journal of Public Health. 7 (3), 130.
- Anna Ratnawati. 2020. Maternity Nursing Care. New library Pres.Yogyakarta
- Astutik, Rahfiludin, M. Z., & Aruben, R. 2017. RISK FACTORS FOR STUNTING IN CHILDREN UNDER THE AGE OF 24-59 MONTHS (Case Study in the Working Area of Puskesmas Gabus II Pati Regency in 2017). Journal of Public Health, 6 (1), 2356-3346.
- Azriful, A. et al. 2018. 'Determinants of Stunting Incidence in Toddlers Aged 24-59 Months in Rangas Village, Banggae District, Majene Regency', Al-sihah: The Public Health Science Journal, 10(2), pp. 192– 203. DOI: 10.24252/AS.V10I2.6874.
- Bardosono, S. 2019. Assessment of Nutritional Status of Toddlers (Anthropometry). Journall Faculty of Medicine UI.
- Burhanudin, M., Istiyani, N., & Widjajanti, A. 2015. Factors Affecting Family Income in Banyuwangi District, Banyuwangi Regency. Student Scientific Articles, 1(1), 1–6.
- Candra, A., Science, B., Faculty, G., University, K., & Semarang, D. 2020. Pathophysiology of Stunting. 8(2), 74–78.
- Cheikh Mbacké Faye, et all. 2018. Factors Associated With Recovery From Stunting Among Under-Five Children In Two Nairobi Informal Settlements. <https://doi.org/10.1371/journal.pone.0215488>.

[13/11/2022]

- Cintya, & Dewi, R. 2015. *Baby Growth Theory & Concept, Toodler; Children and Youth*. Yogyakarta: Nuha Medika
- Dakhi A. 2019. The Relationship between Family Income, Education, and Maternal Knowledge about Nutrition with the Incidence of Stunting in Children Aged 6-23 Months in the Working Area of the Jati Makmur Health Center in North Binjai. *J Health Masy Indones*. VIII:3–77.
- Darmini, N. W., Fitriana, L. B., & Vidayanti, V. 2022. The relationship between the level of maternal knowledge about balanced nutrition and the incidence of stunting in toddlers aged 2-5 years. *Coping: Community of Publishing in Nursing*, 10(2), 160. <https://doi.org/10.24843/coping.2022.v10.i02.p06>
- Goddess, I. 2019. Factors affecting the incidence of stunting in toddlers 24-60 months in the Working Area of the Lakudo Health Center, Central Buton Regency. *Scientific Journal of Health Diagnosis*, 14(1), 85-90.
- Fakhma and Dhevi. 2020. Analysis of factors related to the nutritional status of toddlers in the working area of the South Paringin Health Center, Balangan Regency in 2020. Islamic University of Kalimantan. <http://eprints.uniska-bjm.ac.id/2322/>
- Febriani, C. A., Prime, A. A., & Humairoh, H. 2018. Factors in the incidence of stunting of toddlers aged 6-23 months in Lampung Province. *World Journal of Public Health*, 7(3).
- Febrianti, Y. 2020. Overview of Family Economic Status on Nutritional Status of Toddlers (BB/U) in Rumbai Pesisir District, Pekanbaru City. Thesis, 2(1), 5–7. <http://jurnal.globalhealthsciencegroup.com/index.php/JPPP/article/download/83/65%0Ahttp://www.embase.com/search/results?subaction=viewrecord&from=export&id=L603546864%5Cnhttp://dx.doi.org/10.1155/2015/420723%0Ahttp://link.springer.com/10.1007/978-3-319-76>
- Fikawati, Sandra, et al. 2017. *Child and adolescent nutrition*. Ed. 1. Cet. 1. Depok : Rajawali Press.
- Habimana, S., & Biracyaza. 2019. Risk Factors Of Stunting Among Children Under 5 Years Of Age In The Eastern And Western Provinces Of Rwanda: Analysis Of Rwanda Demographic And Health Survey 2014/2015. *Pediatric* <https://doi.org/10.2147/phmt.s222198>
- Hadisuyitno, J., & Riyadi, B. D. 2021. Determining Factors of Toddler Stunting in Batu City, Indonesia. 12(1), 231–234.
- Hariyani Sulistyoningih, S., & KM, M. 2020. THE RELATIONSHIP BETWEEN PARITY AND EXCLUSIVE BREASTFEEDING WITH STUNTING IN TODDLERS (LITERATURE REVIEW). In *JOURNAL OF NATIONAL SEMINARS* (Vol. 2, No. 01, pp. 1-8).
- Harjatmo TP, Par'i HM, Wiyono S. 2017. *Textbook for Nutritional Status Assessment*. Jakarta: Ministry of

Health of the Republic of Indonesia.

Hondro, I. H. 2021. Overview of mother's knowledge about stimulation of fine motor development in toddlers 0-5 years old in Lololakha Village. *Scientific Papers*, 1(1), 1–20.

Husnaniyah, D., Yulyanti, D., & Rudiansyah, R. 2020. The Relationship Between Maternal Education Level and the Incidence of Stunting. *The Indonesian Journal of Health Science*, 12(1), 57–64. <https://doi.org/10.32528/ijhs.v12i1.4857>

Ika, L., & Ariati, P. 2019. RISK FACTORS CAUSE OF STUNTING IN TODDLERS AGED 23-59 MONTHS. *VI*(1), 28–37.

Imani, N. 2020. *Stunting in Children: Recognize and Prevent Early Childhood*. Hijaz Independent Library.

Beautiful Budiastutik, & Muhammad Zen Rahfiludin. 2019. Risk Factors for Stunting in Children in Developing Countries. *Amerta Nutrition*, 3(3), 122–129. <https://doi.org/10.2473/amnt.v3i3.2019.122-129>

Istiany, Ari & Rusilanti. 2013. *Applied Nutrition*. Bandung: PT Remaja Rosdakarya.

Jayanti, R., & Ernawati, R. 2021. Pregnancy Distance Factors Associated with Stunting Events at Puskesmas Harapan Baru Samarinda Seberang. *Borneo Student Research (BSR)*, 2(3), 1705-1710.

Juwita, S., Andayani, H., Bakhtiar, B., Sofia, S., & Anidar, A. 2019. The Relationship between Total Family Income and Completeness of Basic Immunization with the Incidence of Stunting in Toddlers in Pidie Regency. *Nanggroe Medika Medical Journal*, 2(4), 1-10.

Ministry of Health of the Republic of Indonesia Directorate General of Nutrition Masyarakat. 2018. *Pocket Book of Nutritional Status Monitoring Results*. Jakarta.

Khoeroh, H., & Indriyanti, D. 2017. Evaluation of Nutrition Management of Stunting Toddlers in the Working Area of the Sirampog Health Center. *Unnes Journal of Public Health*, 6 (3), 2252-6781.

Kusumaningtyas, D. E., & Deliana, S. M. 2017. Feeding Patterns on Nutritional Status of 12-24 Months of Age in Working Mothers Abstract. 2(89), 155–167.

Langi, Grace K.L., Day Two, Vera T., Purba, Rudolf B., & Pelanginang, Janeke I. 2019. Nutritional Intake and Family Income Level on the Incidence of Stunting in Children Aged 3-5 Years. Manado: Department of Nutrition Health Polytechnic Ministry of Health Manado, 11(2).

Leo, A. R., Subagyo, H. W., & Kartasurya, M. I. 2018. Risk factors for stunting in children aged 2-5 years in mountain and coastal areas. *J.Gipas*, 2(1), 51-63

Lutviana, E., & Budiono, I. 2019. Prevalence and determinants of undernutrition events in toddlers. *PACKAGING: Journal of Public Health*, 5(2).

- Madina, T. 2019. The Effect of Family Income on Household Consumption Behavior in an Islamic Perspective: A Case Study of Ilir Timur II District, Palembang. *Journal of Sharia Economic Thought and Development*, 4(2), 15–24. <https://ejournal.stebisigm.ac.id/index.php/esha/article/view/153>
- Maharani, S. D. S., Wulandari, S. R., & Melina, F. 2018. The relationship between the incidence of stunting and development in toddlers aged 3-5 years at the Kricak posyandu Yogyakarta. *Health Scientific Journal*, 7(1), 32–36. <https://doi.org/10.52657/jik.v7i1.1095>
- Mentari, S., & Hermansyah, A. 2018. Factors Related to the Stunting Status of Anan Age 24-59 Months in the Working Area of UPK Puskesmas Siantan Hulu. *Pontianak: Pontianak Nutrition Journal (PNJ)*, 01 (01).
- Metwally, A. M., El-Din, E. M. S., Shehata, M. A., Shaalan, A., El Etreby, L. A., Kandeel, W. A., Shaaban, S. A., and Rabah, T. M. 2016. Early life predictors of socio-emotional development in a sample of Egyptian infants', *PLoS ONE*, 11(7): 1–17. doi:10.1371/journal.pone.0158086.
- Mufdlilah. 2017. *Guidebook for Empowering Breastfeeding Mothers in the Exclusive Breastfeeding Program*. Yogyakarta
- Mugianti, S., Mulyadi, A., Anam, A. K., & Najah, Z. L. 2018. Factors Causing Stunting Children Aged 25-60 Months in Sukorejo District, Blitar City. *Journal of Ners and Midwifery*, 5(3), 268–278. <https://doi.org/10.26699/jnk.v5i3.art.p268-278>
- Mustamine. 2018. Maternal Education Level and Exclusive Breastfeeding with the Incidence of Stunting in Toddlers in South Sulawesi Province. *Food Nutrition Media*. 25(1)
- Nasution, D. 2014. The relationship between low birth weight (BBLR) and the incidence of stunting in children aged 6-24 months in Yogyakarta City. Thesis. Gadjah Mada University. Downloaded from <https://repository.ugm.ac.id/id/eprint/129665>
- Ni'mah, C., and Muniroh, L. The relationship between education level, knowledge level and maternal parenting with wasting and stunting in poor toddlers. *Indonesian Nutrition Media*. 10(1): 84-90.
- Ngaisyah, R. D. 2015. Socioeconomic relationship with the incidence of stunting. *Journal of Medical Respati*.10(4): 65– 70.
- Notoatmodjo, Soekidjo. 2018. *Health Research Methodology*. Jakarta: PT Rineka Cipta
- Noviyanti, L.A., Rachmawati, D.A. and Sutejo, I.R., 2020. An Analysis of Feeding Pattern Factors in Infants at Kencong Public Health Center. *JOURNAL AMS*, 6(1): 14-18.
- Nurdin, Ismail and Sri Hartati. 2019. *Social Research Methodology*. Surabaya: Media Friends of Scholars
- Nurmalasari, Y., Anggunan, & Febriany, T. W. 2020. The relationship between maternal education level and family income with the incidence of stunting in children aged 6-59 months. *Journal of*

Midwifery, 6(2): 205-211.

- Palino, Inochi, et al. 2017. Determinants of stunting incidence in toddlers aged 12-59 months in the working area of Puuwatu Health Center, Kendari City in 2016. *Student Scientific Journal of Public Health*. Volume 2, Number 6
- Pongreku, P. S, Sunarsih and Fatmawati. 2020. Factors Associated with the Incidence of Stunting in South Konawe District. *Scientific Journal of Midwifery Stikes Mandala Waluya* Vol 6, No 2 of 2020.
- Rahayu, Seni et al. 2019. The relationship of knowledge, attitudes, behaviors and characteristics of mothers about exclusive breastfeeding to the nutritional status of infants. *AcTion Journal: Aceh Nutrition Journal*, Volume 4, Number 1.
- Rahayu, A., Yulidasari, F., Putri, A. O., & Anggraini, L. 2018. Study Guide - Stunting and Prevention Efforts. Stunting books and prevention efforts. Yogyakarta: CV Mine. Taken from http://kesmas.ulm.ac.id/id/wpcontent/uploads/2019/02/BUKU-REFERENSI-STUDY-GUIDESTUNTING_2018.pdf
- Rahman, A., Munandar, S. A., Fitriani, A., Karlina, Y., & Yumriani. 2022. Understanding Education, Education Science and Elements of Education. *Al Urwatul Wutsqa: Islamic Education Studies*, 2(1), 1–8.
- Rahmawati, N. F., Fajar, N. A., & Idris, H. 2020. stunting of toddlers of poor families receiving PKH in Palembang. 17(1), 23–33. <https://doi.org/10.22146/ijcn.49696>
- Rita Kirana, Aprianti, N. W. H. 2022. The influence of health promotion media on maternal behavior in stunting prevention during the Covid-19 pandemic (in kindergarten children of Kuncup Harapan Banjarbaru). *Journal of Research Innovation*, 2(9), 2899–2906.
- Rusdi, P. H. N. 2021. The relationship between the provision of nutrition and environmental sanitation to the incidence of stunting in toddlers. *Human Care Journal*, 6(3), 731-736.
- Salamung, N, Haryanto, J., & Sustini, F. 2019. Factors Related to Stunting Prevention Behavior During Pregnancy in the Working Area of the Bondowoso Regency Health Center. Surabaya : *Forikes Sound Health Research Journal*, 10(4).
- Salsabila, S., Dewi Noviyanti, R., Pertiwi, D., Kusudaryati, D., & Abstract, K. K. 2022. The Relationship Between Maternal Education Level and Parenting Style with the Incidence of Stunting in Toddlers Aged 12-36 Months in the Sangkrah Health Center Area. *PROFESSION (Islamic Professional): Research Publication Media*, 19(2), 143–151.
- Sanah Nor. 2017. Implementation of the function of puskesmas (community health center) in improving the quality of health services in Long Kali District, Paser Regency. *EJournal Government Sciences*, 5(1), 305–314.

- Sari, I. Y. 2016. Food Consumption and Nutritional Status of Children Under Five (24 – 59 Months) in Puger Wetan Fisherman Village, Puger District, Jember Regency [Jember University]. In Jember University. <https://repository.unej.ac.id/handle/123456789/77677>
- Sari, S. D., & Zelharsandy, V. T. 2022. The Relationship of Family Economic Income and Maternal Education Level to the Incidence of Stunting. *Journal of Midwifery Hope Ibu Pekalongan*, 9(2), 108–113. <https://doi.org/10.37402/jurbidhip.vol9.iss2.200>
- Sebayang, Sofia., Rajagukguk, T. 2019. The Effect of Education, Training and Work Motivation on Teacher Performance in Private Elementary and Junior High School Budi Murni 3 Medan. *METHONOMIX Journal of Management Science*, 2(2), 105–114.
- Setiawan, Y.A. 2019. Relationship Of Mother Factors And Stunting Incidence In Children (24-59 Months) In Buniwangi Village, Work Area Of Pagelaran Public Health Center, Cianjur Regency. *International Seminar on Global Health*. 3(1) : 115-123.
- Setyaningsih A. 2020. Factors Affecting the Occurrence of Stunting in the Working Area of Wanareja 1 Health Center, Cilacap Regency in 2020. Thesis of Public Health Study Program STIKes Bina Cipta Husada Purwokerto.
- Setyo, C., Andhini, D., Nurfajriyani, I., Sadiya, H., & Nurpuji, F. 2022. Stunting Prevention Efforts through Increasing Maternal Knowledge at the Pulasaren Health Center in Cirebon City. 1(4), 381–386.
- Soetjningsih. 2012. *Child Development and Its Problems in Textbook I of Child and Adolescent Development Science*. Jakarta :Sagungseto
- Suci, I., Puspa, M. B., Putri, M. A., & Fatmarani, M. 2019. The Difference Between Hospital and Puskesmas. *Journal of Health*.
- Suryansyah. 2012. Portion Feeding for Infants and Toddlers, <http://Health.detik.com/dokter/768>
- Sutanto, Andina Vita and Fitriana, Yuni. 2018. *Upbringing in pregnancy*. Yogyakarta: New Library.
- Sutarto, Azqinar, T. C., Himayani, R., and Wardoyo. 2020. The Relationship between Maternal Education Level and Family Income with the Incidence of Stunting in Toddlers in the Working Area of the Way Urang Health Center, South Lampung Regency. *World Journal of Public Health*. 9(2): 256-263. <http://ejournalmalahayati.ac.id/index.php/duniakesmas/index>
- Shahida, A. 2019. The Relationship between Education and Parental Work with the Growth of Toddlers in Ingin Jaya Village, Rantau District, Aceh Tamiang Regency in 2018. *Journal of Edukes*, 2(1), 18-28.
- Tongkonoo, I. 2021. The Relationship of Socioeconomic and Environmental Factors with the Incidence of Stunting in Valita Aged 24-49 Months at the Gorontalo Regency Health Center. Thesis, 1(702518006).

- Wanimbo, Erfince, and Wartiningsih, Minarni. 2020. The Relationship between Maternal Characteristics and the Incidence of Baduta Stunting (7-24 Months). Surabaya : Journal of Health Management RS. Dr. Soetomo, 6(1).
- Wati, S. P. 2018. The relationship between education level, maternal knowledge and parental income with the nutritional status of children under the age of 1-5 years in Duwet Village, Wonosari District, Klaten Regency. *University of Muhammadiyah Surakarta*, 1–20.
- Wicaksono, K. E., & Alfianto, A. G. 2020. The positive impact of health education on the level of family knowledge in nutrition management of stunted toddlers. Conference on Innovation and Application Of Science and Technology, 981-986.
- Yanti, T., & Fauziah, D. 2021. The Effect of Family Income on Stunting Incident in Preschool Children at Bogor City During COVID-19 Pandemic. *Indian Journal of Public Health Research & Development*, 12(2), 407–412. <https://doi.org/10.37506/ijphrd.v12i2.14167>
- Yulaikhah, L. 2019. Teaching Book of Obstetric Care of Pregnancy. In *Journal of Chemical Information and Modeling* (Vol. 53).
- Yuliana, E. 2017. Analysis of Student Knowledge About Healthy and Nutritious Food on School Snack Selection. Retrieved from http://repository.ump.ac.id/4114/3/Erlin%20Yuliana_BAB%20II.pdf on November 20, 2022 at 16:17 WIB.
- Yulistiana, Evayanti, 2015. The relationship between maternal knowledge and husband support for pregnant women on the regularity of antenatal care (ANC) visits at the Wates Health Center in Central Lampung in 2015. *Journal of Midwifery* Vol 1, No 2, July 2015: 81-90
- Yunianto, A. E., Fitri, Y., Wagustina, S., Fitriarningsih, E., & Mulyani, N. S. 2020. Mother's height and calcium intake against stunting among children aged 3-5 years and the impact on child development. *Sys Rev Pharm*, 11(10), 606–611.