

Routine Hematology Description and D-dimer COVID-19 Patients at Sidoarjo Regional Hospital

by Turnitin cek4

Submission date: 14-Jun-2024 07:32PM (UTC+0700)

Submission ID: 2199228593

File name: and_D-dimer_COVID-19_Patients_at_Sidoarjo_Regional_Hospital.pdf (304.39K)

Word count: 3675

Character count: 20197

ARTICLE INFO

29 **THOR'S AFFILIATIONS**

Department of Pharmacology,
 Medical Faculty, Universitas
 Wijaya Kusuma Surabaya^{1,2}
 Faculty of Public Health,
 Universitas Muhammadiyah
 Gresik³

29 **RESPONDING AUTHOR**

Nugroho Eko Wirawan Budianto
 Department of Pharmacology,
 Medical Faculty, Universitas
 Wijaya Kusuma Surabaya.

E-mail:

nugrohoewb@uwks.ac.id

Article history

Received 13-01-2024
 Revised 08-03-2024
 Accepted 22-03-2024
 Available online

Please cite this article in APA 7th edition style as:

Budianto, N. E. W, Setiawan, B., Budiono, N. D. P. (2024). Routine hematology description and D-dimer COVID-19 patients at Sidoarjo Regional Hospital. *Jurnal Ilmiah Kedokteran Wijaya kusuma*, 13(1), XX-XXX
<https://doi.org/10.30742/jikw.v13i1.3537>

Routine Hematology Description and D-dimer COVID-19 Patients at Sidoarjo Regional Hospital

Nugroho Eko Wirawan Budianto^{1*}, Budhi Setiawan², Nugrahadi Dwi Pasca Budiono²

Abstract

Background: COVID-19, caused by the SARS-COV-2 virus, is a contagious and potentially fatal disease that is commonly diagnosed using real-time reverse polymerase chain reaction (RT-PCR) assay. Studies have shown that the D-dimer level and the features of hematological exams can offer important prognostic value of the patients. This study aimed to describe data about the routine hematology features and D-Dimer levels of hospitalized COVID-19 patients in Sidoarjo Regional Hospital. **Methods:** This study used a retrospective, cross sectional, observational approach with descriptive analysis. Medical records of 167 COVID-19 patient were collected based on the criteria. The outcomes of the D-dimer and complete blood count of the patients were summarized. **Results:** According to the findings, the number of the patients with COVID-19, 53 (31.7%) were between the ages of 31 and 35, and at least 36 (21.6%) were between the ages of 17 and 30. Prominent hematology routine outcomes were the number of patients with anemia (103 or 61.7%), neutrophilia (105 or 62.9%), and lymphocytopenia (129 or 77.2%). The D-dimer value increased by 93 (55.7%) patients among 167 COVID-19 patients. **Conclusion:** Therefore, monitoring the patient's routine hematological presentation and D-dimer level during treatment is essential for preventing mortality and morbidity.

Keywords: COVID-19, D-dimer, SARS-CoV-2, routine hematology

Original Research Article

INTRODUCTION

COVID-19 infection is a fatal respiratory disease if untreated. The initial name of betacoronavirus disease was SARS-COV-2 Coronaviruses (COV) and the 2019 Coronavirus Novel (2019-nCov). Nevertheless, the World Health Organization (WHO) changed its name to COVID-19, or Coronavirus Disease 2019, instead. The number of cases continues to increase since it first appeared in the city of Wuhan, China, at the end of 2019. It is estimated that the first person infected in Indonesia on March 2, 2020, was a tourist (Handayani et al., 2020).

Studies have demonstrated the prognostic significance of D-dimer levels, which have emerged as a key predictor in COVID-19 patients. discovered a relationship between hsCRP and D-dimer levels in COVID-19 patients, specifically in relation to inflammatory markers (Y. Li et al., 2021; Naymagon et al., 2020; Yu et al., 2020). D-dimmer is a sign of coagulopathy that is often found in COVID-19 patients and is related to the severe condition of the disease (Ardiani et al., 2023). This is because the patient's

D-dimer can increase because of the innate disease carried by the patient (Rostami & Mansouritorghabeh, 2020). D-dimer is known as a strong indicator of morbidity and mortality among COVID-19 patients (Gungor et al., 2021).

The RT-PCR assay and routine hematological profile screening are particularly useful tools for physicians in identifying COVID-19. Numerous COVID-19 studies have employed these tests for disease screening and surveillance (Fitriani, 2020; Susilo et al., 2020; Wu et al., 2020). Leucopenia, lymphopenia, and increased neutrophil counts are a few hematological presentations that may raise the possibility of COVID-19, as well as the quantity of atypical leukocytes (Sabatudung et al., 2023).

Decreased albumin level and increased lactate dehydrogenase (LDH), SGOT, blood sedimentation rate (LED), and C-reactive protein (CRP) levels are also some laboratory abnormalities found in COVID-19 patients. Therefore, the purpose of this study was to assess the D-dimer levels and routine hematologic characteristics of hospitalized COVID-19 patients at Sidoarjo Regional Hospital.

6

MATERIALS AND METHODS

This study was a retrospective cross-sectional secondary data collection study using a quantitative descriptive approach. Secondary data were retrieved from the paper medical record database of the Sidoarjo Regional Hospital, Sidoarjo, Indonesia. Paper-based medical records of patients older than 18-year who were clinically diagnosed with COVID-19 were purposefully selected for further analysis. Administrative, physical examination, and laboratory data of the patients were collected in the excel spreadsheet template. Data on hospitalized COVID-19 patients were collected between May and August 2021. Inclusion criteria involved the COVID-19 patient's medical records that showed positive PCR assay and had both, the results of D-dimer test and routine hematology examination. The exclusion criteria were pregnancy and severe comorbidities such as renal impairment, hepatic diseases, and cardiovascular disorders.

15

RESULTS

Table 1. Characteristics of Patients Based on Age and Gender

Age (year)	Male	Female	%
17-30	18	18	21.6
31-45	25	28	31.7
46-59	20	17	22.1
≥60	24	17	24.6
TOTAL	87	80	100

Table 1 shows the 167 COVID-19 patients who fulfilled the requirements for inclusion. According to table 1, 52.1% of all patients (87 individuals) were male. The highest percentage of patients was 53 (31.7%) in the age group 31 to 45 years, and the lowest percentage was 36 (21.6%) in the age group 17 to 30 years.

Table 2. Patient's Routine Hematology Parameters

Parameter	Results	Total (n)	Percentage (%)
Erythrocyte (RBC)	Decrease	22	13.2
	Normal	143	85.6
	Increase	2	1.2
Hemoglobin (HGB)	Decrease	103	61.7
	Normal	64	38.3
	Increase	0	0

Leucocyte (WBC)	Leukopenia	19	11.4
	Normal	122	73
	Leukocytosis	26	15.6
Monocyte	Monocytopenia	6	3.6
	Normal	106	63.5
	Monocytosis	55	32.9
Neutrophil	Neutropenia	13	7.8
	Normal	49	29.3
	Neutrophilia	105	62.9
Lymphocyte	Lymphocytopenia	129	77.2
	Normal	26	15.6
	Lymphocytosis	12	7.2
Eosinophil	Eosinopenia	0	0
	Normal	160	95.8
	Eosinophilia	7	4.2
Thrombocyte (PLT)	Thrombocytopenia	24	14.4
	Normal	122	73
	Thrombocytosis	21	12.6

Table 2 displays the percentage of COVID-19 patients who had hemoglobin and lymphocyte test results that are higher than normal: 103 (61.7%) and 129 (77.2%), respectively. Furthermore, a substantial number of 129 (77.2%) COVID-19 patients had neutrophil test values that were greater than normal. The results of the tests for erythrocytes, leucocytes, monocytes, eosinophils, and thrombocytes among all the patients indicated that the percentage of patients with normal values was higher than that with abnormal values.

According to table 3, there were 93 patients (55.7%) whose D-dimer level was abnormal or higher than 500 ng/dL. Male and female distributions were comparable, based on D-dimer concentration findings.

Table 3. Gender Overview Based on D-dimer's examination

Gender	Normal	Abnormal (>500 ng/dL)
Male	40	47
Female	34	46
Total	74	93

Table 4. Overview of the results of D-dimer's examination based on age

Age (year)	Abnormal (>500 ng/dL)	Normal
17-30	18	18
31-45	24	29
46-59	20	17
≥60	31	10
Total	93	74

Table 4 shows that out of 167 COVID-19 patients, 93 patients (55.7% of the total) had D-dimer concentrations greater than 500 ng/dL, whereas 74 patients had normal D-dimer levels. Patients who were 60 years of age or older had the highest number (31 or 18.6%), while those in the 17–30 age range had the lowest (18 or 10.8%).

DISCUSSION

D-dimer analysis and routine complete blood counts (CBC) are crucial for the risk classification of COVID-19 patients. In a similar vein, the significance of hematological findings in COVID-19 was highlighted by the systemic effect of the virus on the hematopoietic system and hemostasis (Terpos et al., 2020). Laboratory results of COVID-19 patients with severe symptoms showed an increase in CRP, increased D-dimer, prothrombin time extension, lymphocytopenia, procalcitonin, neutrophilia, albumin, leukopenia, and LDH. In addition, several studies have shown that patients with COVID-19 can experience anemia and thrombocytopenia (Christensen et al., 2020; Shama et al., 2023).

The study findings indicated that while the proportion of COVID-19 patients with anemia was quite high (103, or 61.7%), the majority of patients (143, or 85.6%) had normal erythrocyte counts. The difference in the normal range of the parameters and incidence rate between males and females may be the cause of this inconsistency. In addition, if most of the patients had mild anemia and were only marginally different from the cutoff value, this specific condition might be possible (Bergamaschi et al., 2021).

The interaction between the Sars-COV-2 virus and hemoglobin molecules can occur via erythrocyte receptors, specifically CD147 and CD26. Hemoglobinopathy, which results from this interaction, leads to hemolysis and hemoglobin malfunction. Moreover, ferroportin, which produces hyperferritinemia, and ferroptosis, which causes iron dysmetabolism, are inhibited by the viral hepcidin mimetic effect (Cavezzi et al., 2020). The interpretation of RBC, HGB, and MCV can be normal or slightly lowered in the case of a corona virus infection. The average volume of the corpuscular (MCV), also referred to as the average volume of erythrocytes and hemoglobin, and the number of erythrocytes, or red blood cells (RBC), are known to be correlated (C. Li et al., 2020). Hemolysis, or a drop in hemoglobin, may result from the virus's attack on the heme in the 1-beta hemoglobin chain. This could lead to anemia (Al-kuraishy et al., 2022). Patients with severe COVID-19 disease have a considerably lower hemoglobin values than those with milder variants (Lippi & Mattiuzzi, 2020).

In this study, the number of patients with neutrophilia (105, or 62.9%), and lymphocytopenia (129, or 77.2%) were the prominent outcomes for routine hematology. The neutrophil-to-lymphocyte ratio (NLR) characteristics are derived from the number of neutrophils and lymphocytes. Individuals with a mild prognosis have lower NLR scores than those who are terminally ill and cannot survive. The strength of inflammatory responses is shown by the increase in neutrophils, and immune system damage is indicated by the decline in lymphocytes (Prozan et al., 2021). The controlled immune cell response modifies the virus's inflammatory level. Consequently, the inflammatory level may be predicted using the NLR value. It is feasible to evaluate the COVID-19 risk variables using the greater value of the NLR (Liu et al., 2020). Although lymphocytes are only a risk factor for the disease, the value of NLR shows good performance, which can be used for diagnosis and disease prognosis (Asghar et al., 2022; Peng et al., 2020). If the immune system is weak, the virus can enter the body and cause a decrease the number of lymphocytes. Because a reduction in lymphocyte count may indicate immune system harm, the COVID-19 virus may disrupt the immune system (Mus et al., 2021).

The results of the D-dimer examination in table 3 and table 4 show that 167 patients with COVID-19, there were 93 (55.7%) patients who had abnormal outcomes (higher than 500 ng/dL). According to previous studies, poor radiological imaging and elevated clinical severity are linked to elevated D-Dimer levels (Rostami & Mansouritorghabeh, 2020). Higher D-dimer levels upon hospital admission have been linked to a higher risk of illness and mortality in SARS-COV-2 infected patients (Amelia et al., 2022). Elevated D-dimer levels, illness severity, and mortality were also substantially correlated with older adult age and diabetes (Soni et al., 2020). Furthermore, patients with severe COVID-19 frequently have elevated D-dimer levels, which can indicate the occurrence of ARDS (acute respiratory distress syndrome) will occur (Ryadi et al., 2022).

The result of laboratory examinations in this study showed that the number of the patients with thrombocytopenia was 24 (14.4%). This result differed from that of research done in China in 2020 by Guan et al., who discovered that leukopenia affects 33.7% of patients, thrombocytopenia affects 36.2%

of patients, and lymphocytopenia affects 83.2% of patients (Guan et al., 2020). According to several studies (Guan et al., 2020; Huang et al., 2020; Liu et al., 2020; Zhou et al., 2020), compared with non-severe cases, patients with severe COVID-19 have lower platelet counts. Patients with thrombocytopenia COVID-19 may have disseminated intravascular coagulation (DIC), vascular coagulopathy, significant organ injury, or physiological decompensation as contributing factors (Kaiafa et al., 2022).

This preliminary study was limited in a few ways. Initially, a significant portion of medical records did not fit the inclusion requirements of the study because the patients had various laboratory tests. Second, the study's tiny sample size of medical records and third, the incomplete nature of the test results precluded drawing firm conclusions. This means that more research, including statistical analysis, is required.

CONCLUSION

In summary, data from hospitalized COVID-19 patients revealed a notable presence of anemia, neutrophilia, and lymphocytopenia. It was found that more than 50% of COVID-19 patients had elevated D-dimer concentrations. D-dimer assay and routine hematological examinations are essential components of all-encompassing care for COVID-19 patients.

CONFLICT OF INTEREST

All authors declare that there is no conflict of interest in this study.

ACKNOWLEDGEMENTS

The author would like to thank the research and community service institution to the University of Wijaya Kusuma Surabaya (LPPM) in Surabaya, Indonesia, which supports this work through an internal grant scheme (No. 85/LPPM/UWKS/IV/2022).

REFERENCES

Al-kuraishy, H. M., Al-Gareeb, A. I., Kaushik, A., Kujawska, M., & Batiha, G. E.-S. (2022). Hemolytic Anemia in COVID-19. *Annals of Hematology*, 101(9), 1887–1895. <https://doi.org/10.1007/s00277-022-04907-7>

Amelia, R., Anissa, D. F. N., & Luhulima, D. E. J. (2022). Perbedaan Peningkatan Kadar D-Dimer pada Dewasa dan Lansia Yang Terkonfirmasi COVID-19 di RS Swasta Bintaro. *Journal of Research and Education Chemistry (JREC)*, 4(1), 1–8.

Ardiani, S., Tursinawati, Y., & Wahab, Z. (2023). Hubungan D-Dimer dengan Ketahanan Hidup Pasien Covid-19 Derajat Berat-Kritis di RSUD Tugurejo Semarang. *Health and Medical Journal*, 5(1), 44–49.

Asghar, M. S., Akram, M., Yasmin, F., Najeeb, H., Naeem, G., Gaddam, M., Jafri, M. S., Tahir, M. J., Yasin, I., Mahmood, H., Mehmood, Q., & Marzo, R. R. (2022). Comparative Analysis of Neutrophil to Lymphocyte Ratio and Derived Neutrophil to Lymphocyte Ratio with Respect to Outcomes of Inpatient Coronavirus Disease 2019 Patients: A Retrospective Study. *Frontiers in Medicine*, 9. <https://doi.org/10.3389/fmed.2022.951556>

Bergamaschi, G., Borrelli de Andreis, F., Aronico, N., Lenti, M. V., Barteselli, C., Merli, S., Pellegrino, I., Coppola, L., Cremonese, E. M., Croce, G., Mordà, F., Lapia, F., Ferrari, G., Ballesio, A., Parodi, A., Calabretta, F., Ferrari, M. G., Fumoso, F., Gentile, A., ... Stefani, D. (2021). Anemia in Patients with COVID-19: Pathogenesis and Clinical Significance. *Clinical and Experimental Medicine*, 21(2), 239–246. <https://doi.org/10.1007/s10238-020-00679-4>

Cavezzi, A., Troiani, E., & Corrao, S. (2020). COVID-19: Hemoglobin, Iron, and Hypoxia Beyond Inflammation. A Narrative Review. *Clinics and Practice*, 10(2), 1271. <https://doi.org/10.4081/cp.2020.1271>

17 **rutine hematology description and D-dimer COVID-19 patients at Sidoarjo Regional Hospital**
Nugroho Eko Wirawan Budianto, Budhi Setiawan, Nugrahadi Dwi Pasca Budiono

- 10 Christensen, B., Favaloro, E. J., Lippi, G., & Van Cott, E. M. (2020). Hematology Laboratory Abnormalities in Patients with Coronavirus Disease 2019 (COVID-19). *Seminars in Thrombosis and Hemostasis*, 46(07), 845–849.
- 27 Fitriani, N. I. (2020). Tinjauan pustaka covid-19: Virologi, Patogenesis, dan Manifestasi Klinis. *Jurnal Medika Malahayati*, 4(3), 194–201.
- 4 Guan, W., Ni, Z., Hu, Y., Liang, W., Ou, C., He, J., Liu, L., Shan, H., Lei, C., & Hui, D. S. C. (2020). Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*, 382(18), 1708–1720.
- Gungor, B., Atici, A., Baycan, O. F., Alici, G., Ozturk, F., Tugrul, S., Asoglu, R., Cevik, E., Sahin, I., & Barman, H. A. (2021). Elevated D-Dimer Levels on Admission are Associated with Severity and Increased Risk of Mortality in COVID-19: A Systematic Review And Meta-Analysis. *The American Journal of Emergency Medicine*, 39, 173–179. <https://doi.org/10.1016/j.ajem.2020.09.018>
- 24 Handayani, D., Hadi, D. R., Isbaniah, F., Burhan, E., & Agustin, H. (2020). Corona Virus Disease 2019. *Jurnal Respirologi Indonesia*, 40(2), 119–129.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., ... Cao, B. (2020). Clinical Features of Patients Infected with 2019 Novel Coronavirus in Wuhan, China. *The Lancet*, 395(10223), 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- Kaiafa, G., Savopoulos, C., Karlafti, E., Pantazi, K., Paramythiotis, D., Thomaidou, E., Daios, S., Ztriva, E., Gionis, M., Fyntanidou, V., Argiriadou, H., & Didangelos, T. (2022). Coagulation Profile of COVID-19 Patients. *Life*, 12(10), 1658. <https://doi.org/10.3390/life12101658>
- 9 Li, C., Zhao, C., Bao, J., Tang, B., Wang, Y., & Gu, B. (2020). Laboratory Diagnosis of Coronavirus Disease-2019 (COVID-19). *Clinica Chimica Acta*, 510, 35–46.
- Li, Y., Deng, Y., Ye, L., Sun, H., Du, S., Huang, H., Zeng, F., Chen, X., & Deng, G. (2021). Clinical Significance of Plasma D-Dimer in COVID-19 Mortality. *Frontiers in Medicine*, 8. <https://doi.org/10.3389/fmed.2021.638097>
- 20 Lippi, G., & Mattiuzzi, C. (2020). Hemoglobin Value May be Decreased in Patients with Severe Coronavirus Disease 2019. *Hematology, Transfusion and Cell Therapy*, 42(2), 116–117. <https://doi.org/10.1016/j.htct.2020.03.001>
- Liu, W., Tao, Z.-W., Wang, L., Yuan, M.-L., Liu, K., Zhou, L., Wei, S., Deng, Y., Liu, J., & Liu, H.-G. (2020). Analysis of Factors Associated with Disease Outcomes in Hospitalized Patients with 2019 Novel Coronavirus Disease. *Chinese Medical Journal*, 133(09), 1032–1038.
- 19 Mus, R., Thasliifa, T., Abbas, M., & Sunaidi, Y. (2021). Studi Literatur: Tinjauan Pemeriksaan Laboratorium pada Pasien COVID-19. *Jurnal Kesehatan Vokasional*, 5(4), 242.
- Naymagon, L., Zubizarreta, N., Feld, J., van Gerwen, M., Alsen, M., Thibaud, S., Kessler, A., Venugopal, S., Makki, I., Qin, Q., Dharmapuri, S., Jun, T., Bhalla, S., Berwick, S., Christian, K., Mascarenhas, J., Dembitzer, F., Moshier, E., & Tremblay, D. (2020). Admission D-Dimer Levels, D-Dimer Trends, and Outcomes in COVID-19. *Thrombosis Research*, 196, 99–105. <https://doi.org/10.1016/j.thromres.2020.08.032>
- Peng, J., Qi, D., Yuan, G., Deng, X., Mei, Y., Feng, L., & Wang, D. (2020). Diagnostic Value of Peripheral Hematologic Markers for Coronavirus Disease 2019 (COVID-19): A Multicenter, Cross-Sectional Study. *Journal of Clinical Laboratory Analysis*, 34(10). <https://doi.org/10.1002/jcla.23475>
- Prozan, L., Shusterman, E., Ablin, J., Mitelpunkt, A., Weiss-Meilik, A., Adler, A., Choshen, G., & Kehat, O. (2021). Prognostic Value of Neutrophil-To-Lymphocyte Ratio in COVID-19 Compared with Influenza and Respiratory Syncytial Virus Infection. *Scientific Reports*, 11(1), 21519. <https://doi.org/10.1038/s41598-021-00927-x>
- 25 Rostami, M., & Mansouritorghabeh, H. (2020). D-Dimer Level in COVID-19 Infection: A Systematic Review. *Expert Review of Hematology*, 13(11), 1265–1275.

Ryadi, Y., Rahardjo, S. S., & Ichsan, B. (2022). ¹¹ Meta-Analysis: The Effect of D-Dimer on Mortality in Patients with COVID-19. *Journal of Epidemiology and Public Health*, 7(4), 552–561. <https://doi.org/10.26911/jepublichealth.2022.07.04.11>

Sabatudung, C., Bia, M. B., & Kambuno, N. T. (2023). Gambaran Hematologi Rutin dan D-Dimer Pasien COVID-19 Kota Kupang. *Journal of Indonesian Medical Laboratory and Science (JoIMedLabs)*, 4(1), 14–24.

¹³ Shama, Mahmood, A., Mehmood, S., & Zhang, W. (2023). Pathological Effects of SARS-CoV-2 Associated with Hematological Abnormalities. *Current Issues in Molecular Biology*, 45(9), 7161–7182. <https://doi.org/10.3390/cimb45090453>

Soni, ¹⁴ Gopalakrishnan, R., Vaishya, R., & Prabu, P. (2020). D-Dimer Level is a Useful Predictor for Mortality in Patients with COVID-19: Analysis of 483 Cases. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 14(6), 2245–2249. <https://doi.org/10.1016/j.dsx.2020.11.007>

² Susilo, A., Rumende, C. M., Pitoyo, C. W., Santoso, W. D., Yulianti, M., Herikurniawan, H., Sinto, R., Singh, G., Nainggolan, L., & Nelwan, E. J. (2020). Coronavirus Disease 2019: Tinjauan Literatur Terkini. *Jurnal Penyakit Dalam Indonesia*, 7(1).

Terpos, E., Ntanasis-Stathopoulos, I., Elalamy, I., Kastritis, E., Sergentanis, T. N., Politou, M., Παλτοπούλου, Θ., Gerotziapas, G. T., & Dimopoulos, M. A. (2020). Hematological Findings and Complications of COVID-19. *American Journal of Hematology*, 95(7):834-847 <https://doi.org/10.1002/ajh.25829>

⁵ Wu, S. Y., Yau, H. S., Yu, M. Y., Tsang, H., ²¹ Chan, L. W. C., Cho, W. C. S., Shing Yu, A. C., Yuen Yim, A. K., Li, M. J. W., & Wong, Y. K. E. (2020). The Diagnostic Methods in the COVID-19 Pandemic, Today and in the Future. *Expert Review of Molecular Diagnostics*, 20(9), 985–993.

²⁶ Yu, B., Li, X., Chen, J., Ouyang, ²⁶ M., Zhang, H., Zhao, X., Tang, L., Luo, Q., Xu, M., Yang, L., Huang, G., Liu, X., & Tang, J. (2020). ²⁶ Evaluation of Variation in D-Dimer Levels Among COVID-19 and Bacterial Pneumonia: a Retrospective Analysis. *Journal of Thrombosis and Thrombolysis*, 50(3), 548–557. <https://doi.org/10.1007/s11239-020-02171-y>

¹⁶ Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu, Y., Li, B., Huang, C.-L., Chen, H.-D., Chen, J., Luo, Y., Guo, H., Jian, ²² R.-D., Liu, M.-Q., Chen, Y., Shen, X.-R., Wang, X., ... Shi, Z.-L. (2020). A Pneumonia Outbreak Associated with a New Coronavirus of Probable Bat Origin. *Nature*, 579(7798), 270–273. <https://doi.org/10.1038/s41586-020-2012-7>

Routine Hematology Description and D-dimer COVID-19 Patients at Sidoarjo Regional Hospital

ORIGINALITY REPORT

21%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

12%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Tobias Tritschler, Marie-Eve Mathieu, Leslie Skeith, Marc Rodger et al. "Anticoagulant interventions in hospitalized patients with COVID-19: A scoping review of randomized controlled trials and call for international collaboration", *Journal of Thrombosis and Haemostasis*, 2020
Publication 1%
- 2** ejournal.unisnu.ac.id
Internet Source 1%
- 3** www.ncbi.xyz
Internet Source 1%
- 4** Submitted to London School of Management Education
Student Paper 1%
- 5** pubs.acs.org
Internet Source 1%
- 6** Abdulkadir, M. B., W. B. R. Johnson, and R. M. Ibraheem. "Validity and accuracy of maternal 1%

tactile assessment for fever in under-five children in North Central Nigeria: a cross-sectional study", BMJ Open, 2014.

Publication

7	Antonio Mastroianni, Valeria Vangeli, Luciana Chidichimo, Filippo Urso et al. "Use of canakinumab and remdesivir in moderate-severe COVID-19 patients: A retrospective analysis", International Journal of Immunopathology and Pharmacology, 2023	1 %
Publication		
8	Submitted to Marian University	1 %
Student Paper		
9	Submitted to Darton State College	1 %
Student Paper		
10	researchoutput.csu.edu.au	1 %
Internet Source		
11	garuda.kemdikbud.go.id	1 %
Internet Source		
12	scholar.google.co.id	1 %
Internet Source		
13	Submitted to Australian National University	1 %
Student Paper		
14	Ikhwan Rinaldi, Mira Yulianti, Evy Yuniastuti, Wulyo Rajabto et al. "Factors Associated with All-Cause 30-Day Mortality in Indonesian	1 %

Inpatient COVID-19 Patients at Cipto
Mangunkusumo National General Hospital",
Journal of Clinical Medicine, 2024

Publication

15

Rosalinda Agnestya Indra, Neneng Martini
Neneng Martini, Ariyati Mandiri, Akhmad Yogi
Pramatirta. "Description Of Hemoglobin
Levels In Pregnant Women With Covid-19
Based On Characteristics", JKM (Jurnal
Kebidanan Malahayati), 2024

Publication

16

Submitted to Colorado State University Fort
Collins

Student Paper

17

Nugroho Eko Wirawan Budianto, Nugrahadi
Dwi Pasca Budiono. "Analisis Pengaruh
Pemberian Ekstrak Etanol Buah Rambusa
(Passiflora foetida L.) Terhadap Zona Hambat
Pertumbuhan Bakteri Escherichia coli",
Journal of Herbal, Clinical and Pharmaceutical
Science (HERCLIPS), 2023

Publication

18

Roy Rillera Marzo, Mohamed ElSherif, Muhd
Siv Azhar Merican Bin Abdullah, Hui Zhu Thew
et al. "Demographic and work-related factors
associated with burnout, resilience, and
quality of life among healthcare workers
during the COVID-19 pandemic: A cross

1 %

1 %

1 %

1 %

sectional study from Malaysia", *Frontiers in Public Health*, 2022

Publication

19	jmiki.apfirmik.or.id Internet Source	1 %
20	Submitted to Universitas Sebelas Maret Student Paper	1 %
21	Submitted to University of Minnesota System Student Paper	1 %
22	Submitted to Zdravstveno veleučilište u Zagrebu / University of Applied Health Sciences Student Paper	1 %
23	aquja.alquds.edu Internet Source	1 %
24	ejurnal.ubharajaya.ac.id Internet Source	1 %
25	medicra.umsida.ac.id Internet Source	1 %
26	Lucas C. Godoy, Ewan C. Goligher, Patrick R. Lawler, Arthur S. Slutsky, Ryan Zarychanski. "Anticipating and managing coagulopathy and thrombotic manifestations of severe COVID-19", <i>Canadian Medical Association Journal</i> , 2020 Publication	1 %

27

ahlimedia.com

Internet Source

1 %

28

jurnal.unbrah.ac.id

Internet Source

1 %

29

new.berkalahayati.org

Internet Source

1 %

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography Off