

# 1. The Effect of Bintaro (Cerbera manghas) Leaf Extract on Transmission of Aphids (Homoptera) in Chili (Capsicum annuum) Plants

*by* Dwi Haryanta

---

**Submission date:** 11-Jan-2021 11:57PM (UTC-0800)

**Submission ID:** 1486213600

**File name:** smission\_of\_Aphids\_Homoptera\_in\_Chili\_Capsicum\_annuum\_Plants.pdf (1.4M)

**Word count:** 7407

**Character count:** 33133

# The Effect of Bintaro (*Cerbera manghas*) Leaf Extract on Transmission of Aphids (Homoptera) in Chili (*Capsicum annuum*) Plants

Achmadi Susilo<sup>1</sup>, Dwi Haryanta<sup>2</sup>, Tatuk Tojibatus Sa'adah<sup>3</sup>

<sup>1,2,3</sup>Faculty of Agriculture, University of Wijaya Kusuma Surabaya  
Email: [dwi.haryanta@yahoo.com](mailto:dwi.haryanta@yahoo.com)

52

Received: November 19. Revised: December 18, 2020. Accepted: December 21, 2020.  
Published: December 24, 2020

**Abstract-** One of the threats in the cultivation of chili is the presence of aphids attack. The study aims to determine the effect of bintaro / (*Cerbera manghas*) leaf extract spraying as a bio-pesticide on the transmission patterns of aphids in chili plants. The experimental method refers to the Painter resistance test (Painter, 1951). Factorial experiment with factor I treatment was the concentration of bintaro leaf extract with six levels namely 0.0% concentration; 2.5% concentration; 5.0% concentration; 7.5% concentration; 10.0% concentration; and 12.5% concentration. Factor II treatment was the frequency of spraying namely F1 sprayed once a week (Sunday), and F2 is sprayed twice a week (Sunday and Wednesday). Overall there are 12 treatment combinations. The experimental unit is a polybag planted by one stem of chili per polybag so that it requires 60 plant polybags, carried out in a controlled field from the presence of other animals by covering it with a mosquito net. In the inside of the lid of mosquito net is infested with chili plants which are attacked by aphids. Source of inoculum of one plant stem attacked by aphids for each containment/lid (each experimental group). The number of insects that transmitted to the treatment plants was observed every two days or before spraying, while the damage to plants at the end of the experiment. The results showed that there were two types of aphids that attacked the inoculum source plant namely *Bemesia tabaci* (whitefly) and *Aphis gossypii* (Aphid) and there were symbionts in the form of black ants (*Dolichoderus thoracicus* Smith).

Transmission occurs since one day after infestation. Population density of the three insect species in each treatment was significantly different, there was an interaction between treatment concentrations with the frequency of bintaro leaf extract spraying, but the density was not consistent from one observation to the next. Plant damage occurs in all plants and the frequency of spraying can reduce damage to the chili plants.

**Keywords—** bintaro leaf extract, aphids, chili

## I. INTRODUCTION

In chilli cultivation is inseparable from plant pest organisms/OPT (organisme pengganggu tanaman), namely pests, diseases and weeds. One of the important pests that attack chilli plants either organically and conventionally cultivated is the aphid of *Aphis gossypii* Glover (Homoptera: Aphididae). Aphids is one of the important pests in vegetable crops, especially in the lowlands. curly virus losses incurred as pests reach 35% and as curly virus vectors can result in losses of up to 90%. In controlling pests, farmers generally use chemical pesticides. The widespread and continuous use of pesticides can reduce losses due to pest attacks. However, the use of these chemical pesticides poses serious new problems namely pest resistance and resurgence, loss of natural enemies of pest insects, death of pollinating insects and other hazards to livestock and humans. To reduce this impact the plant pest control can be done by using plants that have the potential as botanical insecticides [1].

The use of plant extracts as bio insecticides or biopesticides has been widely recommended to farmers. Information on biopesticides is generally still limited to plant raw materials and target pests. The recommended technology is still conventional. There are twelve types of plants that are recommended by agricultural extension workers in 20 countries, namely garlic (*Allium sativum*), neem (*Azadirachta*

*indica*), chilli (*Capsicum* spp.), weed (*Chromolaena odorata*), gliriside (*Gliricidia sepium*), *Melia azedarach*, noni (*Moringa oleifera*), tobacco (*Nicotiana tabacum*), *Ocimum gratissimum*), tephrosia (*Tephrosia vogelii*), tree marigold (*Tithonia diversifolia*), and bitter leaf (*Vernonia amygdalina*) [2]. Extract concentration of 50% is the most effective, environmentally friendly, inexpensive and effective method of controlling future pests (Lawal, Aliyu & Adamu, 2015). Botanical insecticides continue to attract attention for small farms and the environment worldwide, considered as an alternative to synthetic insecticides. The use of secondary metabolites by extracting plants is a tradition more than 3000 years ago. Intensive research on various commercial products still cannot meet global demand for biopesticides (Pavela, 2016). The use of plant extracts and botanical insecticides are biopesticides that do not pollute the environment, and can be included in Integrated Pest Control (IPC) programs and organic farming [3].

Bintaro (*Cerbera manghas*) plant which is widely known by the public today is widely used for greening as well as urban ornamental plants. Plant from the genus *Cerbera* has the potential to be antifungal, insecticide, antioxidant, and antitumor [4], and *Cerbera manghas* can give a significant effect on the mortality of termites (*Coptotermes* sp.) with an extract concentration of 10% [5]. This is because bintaro contains cerberine compounds which are toxic and can kill insects [6]. *Cerbera odollam* extract has a significant effect on mortality and inhibits the development of *Eurema* spp. insect pests by giving a concentration of 1%. *Cerbera odollam* seed extract can affect the bioactivity of *Pteroma plagiophleps* larvae and *Spodoptera litura* F. *Cerbera odollam* has secondary metabolite compounds, such as saponins, polyphenols and alkaloids and terpenoids. Secondary metabolite compounds containing N (such as alkaloids and saponins), and the phenol group compounds (such as flavonoids and tannins) are polar so it can be dissolved in polar or semipolar solvent, such as methanol. Each of secondary metabolite compound has a different working power as an insecticide with various mechanisms [7].

Bintaro plants are almost never attacked by pests, so it is hoped that host plants that are sprayed (treated) with bintaro plant extracts will be protected from insect pests as bintaro plants. Need to look for biopesticide ingredients with abundant availability so that they are easily available and inexpensive. This study aims to examine the ability of bintaro leaf extract in rejecting the presence of aphids in chili plants. The results of the study could become information or a reference for the utilization of Bintaro plant waste which abundant in the city of Surabaya as an organic pesticide in urban farming practices.

## II MATERIAL AND METHOD

**Study on the repellency of bintaro leaf extracts toward the Aphids (Homoptera)-** Repellency test of bintaro leaf extract on the hemiptera order insects using *Aphis* sp. in chili plants. The test method uses the Painter plant resistance test approach (Painter, 1951), namely chilli plants treated with various concentrations of bintaro leaf extract infested by *Aphis* sp

insect. Plants that are not visited by insects or the effect of the smallest insects are then considered to have a repellency power.

Factorial experiment with factor I treatment was the concentration of bintaro leaf extract sprayed on six levels of chilli plants namely: K<sub>0</sub>:concentration of 0,0%; K<sub>1</sub>:concentration of 2,5%; K<sub>2</sub>:concentration of 5,0%; K<sub>3</sub>:concentration of 7,5%; K<sub>4</sub>:concentration of 10,0%; and K<sub>5</sub>:concentration of 12,5%. Factor II treatment was the frequency of spraying namely F1 sprayed once a week (sunday), and F2 is sprayed twice a week (Sunday and Wednesday). There are 12 treatment combinations. The experimental unit is a polybag planted by one stem of chili per polybag so that it requires 60 plant polybags, carried out in a controlled field from the presence of other animals by covering it with a mosquito net.

An illustration of the method of implementation is to design a controlled place (with a mosquito net lid), in the middle are placed insects that will be tested with the condition that active moves to get the host, and around it we place the test units according to the treatment. Experimental variable is the response of insects released in the middle in visiting materials that have been treated with bintaro leaf extract.

In addition to the above method (insects are left to choose), experiments are also conducted to determine the response of insects that are given the feed that has been treated with bintaro leaf extract.

Preparation of Test Insect for Chili Plants Aphids. Healthy chili plants aged 6 weeks after planting placed in the middle of a chili garden which indicated to be attacked by aphids. Plants that have been infected by aphids with a population density of 200-400 tails per plant with a population structure of nymphs and adults taken as a source of inoculum in the experiment. It is very possible that aphids that attacks chili plants of prospective inoculant more than one species, or even followed by other insects associated with aphids.

Preparation of Bintaro Leaf Extract. The making of bintaro leaf extracts using water solvents. Fresh healthy bintaro leaves are washed and then air dried. The clean leaves are cut into small sizes and extracted using a *homogenizer* or blender for 15 minutes. The solution is then filtered with a cloth and the liquid is an extract that will be used for the experiment. In order to be more effective in the extract liquid need to be added with the soap amounted to 2 grams of per liter of liquid. Concentration uses a ratio of the weight of leaves and water as a solvent, for example to get a concentration of 5%, then as much as 50 grams of bintaro leaves are extracted with 1000 cc of water.

Experimental variable is the average number of each insect of aphids or insects associated with aphids. Observations were made on the day of treatment (before spraying) and two days after treatment. Because spraying is carried out every Sunday and Wednesday, then observations are conducted on Tuesday, Wednesday, Friday and Sunday. Besides that it also measured the level of plants damage by scoring method at the end of the observation (one month since the source of the inoculum was included in the experimental plant lid). The spraying of chilli plants with bintaro leaf extract according to the treatment is still conducted until the end of the experiment.



### III RESULTS AND DISCUSSION

**Research Result-** There were two types of aphids that attacked the inoculum source plant namely *Bemesia tabaci* (whitefly) and *Aphis gossypii* (Aphid) and there were symbionts in the form of black ants (*Dolichoderus thoracicus* Smith). The insect population at each source of inoculum is very high (more than 500 individuals / plants) with adult and nymph composition. The three organisms are transmitted to the chili plants which are sprayed with bintaro leaf extracts in a lid. Transmission can fly for the winged, or through the media of black ants, which is also quite high in number. A description of the three types of insects is presented in Figure 4. The development of the population of the three types of insects in each treatment plant is presented in Table 1, Table 2, and Table 3.

#### **Insect population of *Bemesia tabaci* (Whitefly)-**

Transmission of the *Bemesia tabaci* insect from the source of the inoculum to the treatment plant occurs starting one day after infestation. Whitefly found in treatment plants (transmission from inoculum sources) consist of the adult and nymphs phase. The average *Bemesia tabaci* insect population in each treatment plant from initial observation to the end is presented in Table 1.a (Figure 1.a.) and Table 1.b.(Figure 1.b.)).

Whitefly are found in all experimental plants. The results of the variance analysis of whitefly population data in each plant showed that there were significant differences between treatments (there was an interaction between concentration and frequency of treatment). In general, it can be seen that the population density on the second day decreased significantly compared to the first day, and occurred in all treatments. The population distribution looks random and inconsistent namely the highest number or the least number shifts at each observation, indicating its presence is not affected by the spraying treatment of bintaro leaf extract.

***Aphis gossypii* (Aphid) insect population-** Transmission of *Aphis gossypii* insects from the source of the inoculum to the treatment plant occurs from one day after infestation. Aphid found in treatment plants (transmission from inoculum sources) consist of adult and nymphs phase. The average insect population of *Aphis gossypii* in each treatment plant from initial observation to the end is presented in Table 2.a Figure 2.a and Table 2.b.(Figure 2.b.).

Aphids are found in all experimental plants. The results of the variance analysis of aphids population data in each plant showed that there were significant differences between treatments (there was an interaction between concentration and frequency of treatment). Population density is relatively stable fluctuates with increases and decreases in a constant range of values. The population distribution looks random and inconsistent namely the highest number or the least number shifts at each observation, indicating its presence is not affected by the spraying treatment of bintaro leaf extract.

#### **Black Ant (*Dolichoderus thoracicus* Smith) Population -**

Transmission of black ant insects from inoculum sources to treatment plants occurs starting one day after infestation. The observed black ants in treatment plants (transmission from

inoculum sources) consist of an adult phase. In addition to those found in plants, there are many black ants that are in the plant environment at the surface of the soil or in polybags that are more numerous than those in plants. The average population of black ant insects in each treatment plant from initial observation to the end is presented in Table 3.a and Table 3.b.

Black ants are found in all experimental plants and in their environment. Black ant mobility is very high, adding to the impression that the population is very high. The results of the variance analysis of black ant population data in each plant showed that there were significant differences between treatments (there was an interaction between concentration and treatment frequency). The population distribution looks random and inconsistent namely the highest number or the least number shifts at each observation, indicating its presence is not affected by the spraying treatment of bintaro leaf extract.

**The Damage of Chili Plant-** Symptoms of plant damage due to aphids infestation are brownish green undulating leaves, plants cannot grow due to the growing point is dead, then followed by the presence of lateral buds. The attacked leaves there are fallen earlier so that the upper leaves look fall earlier than the leaves below. Plants that have died shoots already confirmed the plant will not produce because there are no shoots where the flowers usually appear. If all shoots are damaged, then the damage score is 5 even though the leaves still look fresh and allow new shoots/buds to emerge. Damage to chili was observed 28 days after infestation of pests or on plants aged 60 days after planting. Data on chili crop damage for each treatment presented in Table 4.

All experimental plants suffered damage ranging from minor/light damage to severe/heavy damage. Severe/heavy damage with a greater score of 3 is in control plants (sprayed with a concentration of 0%) and sprayed plants in concentrations of 5% and 10% once a week. Minor/light damage with a score below 2 is found in plants sprayed with a concentration of 7.5% twice a week. The results of the variance analysis of damage data on chili plants showed that there were significant differences between treatments (there was an interaction between concentration and frequency of treatment). In addition, it can be seen that there is a tendency that those which sprayed twice a week are relatively lighter damage compared to those which sprayed once a week.

**Discussion-**Inoculum source plants attacked by two types of aphids, namely *Bemesia tabaci* (whitefly) and *Aphis gossypii* (aphid) and there are symbionts in the form of black ants (*Dolichoderus thoracicus* Smith). This is consistent with the research data [8] in one palm tree there are three types of insects together and symbiosis occurs with several types of black ants. The pattern of distribution of black ant populations is clustered followed the prey (food sources), and symbiosis with insect on plants [9].

Transmission of whitefly occurs a day after infestation. The whitefly population was significantly different between treatments, but it was inconsistent from one observation to the next, and the population tended to decline. The treatment plant group was in a limited space (4 m<sup>2</sup>) so that the treatment effect of the bintaro leaf extract concentration becomes biased, but in cumulative the bintaro leaf extracts can reduce the population

of whitefly. In line with research [10] which concluded that garlic extract concentration of 5% could effectively reduce the population of whitefly in chili the same with the insecticide of Imidacloprid, Bifenthrin and acetamiprid. The single treatment of ginger extract, onion, and chili can suppress the termite population, the mixture treatment of ginger extract with red onion has a termite repellency equal to 88.89% [11]. Research [12] concluded that the spraying of garlic extract and chili extract with a concentration of 10 grams of material for 1000 cc extract solution could reduce aphid insect population and increase mustard production. Brotowali (*Tinospora crispa*) stem extract treatment with the concentration of 3,125% caused mortality of aphids (*Aphis gossypii*) amounted to (82,50%) at 24 hours after treatment, and at concentration of 25% caused mortality equal to 96.25% [11]. According to the conclusion of the study [13] the application of 75 ml of citronella extract added with 100 ml of water can reduce the population of brown aphid on chili almost 100%. The most effective botanical pesticide of soursop extract reduced aphids and mite populations by 80% and 76%. Neem extract decreased aphid population by 73%, and mite population decreased by 56% [14]. Plant extracts are more effective for whitefly on egg and nymphs (pre-adult) stadia [15].

Transmission of Aphis insect occurs a day after infestation. Aphis insect population significantly different between treatments, but it was inconsistent from one observation to the next, and the population tended to be consistent (not decreased) meaning that the treatment of bintaro leaf extract did not affect the decrease in the aphid insect population. In accordance with the research [16] concluded that the biopesticide treatment was unable to suppress the *Myzus persicae* population in tobacco compared to the use of synthetic pesticides. Biopesticides cannot suppress aphid populations as the conclusion of the study [17] the treatment of spraying of biopesticide extracts of two types of plants is only able to cause death 54,4 and 75% of aphids insect far below 4 types of synthetic pesticides which on average cause death above 90% at 48 hours after treatment. Research [18] concluded that gamal leaf extract contains an insecticide that can cause the death of aphids (*Aphis gossypii*) on chili plants. Spraying of leaf extract with a concentration of 15 grams of leaves in 1000 extracts of the solution can cause the death equal to 49,8% in 96 hours after application. Papaya leaf extract can kill aphid infestation on long beans. With a concentration of 25 grams per liter the extract solution can cause the death/mortality up to 75 percent. While Polygonum flower extract with a concentration of 5% and tobacco leaf extract with the concentration of 10% can suppress aphid populations (*Myzus* and *Aphis gossypii*) by 70% and 65%, while the azadirachtin botanical pesticide suppresses the population by 60% [19]. According to research [20] the provision of tuba root extract at a dose of 45 ml/L of water gave the highest result of aphids (*Aphis gossypii*) mortality amounted to 25,28%, and the intensity of crop damage equal to 25,74%, compared with mortality control of 13,61% and intensity of crop damage of 49,45%.

Transmission of black ants occurs a day after infestation. The black ant population is significantly different between treatments, but it is not consistent from one observation to the next. The population of black ants is not pure from the source

of the inoculum, at the place of the experiment there were black ants before infestation. According to research [9] in the yard around the house there are several types of ants, including black ants in symbiosis with plant insect. The distribution of ants in the yard following the availability of food, for black ants following the presence of plant insect, after conducted an infestation the plants which attacked by aphid automatically cause the black ant came directly.

The presence of two types of aphids namely *Bemisia tabaci* (whitefly) and *Aphis gossypii* (aphid) caused damage to the experimental plant. The treatment of bintaro leaf extract has no significant effect on population fluctuations of two types of aphids, namely *Bemisia tabaci* (whitefly) and *Aphis gossypii* (aphid), however, it is generally seen that plants that are sprayed twice a week have less damage than those that are sprayed once a week. This can be interpreted that the treatment of bintaro leaf extract affects the eating behavior of two aphids on chili plants. The treatment of *Polygonum persicaria* plant extract as a biopesticide significantly decreased the appetite of the *Nomophila indistinctalis* insect. The spraying of garlic extract and chili extract with a concentration of 10 grams of material for 1000 cc extract solution can reduce damage and increase mustard production [8].

#### IV CONCLUSION

The treatment of Bintaro Leaf Extract (*Cerbera manghas*) did not significantly affect on the transmission and development of the population of *Bemisia tabaci* (whitefly) and *Aphis gossypii* (aphid) aphids on the chili plants. The presence of aphids is followed by a symbiont namely black ants. The treatment of bintaro leaf extract influence on the appetite of aphids seen from the intensity of plant damage. Plants with twice a week of treatment are less damaged than plants with once a week of treatment. Needs to be conducted the further research in the field with the extract concentration increased to 50% (in this study the highest was only 12,5%).

#### V ACKNOWLEDGEMENTS

Manuscript material is part of the scheme of penelitian dasar unggulan perguruan tinggi (PDUPT) year II from the 3 year plan. On this occasion, we would like to thank the Directorate of Research and Community Service Directorate General of Research and Development Strengthening The Ministry of Research Technology and Higher Education who has provided funding for research implementation.

#### References

- [1] R. Mustiarif, "EFEKTIVITAS EKSTRAK BIJI BINTARO (*Cerbera odollam* G.) TERHADAP MORTALITAS KUTU DAUN (*Aphis gossypii* GLOVER) PADA TANAMAN CABAI (*Capsicum annum* L.) SKRIPSI Oleh: Rial Mustiarif NPM. E1J012079 PROGRAM STUDI AGROEKOTEKNOLOGI JURUSAN BUDIDAYA PERTANIAN FAKULTAS PERTANIAN UNIVERSITAS BENGKULU 2016."



- [2] J. Dougoud, S. Toepfer, M. Bateman, and W. H. Jenner, "Efficacy of homemade botanical insecticides based on traditional knowledge. A review," *Agronomy for Sustainable Development*, vol. 39, no. 4, p. 37, 2019.
- [3] S. K. Ghosh, "Integrated field management of aphids (*Myzus persicae* Sulz. and *Aphis gossypii* Glov. Together) on potato (*Solanum tuberosum* L.) using bio-pesticides," *International Journal of Science, Environment and Technology*, vol. 4, no. 3, pp. 682–689, 2015.
- [4] Y. XU, T. ENG, and W. TONG, "Chemical Constituents and Bioactivity of Mangrove Plants in the Genus *Cerbera* [J]," *Journal of Guangxi Academy of Sciences*, vol. 59, 2011.
- [5] D. Tarmadi, H. Prianto, I. Guswenrivo, T. Kartika, and S. Yusuf, "Pengaruh Ekstrak Bintaro (*Carbera odollam* Gaertn) dan Kecubung (*Brugmansia candida* Pers) terhadap Rayap Tanah *Coptotermes* sp Influence of Bintaro (*Carbera odollam* Gaertn) and Kecubung (*Brugmansia candida* Pers) act against Subterranean Termite *Coptotermes* sp," *Jurnal Ilmu dan Teknologi Kayu Tropis*, vol. 5, no. 22, pp. 38–42, 2007.
- [6] N. Nur Apriliyanti, "Uji Potensi Racun Ekstrak Daun Bintaro (*Cerbera odollam* Gaertn) Sebagai Bioinsektisida Terhadap Mortalitas Larva *Plutella Xylostella* Pada Tanaman *Brassica oleracea*," 2016.
- [7] S. Utami, "Aktivitas insektisida bintaro (*Cerbera odollam* Gaertn) terhadap hama *Eurema* spp. pada skala laboratorium," *Jurnal Penelitian Hutan Tanaman*, vol. 7, no. 4, pp. 211–220, 2010.
- [8] Z. AMILASARIY, H. Herlina, and N. A. Fithri, "Uji toksisitas akut ekstrak biji palem putri (*Adonidia merrillii* (Becc.) Becc) terhadap tikus putih betina galur wistar dengan fixed procedure," 2015.
- [9] R. Riyanto, "Kepadatan, Pola Distribusi dan Peranan Semut pada Tanaman di Sekitar Lingkungan Tempat Tinggal," *Jurnal Penelitian Sains (JPS)*, vol. 10, no. 2, pp. 241–253, 2007.
- [10] M. Iftikhar, M. A. Khan, and S. Haider, "Comparative Efficacy of Insecticides and Plant Extracts for Management of ChiLCV Disease in Relation to Epidemiology," 2017.
- [11] M. S. D. Permadi and H. Fitrihidajati, "Pengaruh Pemberian Ekstrak Batang Brotowali (*Tinospora crispa*) terhadap Mortalitas Kutu Daun (*Aphis gossypii*)," *AnteraBio*, vol. 8, no. 2, 2019.
- [12] R. Pavela, "History, presence and perspective of using plant extracts as commercial botanical insecticides and farm products for protection against insects—a review," *Plant Protection Science*, vol. 52, no. 4, pp. 229–241, 2016.
- [13] F. Arfianto, "Pengendalian hama kutu daun coklat pada tanaman cabe menggunakan pestisida organik ekstrak serai wangi," *Anterior Jurnal*, vol. 16, no. 1, pp. 57–66, 2014.
- [14] M. S. Saleem, T. S. Batool, M. F. Akbar, S. Raza, and S. Shahzad, "Efficiency of botanical pesticides against some pests infesting hydroponic cucumber, cultivated under greenhouse conditions," *Egyptian Journal of Biological Pest Control*, vol. 29, no. 1, p. 37, 2019.
- [15] A. Cruz-Estrada, M. Gamboa-Angulo, R. Borges-Argáez, and E. Ruiz-Sánchez, "Insecticidal effects of plant extracts on immature whitefly *Bemisia tabaci* Genn.(Hemiptera: Aleyroideae)," *Electronic Journal of Biotechnology*, vol. 16, no. 1, pp. 6–6, 2013.
- [16] A. Khan *et al.*, "Efficacy of a biopesticide and synthetic pesticides against tobacco aphid, *Myzus persicae* Sulz.(Homoptera, Aphididae), on tobacco in Peshawar," *Journal of Entomology and Zoology Studies*, vol. 4, pp. 337–373, 2015.
- [17] K. Elango, S. Sridharan, P. Saravanan, and S. Balakrishnan, "Laboratory evaluation of insecticides and biopesticides against pomegranate aphid *Aphis punicae* Passerini," *IJCS*, vol. 5, no. 5, pp. 1810–1812, 2017.
- [18] N. Agustina, M. I. Pramudi, and N. Aidawati, "Pengaruh Larutan Daun Gamal (*Gliricidia sepium*) Terhadap Mortalitas Kutu Daun *Aphis gossypii* pada Cabai (*Capsicum annum* L)," *Jurnal Proteksi Tanaman TROPIKA*, vol. 2, no. 1, pp. 86–91, 2019.
- [19] L. Quesada-Romero, C. Fernández-Galleguillos, J. I. J. mann, M. A. Bravo, and E. Fuentes-Contreras, "Antifeedant and insecticidal activity of *Polygonum persicaria* extracts on *Nomophila indistinctalis*," *Journal of Pharmacy & Pharmacognosy Research*, vol. 5, no. 3, pp. 167–173, 2017.
- [20] R. R. Kinansi, S. W. Handayani, D. Prastowo, and A. O. Y. Sudarno, "Efektivitas Ekstrak Etanol Akar Tuba (*Derris elliptica*) terhadap Kematian Periplaneta americana dengan Metode Spraying," *Balaba: Jurnal Litbang Pengendalian Penyakit Bersumber Binatang Banjarnegara*, pp. 147–158, 2018.

#### Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

[https://creativecommons.org/licenses/by/4.0/deed.en\\_US](https://creativecommons.org/licenses/by/4.0/deed.en_US)

Table 1a. The average population of *Bemisia tabaci* insects in chili plants with various concentrations treatment of bintaro leaf extract in 1-10 days after pest infestation

Treatment	observation on the length time (days) after inoculum source infestation							
	1	3	5	6	8	10		
K <sub>0</sub> F <sub>1</sub>	35.6 bc	5.4 ab	3.6 cd	4.6 a	5.4 ab	7.2 bc		
K <sub>1</sub> F <sub>1</sub>	109.6 a	17.8 a	34.4 a	27.6 a	30.4 a	42.8 a		
K <sub>2</sub> F <sub>1</sub>	43.7 bc	7.0 ab	0.0 d	13.2 a	13.4 ab	21.0 ab		
K <sub>3</sub> F <sub>1</sub>	28.4 c	2.6 b	1.6 d	3.0 a	1.6 b	1.8 c		
K <sub>4</sub> F <sub>1</sub>	33.6 bc	4.4 ab	0.0 d	16.4 a	22.0 ab	25.2 abc		
K <sub>5</sub> F <sub>1</sub>	44.0 bc	1.6 b	0.6 d	5.0 a	5.4 ab	9.4 bc		
K <sub>0</sub> F <sub>2</sub>	38.6 bc	8 ab	15.0 abc	14.0 a	17.6 ab	24.0 ab		
K <sub>1</sub> F <sub>2</sub>	62.0 bc	3.4 ab	5.4 bcd	8.0 a	11.6 ab	11.8 bc		
K <sub>2</sub> F <sub>2</sub>	28.8 bc	4.4 ab	4.2 cd	6.2 a	7.2 ab	19.6 abc		
K <sub>3</sub> F <sub>2</sub>	30.2 c	11.6 ab	19.4 ab	21.8 a	28.4 ab	27.2 ab		
K <sub>4</sub> F <sub>2</sub>	35.0 bc	6.6 ab	5.6 bcd	9.6 a	16.4 ab	14.2 abc		
K <sub>5</sub> F <sub>2</sub>	76.8 ab	3.6 ab	6.4 bcd	9.2 a	11.4 ab	20.2 abc		




Figure 1.a. The average population of *Bemisia tabaci* insect in chili plants with various concentration of bintaro leaf extract in 1-10 days after infestation

Table 1b. The average population of *Bemisia tabaci* insects in chili plants with various concentrations treatment of bintaro leaf extract in 12-20 days after pest infestation

Treatment	observation on the length time (days) after inoculum source infestation					
	12	13	15	17	19	20
K <sub>0</sub> F <sub>1</sub>	8.0 ab	7.6 ab	8.0 ab	7.8	5.8 ab	2.6 b
K <sub>1</sub> F <sub>1</sub>	16.8 a	16.8 ab	17.4 a	20.0	12.2 ab	7.0 ab
K <sub>2</sub> F <sub>1</sub>	16.7 ab	16.6 ab	15.0 a	17.6	11.0 ab	14.0 a
K <sub>3</sub> F <sub>1</sub>	2.6 b	2.0 b	3.2 b	7.4	4.2 ab	6.0 ab
K <sub>4</sub> F <sub>1</sub>	15.0 ab	16.8 ab	8.4 ab	12.0	13.0 a	11.4 ab
K <sub>5</sub> F <sub>1</sub>	7.6 ab	7.56 b	9.8 ab	13.6	12.0 ab	12.8 a
K <sub>0</sub> F <sub>2</sub>	11.6 ab	15.36 b	13.2 ab	13.4	7.6 ab	8.2 ab
K <sub>1</sub> F <sub>2</sub>	13.4 ab	16.55 b	13.0 ab	17.2	2.4 b	5.0 ab
K <sub>2</sub> F <sub>2</sub>	12.8 ab	12.0 63	3.8 ab	5.4	5.0 ab	9.6 ab
K <sub>3</sub> F <sub>2</sub>	15.2 ab	13.4 ab	8.4 ab	8.4	6.4 ab	4.8 ab
K <sub>4</sub> F <sub>2</sub>	14.18 b	17.4 a	11.8 ab	13.2	7.0 ab	10.8 ab
K <sub>5</sub> F <sub>2</sub>	4.8 ab	5.20 ab	6.4 ab	6.0	8.4 ab	8.2 ab

Information: - the average value in a column followed by the same letter is not significantly different based on the BNT/LSD test 5%.  
- before analyzed the data conducted transformation  $\sqrt{x+0,5}$




Figure 1.b.. The average population of *Bemisia tabaci* insect in chili plants with various concentration of bintaro leaf extract in 12-20 days after infestation



Table 2.a The average population of *Aphis gossypii* insects in chili plants with various concentrations treatment of bintaro leaf extract in 1-10 days after pest infestation

Treatment	observation on the length time (days) after inoculum source infestation					
	1	3	5	6	8	10
K <sub>0</sub> F <sub>1</sub>	35.6 bc	38.8 ab	46.2 ab	53.2 a	28.0 ab	29.0 abc
K <sub>1</sub> F <sub>1</sub>	109.6 a	53.6 ab	64.27 ab	68.4 54	30.2 b	47.8 ab
K <sub>2</sub> F <sub>1</sub>	43.7 c	59.0 ab	55.2 ab	42.2 ab	33.0 ab	39.4 abc
K <sub>3</sub> F <sub>1</sub>	28.4 c	24.49 b	26.0 ab	57.8 a	22.2 ab	50.8 abc
K <sub>4</sub> F <sub>1</sub>	33.6 bc	32.2 ab	46.4 ab	49.2 ab	21.4 b	29.0 abc
K <sub>5</sub> F <sub>1</sub>	44.0 bc	38.0 ab	33.8 ab	35.2 ab	24.42 b	24.2 bc
K <sub>0</sub> F <sub>2</sub>	38.6 bc	50.4 ab	41.0 ab	61.0 a	29.0 ab	37.4 abc
K <sub>1</sub> F <sub>2</sub>	62.0 bc	46.8 ab	48.6 ab	38.8 ab	23.4 ab	35.0 abc
K <sub>2</sub> F <sub>2</sub>	28.8 bc	36.31 b	45.8 ab	66.2 a	37.8 ab	42.4 abc
K <sub>3</sub> F <sub>2</sub>	30.2 c	18.2 ab	24.8 ab	30.8 ab	21.8 ab	31.8 abc
K <sub>4</sub> F <sub>2</sub>	35.27 bc	22.8 b	15.8 b	9.2 b	11.4 b	16.6 c
K <sub>5</sub> F <sub>2</sub>	76.8 ab	59.8 a	72.2 a	70.2 a	56.4 a	57.6 a




Figure 2.a.The average population of *Aphis gossypii* insects in chili plants with various concentrations treatment of bintaro leaf extract in 1-10 days after pest infestation

Table 2.b. The average population of *Aphis gossypii* insects in chili plants with various concentrations treatment of bintaro leaf extract in 12-20 days after pest infestation

Treatment	observation on the length time (days) after inoculum source infestation					
	12	13	15	17	19	20
K <sub>0</sub> F <sub>1</sub>	23.0 ab	18.6 abc	18.2 c	23.4 cd	14.2 b	24.2 ab
K <sub>1</sub> F <sub>1</sub>	36.8 a	37.8 a	47.8 ab	53.6 ab	23.4 ab	29.2 ab
K <sub>2</sub> F <sub>1</sub>	48.6 ab	24.0 abc	31.0 abc	28.0abcd	19.44 b	20.6 ab
K <sub>3</sub> F <sub>1</sub>	22.6 ab	31.8 ab	42.2 ab	36.8abcd	23.4 ab	20.0 ab
K <sub>4</sub> F <sub>1</sub>	20.4 ab	19.8 abc	21.0 c	17.4 d	15.8 b	23.8 ab
K <sub>5</sub> F <sub>1</sub>	18.8 b	16.0 bc	31.2 abc	31.0abcd	21.4 ab	28.4 ab
K <sub>0</sub> F <sub>2</sub>	26.0 ab	22.6 abc	48.2 a	44.2 abc	25.6 ab	30.8 ab
K <sub>1</sub> F <sub>2</sub>	30.0 ab	30.6 ab	49.6 a	59.8 a	33.8 a	62.4 a
K <sub>2</sub> F <sub>2</sub>	34.8 ab	33.4 ab	46.4 ab	53.0 ab	34.2 a	32.8 ab
K <sub>3</sub> F <sub>2</sub>	19.6 b	17.6 abc	42.8 ab	40.8abcd	21.0 ab	25.2 ab
K <sub>4</sub> F <sub>2</sub>	13.0 ab	10.4 c	24.4 bc	24.8 bcd	14.41 b	18.8 b
K <sub>5</sub> F <sub>2</sub>	29.8 ab	31.0 ab	46.2 a	45.8 ab	26.4 ab	24.8 ab

Information: - the average value in a column followed by the same letter is not significantly different based on the BNT/LSD test 5%.  
- before analyzed the data conducted transformation  $\sqrt{x+0,5}$




Figure 2.b. The average population of *Aphis gossypii* insects in chili plants with various concentrations treatment of bintaro leaf extract in 12-20 days after pest infestation

Table 3.a.The average population of black ant in chili plants with various concentrations treatment of bintaro leaf extract in 1-10 days after pest infestation

Treatment	observation on the length time (days) after inoculum source infestation					
	1	3	5	6	8	10
K <sub>0</sub> F <sub>1</sub>	8.2 ab	4.4 a	8.6 a	8.2 ab	8.43 b	4.8 a
K <sub>1</sub> F <sub>1</sub>	18.2 a	8.6 a	11.4 a	16.2 ab	12.6 ab	6.4 a
K <sub>2</sub> F <sub>1</sub>	8.0 ab	11.4 a	9.4 a	8.2 ab	11.2 ab	6.0 a
K <sub>3</sub> F <sub>1</sub>	1.6 b	0.8 a	2.4 a	7.6 ab	7.6 ab	13.6 a
K <sub>4</sub> F <sub>1</sub>	10.4 ab	12.0 a	11.2 a	15.2 ab	8.0 ab	4.6 a
K <sub>5</sub> F <sub>1</sub>	12.0 ab	4.4 a	7.6 a	6.8 ab	4.0 ab	7.6 a
K <sub>0</sub> F <sub>2</sub>	9.6 ab	5.2 a	16.0 a	1.18 ab	6.6 a 18	3.6 a
K <sub>1</sub> F <sub>2</sub>	6.2 b	0.0 a	1.4 a	5.8 ab	0.8 b	5.0 a
K <sub>2</sub> F <sub>2</sub>	11.2 ab	8.2 a	13.4 a	16.0 ab	7.4 ab	7.6 a
K <sub>3</sub> F <sub>2</sub>	5.2 ab	4.4 a	3.4 a	3.6 ab	5.8 ab	5.2 a
K <sub>4</sub> F <sub>2</sub>	1.6 b	1.6 a	1.0 a	1.2 b	1.8 b	0.0 a
K <sub>5</sub> F <sub>2</sub>	11.0 ab	9.6 a	9.6 a	16.8 a	15.6 a	8.6 a

Table 3.b.The average population of black ant in chili plants with various concentrations treatment of bintaro leaf extract in 12-20 days after pest infestation

Treatment	observation on the length time (days) after inoculum source infestation					
	12	13	15	17	19	20
K <sub>0</sub> F <sub>1</sub>	3.0 ab	3.2 ab	2.2 bc	1.6 ab	0.0 c	1.8 ab
K <sub>1</sub> F <sub>1</sub>	7.47 a	8.6 ab	11.0 a	7.4 ab	6.8 abc	3.2 ab
K <sub>2</sub> F <sub>1</sub>	4.2 ab	4.8 ab	5.0 abc	5.61 b	8.8 a	2.6 ab
K <sub>3</sub> F <sub>1</sub>	5.2 ab	11.26 a	2.6 bc	3.6 ab	3.2 abc	2.0 ab
K <sub>4</sub> F <sub>1</sub>	5.0 ab	4.8 ab	36 c	0.8 b	0.4 bc	0.0 b
K <sub>5</sub> F <sub>1</sub>	4.4 53	4.0 ab	3.0 abc	2.4 ab	3.4 abc	3.4 ab
K <sub>0</sub> F <sub>2</sub>	1.6 ab	2.0 ab	6.2 29	3.8 ab	2.8 abc	1.6 ab
K <sub>1</sub> F <sub>2</sub>	2.4 ab	1.2 b	6.0 ab	4.8 ab	3.4 abc	3.4 ab
K <sub>2</sub> F <sub>2</sub>	5.4 ab	6.8 ab	4.6 64	3.8 ab	5.6 abc	3.2 ab
K <sub>3</sub> F <sub>2</sub>	0.4 b	1.6 26	2.4 abc	4.2 ab	3.8 abc	4.0 a
K <sub>4</sub> F <sub>2</sub>	1.6 ab	2.4 ab	0.6 c	1.6 ab	1.4 abc	1.8 ab
K <sub>5</sub> F <sub>2</sub>	6.2 a	8.6 ab	6.6 abc	8.6 a	4.2 ab	2.8 ab

Information: - the average value in a column followed by the same letter is not significantly different based on the BNT/LSD test 5%.  
- before analyzed the data conducted transformation  $\sqrt{x+0,5}$



Table 4. The average level of damage to chili due to aphids (*Aphids gossypii* and *Bemesia tabaci*) with the treatment of various concentrations of bintaro leaf extract

Treatment	Average damage score	information
K <sub>0</sub> F <sub>1</sub> (concentration of 0% frequency once a week)	3.2 a	heavily damaged
K <sub>1</sub> F <sub>1</sub> (concentration of 2.5% frequency once a week)	2.5 ab	moderately damaged
K <sub>2</sub> F <sub>1</sub> (concentration of 5.0% frequency once a week)	3.0 ab	heavily damaged
K <sub>3</sub> F <sub>1</sub> (concentration of 7.5% frequency once a week)	2.6 ab	moderately damaged
K <sub>4</sub> F <sub>1</sub> (concentration of 10% frequency once a week)	3.4 a	heavily damaged
K <sub>5</sub> F <sub>1</sub> (concentration of 12.5% frequency once a week)	2.8 ab	moderately damaged
K <sub>0</sub> F <sub>2</sub> (concentration of 0% frequency twice a week)	2.4 ab	moderately damaged
K <sub>1</sub> F <sub>2</sub> (concentration of 2.5% frequency twice a week)	2.8 ab	moderately damaged
K <sub>2</sub> F <sub>2</sub> (concentration of 5.0% frequency twice a week)	2.2 ab	moderately damaged
K <sub>3</sub> F <sub>2</sub> (concentration of 7.5% frequency twice a week)	1.6 b	lightly damaged
K <sub>4</sub> F <sub>2</sub> (concentration of 10% frequency twice a week)	2.4 ab	moderately damaged
K <sub>5</sub> F <sub>2</sub> (concentration of 12.5% frequency twice a week)	2.8 ab	moderately damaged

Information: - the average value in a column followed by the same letter is not significantly different based on the BNT/LSD test 5%.

- before analyzed the data conducted transformation  $\sqrt{x+0.5}$

Score 0: no damaged plants: healthy

Score 1: damaged plants 1 s/d 20% : Very light damage

Score 2: damaged plants 21 s/d 40% : light damage

Score 3: damaged plants 41 s/d 60% : Medium damage

Score 4: damaged plants 61 s/d 80% : Heavy damage


Score 5: damaged plants 81 s/d 100% : Very heavy damage



Figure 1. Experimental material for chilli plants in a mosquito net lid



Figure 2. Mosquito net contain one test run



Information:

- : inoculum source (release of test insects)
- : host / feed that has been given treatment

Figure 3. Design of the placement of inoculum sources toward the treated plants



Figure 4. chili leaves with the pest of *Bemisia tabaci* (whitefly), *Aphis gossypii* (aphid) and *Dolichoderus thoracicus* (black ant)

# 1. The Effect of Bintaro (*Cerbera manghas*) Leaf Extract on Transmission of Aphids (*Homoptera*) in Chili (*Capsicum annuum*) Plants

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

1

Submitted to School of Business and Management ITB

Student Paper

3%

2

[docobook.com](http://docobook.com)

Internet Source

1%

3

[www.iosrjournals.org](http://www.iosrjournals.org)

Internet Source

1%

4

[www.naun.org](http://www.naun.org)

Internet Source

1%

5

[repository.unib.ac.id](http://repository.unib.ac.id)

Internet Source

<1%

6

[www.hindawi.com](http://www.hindawi.com)

Internet Source

<1%

7

[link.springer.com](http://link.springer.com)

Internet Source

<1%

8

[jurnalmapeki.biomaterial-lipi.org](http://jurnalmapeki.biomaterial-lipi.org)

Internet Source

<1%



9

Charilaos Giotis. "Effect of soil amendments and biological control agents (BCAs) on soil-borne root diseases caused by *Pyrenochaeta lycopersici* and *Verticillium albo-atrum* in organic greenhouse tomato production systems", European Journal of Plant Pathology, 04/2009

Publication

<1 %

10

[www.redalyc.org](http://www.redalyc.org)

Internet Source

<1 %

11

[www.ocl-journal.org](http://www.ocl-journal.org)

Internet Source

<1 %

12

[ejournal2.litbang.kemkes.go.id](http://ejournal2.litbang.kemkes.go.id)

Internet Source

<1 %

13

[dergipark.org.tr](http://dergipark.org.tr)

Internet Source

<1 %

14

Norhidayah Che Soh, Noor Shahira Md, Suhaizan Lob, Nurul Faziha Ibr, Husni Hayati Moh, Johar Mohamed. "Effect of Neem Extract on Growth Performance and Post-harvest Quality of Chili", Asian Journal of Plant Sciences, 2020

Publication

<1 %

15

[www.scirp.org](http://www.scirp.org)

Internet Source

<1 %

16

Nusrat Iqbal, Ranju Sharma, Dipak Kumar

<1 %

Hazra, Saurabh Dubey, Natish Kumar, Amrish Aggrawal, Jitendra Kumar. "Successful utilization of waste cooking oil in Neem oil based fungicide formulation as an economic and eco-friendly green solvent for sustainable waste management", Journal of Cleaner Production, 2020

Publication

17

[ejurnal.litbang.pertanian.go.id](http://ejurnal.litbang.pertanian.go.id)

Internet Source

<1 %

18

Olusola Omuetti, I. D. Morton. "Development by extrusion of soyabari snack sticks: a nutritionally improved soya—maize product based on the Nigerian snack (kokoro)", International Journal of Food Sciences and Nutrition, 2009

Publication

<1 %

19

[journal.umpalangkaraya.ac.id](http://journal.umpalangkaraya.ac.id)

Internet Source

<1 %

20

[jtam.ulm.ac.id](http://jtam.ulm.ac.id)

Internet Source

<1 %

21

[repository.unsri.ac.id](http://repository.unsri.ac.id)

Internet Source

<1 %

22

[repository.institutpendidikan.ac.id](http://repository.institutpendidikan.ac.id)

Internet Source

<1 %

23

[mafiadoc.com](http://mafiadoc.com)

Internet Source

<1 %

24

[ejournal.unesa.ac.id](http://ejournal.unesa.ac.id)

Internet Source

<1 %

25

Submitted to Universidad Industrial de Santander UIS

Student Paper

<1 %

26

Anne W. de Valena, Steven J. Vanek, Katherin Meza, Raul Ccanto et al. "Land use as a driver of soil fertility and biodiversity across an agricultural landscape in the Central Peruvian Andes", Ecological Applications, 2017

Publication

<1 %

27

Jenny Carlsson. "Mitochondrial genotypes with variable parts of Arabidopsis thaliana DNA affect development in Brassica napus lines", Theoretical and Applied Genetics, 08/24/2007

Publication

<1 %

28

[talenta.usu.ac.id](http://talenta.usu.ac.id)

Internet Source

<1 %

29

L. S. S. BARROS, L. A. AMARAL, C. S. LORENZON, J. L. JUNIOR, J. G. MACHADO NETO. "Potential microbiological contamination of effluents in poultry and swine abattoirs", Epidemiology and Infection, 2006

Publication

<1 %

30

[www.agriculturejournals.cz](http://www.agriculturejournals.cz)

Internet Source

<1 %



- 31 Ralph L. Obendorf, Elizabeth M. Sensenig, Erin M. Byrt, Anna B. Owczarczyk, Minori Ohashi, Steven R. Schnebly. " Cyclitol galactosides in low-raffinose, low-stachyose soybean embryos after feeding - -inositol, -inositol or -pinitol ", Seed Science Research, 2013  
Publication <1 %
- 
- 32 [www.journalofscience.org](http://www.journalofscience.org)  
Internet Source <1 %
- 
- 33 Oktavianus Lumban Tobing, Yanyan Mulyaningsih. "THE RECONDITIONING GROWTH AND PRODUCTION OF CHILI THROUGH THE BANANA HUMP AND MIMBA LEAF EXTRACT", Indonesian Journal of Applied Research (IJAR), 2020  
Publication <1 %
- 
- 34 Submitted to International School of Azerbaijan  
Student Paper <1 %
- 
- 35 N A Tiffani, H Ramdan, Dungani. " The Characteristics of Stands at Site 23 B, RPH Maribaya, BKPH Parung Panjang, KPH Bogor which is attacked by pests and diseases ", IOP Conference Series: Earth and Environmental Science, 2020  
Publication <1 %
- 
- 36 Elisabeth Olsen, Gjermund Vogt, Annette <1 %

Veberg, Dag Ekeberg, Astrid Nilsson. "Analysis of Early Lipid Oxidation in Smoked, Comminuted Pork or Poultry Sausages with Spices", Journal of Agricultural and Food Chemistry, 2005

Publication

---

37

[digilib.its.ac.id](http://digilib.its.ac.id)

Internet Source

<1 %

---

38

S. F. CARNEGIE. "Contamination of seed potato tubers grown from stem cutting or microplant-derived tubers by *Phoma foveata* and *Polyscytalum pustulans* in Scotland: the influence of site and fungicide treatment at harvest", Annals of Applied Biology, 2/1991

Publication

<1 %

---

39

[worldwidescience.org](http://worldwidescience.org)

Internet Source

<1 %

---

40

[www.dhsspsni.gov.uk](http://www.dhsspsni.gov.uk)

Internet Source

<1 %

---

41

Bernadette Earley. "Effect of road transport for up to 24 hours followed by twenty-four hour recovery on live weight and physiological responses of bulls", BMC Veterinary Research, 2010

Publication

<1 %

---

42

J. E. Quisenberry, W. R. Jordan, B. A. Roark, D.

<1 %

## W. Fryrear. " Exotic Cottons as Genetic Sources for Drought Resistance ", Crop Science, 1981

Publication

43

P. Quénéhervé. "Nematode resistance in bananas: screening results on some wild and cultivated accessions of Musa spp.", Euphytica, 01/2009

Publication

<1 %

44

Yuichi Yoshii. "The effect of wrist position on the relative motion of tendon, nerve, and subsynovial connective tissue within the carpal tunnel in a human cadaver model", Journal of Orthopaedic Research, 08/2008

Publication

<1 %

45

[www.futuremedicine.com](http://www.futuremedicine.com)

Internet Source

<1 %

46

[garuda.ristekbrin.go.id](http://garuda.ristekbrin.go.id)

Internet Source

<1 %

47

VNUA

Publication

<1 %

48

Eduardo Schneid, Poliana Dias de Moraes. "Grain angle and temperature effect on embedding strength", Construction and Building Materials, 2017

Publication

<1 %

49

Colkesen, M, A Kamalak, O Canbolat, Y



Gurbuz, and CO Ozkan. "Effect of cultivar and formaldehyde treatment of barley grain on rumen fermentation characteristics using *in vitro* gas production", South African Journal of Animal Science, 2007.

Publication

<1 %

50

Matthew G. Blanchard. "Benzyladenine Promotes Flowering in Doritaenopsis and Phalaenopsis Orchids", Journal of Plant Growth Regulation, 06/2008

Publication

<1 %

51

[balingtang.litbang.pertanian.go.id](http://balingtang.litbang.pertanian.go.id)

Internet Source

<1 %

52

Keita Shinogaya. "Effective Ways of Enhancing the Quality of Question Generating and Spontaneous Information Search Outside the Classroom", International Journal of Higher Education, 2020

Publication

<1 %

53

Jin Choi, Zahra Aminikhoie, Kyoung-Duck Kim, Sang-Min Lee. "Growth and Fatty Acid Composition of Juvenile Olive Flounder *Paralichthys olivaceus* Fed Diets Containing Different Levels and Ratios of Eicosapentaenoic Acid and Docosahexaenoic Acid", Fisheries and aquatic sciences, 2014

Publication

<1 %

54

Beyene, Solomon T.. "Rangeland degradation in Swaziland: Dip tank use effects on range condition and grazing capacity in three soil types", African Journal of Ecology, 2014.

Publication

&lt;1 %

55

Imre J. Holb. "Effect of six sanitation treatments on leaf litter density, ascospore production of *Venturia inaequalis* and scab incidence in integrated and organic apple orchards", European Journal of Plant Pathology, 2006

Publication

&lt;1 %

56

Tang, X.Y.. "Air-drying and liming effects on exchangeable cadmium mobilization in contaminated soils: A repeated batch extraction study", Geoderma, 20110215

Publication

&lt;1 %

57

M. Benady, , J. E. Simon, D. J. Charles, and G. E. Miles. "Fruit Ripeness Determination by Electronic Sensing of Aromatic Volatiles", Transactions of the ASAE, 1995.

Publication

&lt;1 %

58

A. C. Gama-Rodrigues, N. F. Barros, E. S. Mendonça. "Alterações edáficas sob plantios puros e misto de espécies florestais nativas do sudeste da Bahia, Brasil", Revista Brasileira de Ciência do Solo, 1999

Publication

&lt;1 %

59

docplayer.net

Internet Source

&lt;1 %

60

pps.uny.ac.id

Internet Source

&lt;1 %

61

Eder Jorge de Oliveira, Taliane Leila Soares, Cristiane de Jesus Barbosa, Hermes Peixoto Santos-Filho, Onildo Nunes de Jesus.

"Severidade de doenças em maracujazeiro para identificação de fontes de resistência em condições de campo", Revista Brasileira de Fruticultura, 2013

Publication

&lt;1 %

62

"10.1007/BF00185650", CrossRef Listing of Deleted DOIs, 2011

Publication

&lt;1 %

63

Guillermo Martínez Pastur, Miriam Arena, Luis Hernandez, Néstor Curvetto, Eleonora Eliasco. "Histological events during rooting of (Fagaceae)", New Zealand Journal of Botany, 2005

Publication

&lt;1 %

64

Harshdeep Singh, Neelam Joshi. "Management of the aphid, Myzus persicae (Sulzer) and the whitefly, Bemisia tabaci (Gennadius), using biorational on capsicum under protected cultivation in India", Egyptian Journal of Biological Pest Control, 2020

Publication

&lt;1 %

---

65

Julien Dougoud, Stefan Toepfer, Melanie Bateman, Wade H. Jenner. "Efficacy of homemade botanical insecticides based on traditional knowledge. A review", Agronomy for Sustainable Development, 2019

Publication

---

<1%

---

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off