

2._The_Effect_of_Spirulina_on_Apoptosis_Through_the.pdf

by Harry Kurniawan Gondo

Submission date: 30-Nov-2021 12:53PM (UTC+0700)

Submission ID: 1716097068

File name: 2._The_Effect_of_Spirulina_on_Apoptosis_Through_the.pdf (405.48K)

Word count: 3410

Character count: 19313

The Effect of Spirulina on Apoptosis Through the Caspase-3 Pathway in a Preeclamptic Wistar Rat Model

Harry Kurniawan Gondo

Department of Obstetrics and Gynecology, Faculty of Medicine, Wijaya Kusuma University, Surabaya

Abstract

Objective: To determine the effects of spirulina expression on trophoblast cell apoptosis and caspase-3 expression in a preeclamptic wistar rat model. The preeclampsia model was administered to pregnant rats by interleukin-6 induction. **Materials and Methods:** This research is a Post Test Only Control Group Design. A total of 25 rats with preeclampsia model were divided in 36 groups, namely negative control group, positive control group, treatment group with IL-6 administration, and spirulina with 10 mg, 20 mg, and 40 mg/100 g body weight. Research has shown that spirulina reduces trophoblast cell expression through apoptosis in a rat model of preeclampsia. The positive control group showed the highest mean expression of apoptosis (62.38) and caspase-3 (804.33). The lowest average expression of apoptosis was 44.26, with 592.24 expressions of caspase-3 in group P3, Spirulina treatment at a dose of 40mg/day /kgBW **Conclusion:** Spirulina inhibits trophoblast apoptosis in preeclampsia through caspase-3 signaling.

Keywords: Apoptosis, caspase-3, interleukin-6, preeclampsia, spirulina, trophoblast.

INTRODUCTION

The risk of death from preeclampsia is about 300 times higher in developing countries than it is in developed [1]. It has reached 305 per 100,000 live births based on the 2015 Intercensus Population Survey. The newborn mortality rate is 15 per 1,000 live births, based on the Indonesian Health Demographic Survey [2]. Preeclampsia is characterized by the early onset of hypertension, edema, proteinuria, chronic immune activation and endothelial dysfunction, after 20 weeks of gestation [3]. This pregnancy complication is indicated by a systolic/diastolic blood pressure of 140/90 mmHg or higher, proteinuria of more than 300 mg or a protein-creatinine ratio higher than 0.3 in 24-hour urine [4]. In preeclampsia, trophoblasts experience increased oxidative stress on trophoblasts along with increased inflammatory response along with decreased antioxidants [5]. The pathogenesis of preeclampsia stems from cytotrophoblast differentiation, invasion of the superficial uterine cytotrophoblast followed by decreased maternal blood flow to the placenta [6].

Cysteine-dependent aspartate-driven proteases (caspases) are synthesized in cells as zymogens.

Apoptosis is associated with Caspase activation [7]. Apoptosis is defined as programmed cell death accompanied by morphological changes [23]. The molecular changes in apoptosis are the release of cytochrome c from the mitochondria and the activation of caspase 3 in the cytosol [9]. Intrinsic apoptosis-mediated permeabilization of the mitochondrial outer membrane leads to apoptosis formation, activation of caspase-9, and effector caspase. Caspase-3, caspase-7 affect the regulation of ROS production during intrinsic cell death [10]. Effector caspases including caspase-3 and -7 which are located in the cytosol, cytoskeleton, and nucleus, during apoptosis can cleave the P57 subunit in mitochondria which damages the electron transport chain and ends with a reorganization of the actin cytoskeleton and membrane blebbing [11].

Address for Correspondence: Harry Kurniawan Gondo, Department of Obstetrics and Gynecology, Faculty of Medicine, Wijaya Kusuma University, Surabaya
Email: gondo.hk@gmail.com

Submitted: 22th July, 2021

Received: 25th July, 2021

Accepted: 14th August, 2021

Published: 21st Oct, 2021

14

Access this article online

Quick Response Code:



Website:

www.jnsbm.org

DOI:

10.4103/jnsbm.JNSBM_12_3_2

12 is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Harry Kurniawan Gondo H K. The effect of spirulina on apoptosis through the caspase-3 pathway in a preeclamptic wistar rat model. J Nat Se Biol Med 2021;12:280-4.

Preeclampsia causes a maternal systemic inflammatory response syndrome involving leukocytes and endothelial cells, oxidative stress that damages maternal syncytiotrophoblast cells, and apoptosis [12]. Microdeposition of the syncytiotrophoblast-microvillus membrane occurs in systemic endothelial damage [13]. In addition, apoptosis in trophoblast cells is a clinical manifestation of preeclampsia [14]. A higher inflammatory response indicates an increased release of proinflammatory cytokines such as interleukin-6 (IL-6) [15].

Interleukin-6 (IL-6) was found to be widely expressed in the female reproductive tract as well as in pregnancy tissue, regulating the function of embryo implantation and placental development, immune adaptation to gestational tolerance [16]. Interleukin-6 is involved in chronic inflammation, and autoimmunity [17]. Furthermore, adaptive immunity is stimulated by excessive apoptosis. Monocytes are the largest type of leukocyte and show increased expression of IL-6 in the uterus, decidual cells, and the placenta [18].

Spirulina is a blue-green algae that contains 60%-70% protein, 15%-25% carbohydrates, 6%-13% fat, and 8%-10% fiber [19]. It has a microscopic filamentous cyanobacterium containing C-phycocyanin (CPC) [20]. Moreover, it induces apoptosis by DNA fragmentation and nuclear condensation. C-phycocyanin from spirulina reduces regulation of antiapoptotic proteins and upregulates proapoptotic proteins [21]. It functions as an antioxidant. Furthermore, it inhibits the development of immunomodulators [22,23]. In this study, we investigated the effect of spirulina on apoptosis in trophoblast cells and Caspase 3 expression in the IL-6-induced rat model of preeclampsia.

MATERIAL AND METHODS

This research has been reviewed by the appropriate ethics committee of the Faculty of Medicine, Wijaya Kusuma University, Surabaya No. 10198/SLE/FK/UWKS/2018). This study was a Test Only Control Group Design; twenty five rats were divided into 5 groups. These groups are similar to the research conducted by Soekidjo [24].

Negative control: pregnant rats were given aquadest
P0 (positive control): induced by IL-6 at a dose of 5 mg/day (intravenous [IV]),
P1 (treatment group 1): induced by IL-6 and given by spirulina at a dose of 10 mg/day (oral),
P2 (treatment group 2): induced by IL-6 and administered by spirulina at a dose of 20 mg/day (oral),
P3 (treatment group 3): induced by IL-6 and administered by spirulina at a dose of 40 mg/day (oral),

Two to three month old female rats weighing between

150 and 250 gm were separated from male rats for 2 weeks. Male and female rats were then put together into cages after 2 weeks. A vaginal plug-in female mice was examined after night. Pregnant rats were divided into five groups, each group containing five rats. Mice treated with AIN 93G food from Dyets Inc. and distilled water. The ingredients of AIN 93G are as follows: 40% flour, 20% casein, 13% maltodextrin, 10% sucrose, 7% soybean oil, 5% fiber, 3.5% minerals, 1% vitamins, 0.3% L-cystine, 0.25% choline bitartrate, and 0.0014% tert-butylhydroquinone. Mice were treated at a dose of 5mg/100 g using a wing needle in the tail vein for 3 days. Systolic and diastolic blood pressure was measured with a transducer three times (Kent Scientific CODA) at the Physiology Laboratory, Faculty of Medicine, Universitas Brawijaya. The average value of systolic and diastolic pressure was measured.

Spirulina is injected at a dose of 10 (P1), 20 (P2), and 40 (P3) mg/day. The control group and the treatment group were acclimatized with ketamine 0.5 mg orally after treatment. The rat placenta was placed in an Eppendorf tube containing 10% formalin. Then, the placenta was placed on a glass slide and stained using Bax polyclonal antibody with FITC and Rhodamine staining reagents. Observations were made using a confocal microscope. Expression of apoptotic cells and caspase-3 in placenta using Bax polyclonal antibody staining with FITC and Rhodamine.

RESULTS AND DISCUSSION

The results of this study were analyzed using comparative and homogeneity tests. Caspase-3 expression and apoptosis were measured using one-way ANOVA and Post Hoc Fisher's Least Significant Difference comparative test. As presented in Table 1, caspase-3 in the preeclampsia model is represented in P0 which has the highest level compared to other groups. Between treatment groups (P1, P2, P3), the expression of caspase-3 at P3 refers to the lowest level, close to P0.

Table 1. Caspase-3 Expression and Apoptosis

Caspase3	N	Mean	Standard Deviation	P
Negative Control	5	202.77	16.72	.003
P0	5	804.33	19.40	.080
P1	5	726.00	21.29	.001
P2	5	667.63	23.04	.060
P3	5	592.24	19.40	.009
Apoptosis				
Negative Control	5	46.32	6.08	.000
P0	5	62.38	13.28	.090
P1	5	62.96	2.38	.000
P2	5	52.10	12.42	.000
P3	5	44.26	9.41	.018

Apoptotic expression showed a slightly different trend with caspase-3 expression. The lowest level of apoptotic cells in the placenta was at P3 which was given with

spirulina at a dose of 40 mg/day (the highest dose). This indicates that spirulina at a dose of 40 mg/day can restore apoptosis in preeclampsia model rats (44,26) compared to negative controls (46,32). The spirulina treatment group at a dose of 10 mg/day (P1) significantly increased the rate of apoptosis.

Elevated levels of IL-6³⁸ have been reported in preeclamptic pregnancies. IL-6 increases endothelial cell permeability in pregnancy pathology by altering the shape and rearrangement of intracellular actin fibers. Furthermore, it reduces prostacyclin synthesis by inhibiting the cyclooxygenase enzyme and increasing the ratio of thromboxane A2 to prostacyclin. IL-6 also stimulates platelet-derived growth factors that cause preeclampsia^[25]. In a previous study, the induction of IL-6 in experimental animals led to an increase in the expression of apoptosis. Increased IL-6 production through monocyte activation and Th1 predominance induces inflammation due to dysregulation of the maternal immune system^[26]. In addition, it regulates immune and inflammatory responses^[27]. Therefore, endothelial cell dysfunction induces preeclampsia by increasing blood plasma levels and impairing the immune system^[28]. This is indicated by the average expression of caspase-3 in the positive control group. In the negative control, the expression of apoptosis was lower than the positive control.

This phenomenon is similar to a study^[29] entitled Phycocyanin ameliorates trophoblast apoptosis in a mouse model of IL-6-induced preeclampsia. Spirulina's bioactive content is carotenoids, phycocyanin, beta-carotene, zeaxanthin, SOD, polysaccharides, calcium, GLA, Vitamin B-12, Vitamin K1 and K2, it also contains high fiber and protein which are used for health maintenance. Spirulina is useful in preventing and controlling preeclampsia by reducing levels of proinflammatory cytokines^[20,22,29]. Spirulina offers a more complete nutritional profile than phycocyanin, which is why there is a hope that it will be better than both. The following are the histopathological results of caspase 3.




Figure 1. Histopathological results on Caspase-3

After administering spirulina to a pregnant rat model of preeclampsia that had been induced by IL-6, there was a decrease in apoptosis and caspase-3 expression as shown in Figure 1. The test was performed by counting the number of cells containing caspase-3 expression with 400x magnification in 10 fields of view. On examination, it was found that the expression of caspase-3 after administration of spirulina at treatment 3 doses of 40 mg/day/kg BW decreased apoptotic expression more than before the administration of spirulina.

The doses of P1 and P2 were significantly different ($p > 0.05$). In the P1 group, the average expression of caspase-3 was 726.00 and apoptosis 62.38. together with P2, the mean expression values of caspase-3 and apoptosis were 667.63 and 52.10, respectively. These results indicate that caspase-3 inhibition is significantly reduced by apoptosis. Spirulina therapy contains CPC, which suppresses caspase 3. Therefore, the apoptotic pathway can also be suppressed^[29,30]. We found that spirulina at a dose of 40 mg/day reduced trophoblast cell apoptosis. These results are in agreement with the theory which states that Spirulina plantesis may help the immune system to fight against infection. This alga also contains gamma-linolenic acid (GLA), and also provides alpha-linolenic acid (ALA), linolenic acid (LA), stearidonic acid (SDA), eicosapentaenoic (EPA), docosahexaenoic acid (DHA), arachidonic acid (AA). The vitamins contained in it are vitamins B1, B2, B3, B6, B9, B12, Vitamin C, Vitamin D²⁷ and Vitamin E. 2,3. In addition it is also a source of potassium, calcium, chromium, copper, iron, magnesium, manganese, phosphorus, selenium, sodium, and zinc which are used by the wider community to maintain a healthy body, especially the content of C-phycocyanin which is known as C-PC, a substance with potent cancer chemopreventive activity^[20] to prevent and control the occurrence of preeclampsia, by decreasing levels of pro-inflammatory cytokines^[29].

CONCLUSION

The administration of spirulina 40 mg/day orally was effective in repairing trophoblasts and reducing the expression of apoptosis and caspase 3 in an IL-6-induced rat model of preeclampsia.

SUGGESTION

Further studies should be conducted with a longer treatment period of spirulina administration to elucidate the significant effect of spirulina on the reduction of apoptosis. Additionally, spirulina's dose should be considered to determine its toxicity.

CONFESSION

This work was supported by the Faculty of Medicine, Wijaya Kusuma University, Surabaya, Indonesia.

REFERENCES

1. Ragasudha C, Madhavi AP, Sharon PS, Priya SS, Shehnaz S. A study of maternal deaths from preeclampsia and eclampsia in a tertiary care centre. *IAIM*. 2018; 5(1): 6-10. Available from: https://iaimjournal.com/wp-content/uploads/2018/01/iaim_2018_0501_02.pdf.
2. Mangla K, Hoffman MC, Trumpff C, O'Grady S, Monk C. Maternal self-harm deaths: an unrecognized and preventable outcome. *American journal of obstetrics and gynecology*. 2019; 221(4): 295-303. doi: <https://doi.org/10.1016/j.ajog.2019.02.056>.
3. Cornelius DC. Preeclampsia: from inflammation to immunoregulation. *Clinical medicine insights: Blood disorders*. 2018; 11: 1-6. doi: <https://doi.org/10.1177/2F1179545X17752325>.
4. Tenório MB, Ferreira RC, Moura FA, Bueno NB, de Oliveira ACM, Goulart MOF. Cross-Talk between Oxidative Stress and Inflammation in Preeclampsia. *Oxidative medicine and cellular longevity*. 2019; 2019: 8238727. doi: <https://doi.org/10.1155/2019/8238727>.
5. Mihiu D, Razvan C, Malutan A, Mihaela C. Evaluation of maternal systemic inflammatory response in preeclampsia. *Taiwanese Journal of Obstetrics and Gynecology*. 2015; 54(2): 160-66. doi: <https://doi.org/10.1016/j.tjog.2014.03.006>.
6. Li J, Ding Z, Yang Y, Mao B, Wang Y, Xu X. Lycium barbarum polysaccharides protect human trophoblast HTR8/SVneo cells from hydrogen peroxide-induced oxidative stress and apoptosis. *Molecular medicine reports*. 2018; 18(3): 2581-88. doi: <https://doi.org/10.3892/mmr.2018.9274>.
7. Parrish AB, Freel CD, Kornbluth S. Cellular mechanisms controlling caspase activation and function. *Cold Spring Harbor perspectives in biology*. 2013; 5(6): a008672. doi: <https://www.doi.org/10.1101/cshperspect.a008672>.
8. Tang D, Kang R, Berghe TV, Vandenabeele P, Kroemer G. The molecular machinery of regulated cell death. *Cell research*. 2019; 29(5): 347-64. doi: <https://doi.org/10.1038/s41422-019-0164-5>.
9. Lossi L, Castagna C, Merighi A. Caspase-3 mediated cell death in the normal development of the mammalian cerebellum. *International journal of molecular sciences*. 2018; 19(12): 3999. doi: <https://doi.org/10.3390/ijms19123999>.
10. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH. Caspase-9, caspase-3 and caspase-7 Have Distinct Roles During Intrinsic Apoptosis. *BMC cell biology*. 2013; 14(1): 1-9. doi: <https://doi.org/10.1186/1471-2121-14-32>.
11. Redza-Dutordoir M, Averill-Bates DA. Activation of apoptosis signalling pathways by reactive oxygen species. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*. 2016; 1863(12): 2977-92. doi: <https://doi.org/10.1016/j.bbamcr.2016.09.012>.
12. Sargent IL, Borzychowski AM, Redman CW. Immunoregulation in normal pregnancy and preeclampsia: an overview. *Reproductive biomedicine online*. 2006; 13(5): 680-86. doi: [https://doi.org/10.1016/S1472-6483\(10\)60659-1](https://doi.org/10.1016/S1472-6483(10)60659-1).
13. Wang W, Wang R, Zhang Q, Mor G, Zhang H. Benzo (a) pyren-7, 8-dihydrodiol-9, 10-epoxide induces human trophoblast Swan 71 cell dysfunctions due to cell apoptosis through disorder of mitochondrial fission/fusion. *Environmental Pollution*. 2018; 233: 820-32. doi: <https://doi.org/10.1016/j.envpol.2017.11.022>.
14. Mohammadpour-Gharehbagh A, Eskandari M, Sadegh MH, et al. Genetic and epigenetic analysis of the BAX and BCL2 in the placenta of pregnant women complicated by preeclampsia. *Apoptosis*. 2019; 24(3): 301-11. doi: <https://doi.org/10.1007/s10495-018-1501-8>.
15. Qu H-M, Qu L-P, Li X-Y, Pan X-Z. Overexpressed HO-1 is associated with reduced STAT3 activation in preeclampsia placenta and inhibits STAT3 phosphorylation in placental JEG-3 cells under hypoxia. *Archives of medical science: AMS*. 2016; 14(3): 597-607. doi: <https://dx.doi.org/10.5114%2Faoms.2016.63261>.
16. Prins JR, Gomez-Lopez N, Robertson SA. Interleukin-6 in pregnancy and gestational disorders. *Journal of reproductive immunology*. 2012; 95(1-2): 1-14. doi: <https://doi.org/10.1016/j.jri.2012.05.004>.
17. Ishihara K, Hirano T. IL-6 in autoimmune disease and chronic inflammatory proliferative disease. *Cytokine & growth factor reviews*. 2002; 13(4-5): 357-68. doi: [https://doi.org/10.1016/S1359-6101\(02\)00011-8](https://doi.org/10.1016/S1359-6101(02)00011-8).
18. Naugler WE, Karin M. The wolf in sheep's clothing: the role of interleukin-6 in immunity, inflammation and cancer. *Trends in molecular medicine*. 2008; 14(3): 109-19. doi: <https://doi.org/10.1016/j.molmed.2007.12.007>.
19. Wan D, Wu Q, Kuča K, Střelina. Nutraceuticals. *Elsevier*; 2021:959-74. doi: <https://doi.org/10.1016/B978-0-12-821038-3.00057-4>.
20. Ravi M, De SL, Azharuddin S, Paul SF. The beneficial effects of Spirulina focusing on its immunomodulatory and antioxidant properties. *Nutrition and Dietary Supplements*. 2022; 2: 73-83. Available from: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.1022.7279&rep=rep1&type=pdf>.
21. Belay A. The potential application of Spirulina (Arthrospira) as a nutritional and therapeutic supplement in health management. *J Am Nutraceutical Assoc*. 2002; 5: 27-48. Available

- from: <https://ci.nii.ac.jp/naid/10012910824/>.
22. Prabakaran G, Sampathkumar P, Kavisri M, Moovendhan M. Extraction and characterization of phycocyanin from *Spirulina platensis* and evaluation of its anticancer, antidiabetic and antiinflammatory effect. *International journal of biological macromolecules*. 2020; 153: 256-63. doi: <https://doi.org/10.1016/j.ijbi.2020.03.009>.
23. Adams M. Superfoods for optimum health: chlorella and spirulina. New York: Truth Publishing International. 2005.
24. Isaacs AN. An overview of qualitative research methodology for public health researchers. *International Journal of medicine and public health*. 2014; 4(4): 318-23. doi: <http://dx.doi.org/10.4103/2230-8598.144055>.
25. Sorokin Y, Romero R, Mele L, et al. Maternal serum interleukin-6, C-reactive protein, and matrix metalloproteinase-9 concentrations as risk factors for preterm birth < 32 weeks and adverse neonatal outcomes. *American journal of perinatology*. 2010; 27(08): 631-40. Available from: <https://www.thieme-connect.com/products/ejournals/abstract/10.1055/s-0030-1249366>.
26. Geldenhuys J, Rossouw TM, Lombaard HA, Ehlers MM, Kock MM. Disruption in the regulation of immune responses in the placental subtype of preeclampsia. *Frontiers in immunology*. 2018; 9: 1659. doi: <https://doi.org/10.3389/fimmu.2018.01659>.
27. NOYAN T, BURSAL E, ŞEKEROĞLU MR, DÜLGER H, KAMACI M. The serum interleukin-6 and tumor necrosis factor-alpha levels and their relationship with antithrombin-III and von Willebrand factor in preeclampsia. *Journal of the Turkish-German Gynecological Association*. 2006; 7(1): 39-44. Available from: <https://app.trdizin.gov.tr/publication/paper/detail/TmpVeU5UVTE>.
28. Lockwood CJ, Yen C-F, Basar M, et al. Preeclampsia-related inflammatory cytokines regulate interleukin-6 expression in human decidual cells. *The American journal of pathology*. 2008; 172(6): 1571-79. doi: <https://doi.org/10.2353/ajpath.2008.070629>.
29. Gondo HK, Kusworini H, Arsana W, Sarjoto T. Phycocyanin Ameliorate Trophoblast Apoptosis In Il-6-Induced Preeclamptic Rat Models. *International Journal of Pharmacognosy and Phytochemical Research*. 2017; 9(3): 424-27. Available from: <http://erepository.uwks.ac.id/id/eprint/2876>.
30. Garbuzova-Davis S, Bickford PC. Neuroprotective Effect of Spirulina in a Mouse Model of ALS. *Open Tissue Engineering and Regenerative Medicine Journal*. 2010; 3(1): 36-41. doi: <http://dx.doi.org/10.2174/1875043501003010036>.

2._The_Effect_of_Spirulina_on_Apoptosis_Through_the.pdf

ORIGINALITY REPORT

17%

SIMILARITY INDEX

14%

INTERNET SOURCES

8%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Wales Institute, Cardiff Student Paper	1%
2	www.medrxiv.org Internet Source	1%
3	www.tjogonline.com Internet Source	1%
4	dspace2.creighton.edu Internet Source	1%
5	ijmedph.org Internet Source	1%
6	Submitted to 2702 Student Paper	1%
7	Olivia K Travis, Cedar Baik, Geilda A Tardo, Lorena Amaral et al. "Adoptive Transfer of Placental Ischemia Stimulated Natural Killer Cells Causes a Preeclampsia - Like Phenotype in Pregnant Rats", American Journal of Reproductive Immunology, 2020 Publication	1%

8	ir.lib.uwo.ca Internet Source	1 %
9	www5.uclan.ac.uk Internet Source	1 %
10	Ji-Yeon Hwang. "Hypolipidemic effect of Salicornia herbacea in animal model of type 2 diabetes mellitus", Nutrition Research and Practice, 2007 Publication	1 %
11	invert.amphibians.org Internet Source	<1 %
12	www.e-journal.unair.ac.id Internet Source	<1 %
13	medcraveonline.com Internet Source	<1 %
14	mafiadoc.com Internet Source	<1 %
15	repository.unsoed.ac.id Internet Source	<1 %
16	Hana Katharina Heymes, Hanna Schröder, Andreas Follmann, Anja Sommer et al. "How Unexpected Is Out-of-hospital Sudden Cardiac Arrest? A Retrospective Cohort Study", Research Square Platform LLC, 2021 Publication	<1 %

17	mds.marshall.edu Internet Source	<1 %
18	www.kennispoort-verloskunde.nl Internet Source	<1 %
19	Submitted to Chulalongkorn University Student Paper	<1 %
20	repository.usd.ac.id Internet Source	<1 %
21	www.cellbiopharm.com Internet Source	<1 %
22	digilib.unila.ac.id Internet Source	<1 %
23	docplayer.net Internet Source	<1 %
24	jasn.asnjournals.org Internet Source	<1 %
25	www.jurnal.unsyiah.ac.id Internet Source	<1 %
26	www.mif-ua.com Internet Source	<1 %
27	www.nutrivital.co.uk Internet Source	<1 %
28	www.qscience.com Internet Source	<1 %

29

Submitted to AUT University

Student Paper

<1 %

30

D. J. Kominsky, R. J. Bickel, K. L. Tyler.
"Reovirus-Induced Apoptosis Requires
Mitochondrial Release of Smac/DIABLO and
Involves Reduction of Cellular Inhibitor of
Apoptosis Protein Levels", Journal of Virology,
2002

Publication

<1 %

31

rasayanjournal.co.in

Internet Source

<1 %

32

ttfnc.org

Internet Source

<1 %

33

www.endometriozisdernegi.org

Internet Source

<1 %

34

Mithat Aydın, Türkan Çelik. "Impact of the
Digital Literacy Courses Taken by the
Prospective Social Studies Teachers by
Distance Learning on Digital Citizenship
Skills", Research on Education and Media,
2020

Publication

<1 %

35

ajp.amjpathol.org

Internet Source

<1 %

36

paperity.org

Internet Source

<1 %

37	research.vu.nl Internet Source	<1 %
38	Papa, A.. "Cytokine levels in Crimean-Congo hemorrhagic fever", Journal of Clinical Virology, 200608 Publication	<1 %
39	myassignmenthelp.com Internet Source	<1 %
40	portal.research.lu.se Internet Source	<1 %
41	portlandpress.com Internet Source	<1 %
42	Edward Mullins. "Changes in the maternal cytokine profile in pregnancies complicated by fetal growth restriction", American Journal Of Reproductive Immunology, 04/2012 Publication	<1 %
43	Linyan Wang, Juan Zhang. "Long intergenic ncRNA 00473 improves the invasion of trophoblastic cells via miR-16-5p", Pregnancy Hypertension, 2020 Publication	<1 %
44	Yin-Xu Zhang, Xiao-Mei Liu, Jing Wang, Jun Li, Ying Liu, Hua Zhang, Xue-Wen Yu, Ning Wei. "Inhibition of AKT/FoxO3a signaling induced PUMA expression in response to p53-	<1 %

independent cytotoxic effects of H1: A
derivative of tetrandrine", Cancer Biology &
Therapy, 2015

Publication

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	Off		