

# 15

*by* Emilia D

---

**Submission date:** 13-Jan-2020 06:50PM (UTC-0800)

**Submission ID:** 1241690230

**File name:** 15.\_PROSEDING.WM.2010.docx (22.6K)

**Word count:** 1723

**Character count:** 9173

**THE EFFECT OF KATU INFUSE (*SAUROPIUS ANDROGYNUS*)  
LEAVES ON SPERMATOGENESIS TESTIS  
SEMINIFEROUS TUBULES OF MICE  
(*MUS MUSCULUS*)**

Mas Loegito<sup>1</sup>, Mas Mansyur<sup>2</sup>, E Devi Dwi Rianti<sup>3\*</sup>

<sup>1</sup>Department of Biology molleculer, <sup>2</sup>Faculty of Medicine, Wijaya Kusuma University Surabaya.

<sup>2,3</sup>Department of medical physisc, Faculty of Medicine Wijaya Kusuma University Surabaya,

\*corresponding author, e-mail: [mbak.devi@gmail.com](mailto:mbak.devi@gmail.com)

**Abstract** : Katu leaves contain steroid substances, Steroid substances in blood can deinhibids indirectly hypothalamus activity and inhibits secretion of GnRhH. Reduction of GnRH in the blood from hypothalamus cause inhibits FSH and LH secretion from pituitayr gland, resultion in the disruption of spermatogenesis proces. Material consist of infuse sauopus androgyans leaves and 60 male mice of 8 weeks old. With 20-30 grams body weight, were devided into control. Group and streatment groups. Data were analyzed using Anova test. If the results indicated significant of deifferent of 95%, the analyzed was continued with Duncan test. The result of this exptiment are: The heightening of the infuse concentration of anropus androgygnus leaves were treated 0,5 cc infuse per oral / per day, caused decreases of the cell association number, the spermatosit number, and the spermatid number in testis seminiferous tubules of mice. So the infuse of samropus andrognus leaves will be consumed as contraceptive substance for men.

**Keywords** : the cell association number, the spermatosit number, the spermatid number, the infus concentration.

## INTRODUCTION

Katu (*Sauropus androgygnus*, Merr) is a herb belonging to the family Euphorbiaceae. The leaves of this plant are usa as herbal medication for fever, franbosia, ulcer venereal diseaces, influence breast milk production and consumed as vegetable and to colour of food (Heyne K,1987, Soedarman and Harsono, 1968).

Katu leaves contain steroid substances (Pradjonggo, 1983). Steroid substances in blood cause inhibids indirectly hypthalamus activity and inhibids secretion of GnRH (Evan, RM,1988). Steroid substances will gives negative feedbact action and inhibids secretion of gonatropin hormone (wade, 1996).

Reduction of GnRH in the blood from hypothalamus cause inhibids FSH and LH secretion from pituitary gland, resulting in the disruption of spermatogenesis proses (Wignjosastro, 1997). Based on these facts, we want to investigation to the effect of infuse katu leaves to spermatogenesis process in tubules seminifefous testis of mice, especially on the cell associatiopn number, spermosit number, and spermatid number. Wether these are correlation between increases of infuse consentration and inhibits spermatogenesis in its testis.

## MATERIALS AND METHODS

This research was conducted at the Labolatory of reproduction Biology, Faculty of Sains and Tecnology, Airlangga Universty, Surabaya.

Mereial consist of infuse katu leaves and 60 male mice of 8 weeks old with 20-30 grams body weight, were devided into six groups, i.e.control group were given 0,5 cc per oral per day aquadest; P1 were given 0,5 cc per oral per day 5% infuse; P2 were given 0,5 cc per oral per day 10% infuse; P3 were given 0,5 cc per oral per day 15 % infuse; P4 were given 0,5 % per oral per day 20 % infuse; P5 were given 0,5 cc per oral per day 25 % infuse. Treatment was carried out for 35 days. The testis were removed to exemine the structure of the histological tubules seminiferous of the testis.

Data were obtained by enumerating the number of cell association, spermatosit, and spermatid in testis seminiferous tubules of mice from controle and each treatment gruops. Data were analyzed was continued with Duncan Test (Duncan Multiple Range Test).

## RESULTS

The experiment dates before of anove test, we make of normal distribution test with “ One Sampel Kolmogorof – Smirov Test”. The results symp sig test more than 0.60, so the normal distribution test so we can do anova test. Base on the anove test of the date there are different of cell association number in tubules seminiferous on any threatment groups see table 1 and 2.

Table1. The result anove test of different cell association number in seminiferous tubules for any threatment and controle

Source	Sum Square	df	Mean Square	F	Sig.
Corrected Model	1238.908	5	247.782	98.615	0.000
Intercept	1811.702	1	1811.702	721.043	0.000
Treatment	1238.908	5	247.782	98.615	0.000
Error	135.681	54	2.513	-	-
Total	3186.290	60	-	-	-
Corrected total	1374.589	59	-	-	-

Table 2. The date of the influence of infuse katu leaves which defferent of mean cell asscition number

No	Threatment	Mean of cell association number
1	Control	14.76±3.54
2	P1 5%	7.24±0.80
3	P2 10%	4.77±0.70
4	P3 15%	2.46±0.85
5	P4 20%	1.98±0.77
6	P5 25%	1.80±0.27

Base on the anove test of the date there are different of spermatocyte number in seminiferous tubules on any threatment groups see table 3 and 4.

Table 3. The result anove test of different spermatocyte of number in seminiferous tubules for any threatment and controle

Source	Sum Square	df	Mean Square	F	Sig.
Corrected Model	1302.163	5	260.433	82.067	0.000
Intercept	12229.393	1	12229.353	3853.710	0.000
Treatment	1302.163	5	260.433	82.067	0.000
Error	171.364	54	3.173	-	-
Total	13702.920	60	-	-	-
Corrected total	1473.527	59	-	-	-

Table 4. The date of the influence of infuse katu leaves which defferent of mean spermatocyte number.

No.	Threatment	Mean of cell association number
1	Control	22.72± 2.74
2	P1 5%	17.08±1.63
3	P2 10%	14.48±1.32
4	P3 15%	12.44±1.58
5	P4 20%	10.32±1.03
6	P5 25%	8.62±1.29

Base on the anove test of the date there are different of spermatid number in tubules seminiferous on any threatment groups see table 5 and 6.

Table 5. The result anove test of different spermatid number in seminiferous tubules for any threatment and control

Source	Sum Square	df	Mean Square	F	Sig.
Corrected Model	1382.203	5	76.441	5.714	0.000
Intercept	10864.913	1	10864.913	812.202	0.000
Treatment	382.203	5	76.441	5.714	0.000
Error	722.364	54	13.377	-	-
Total	11969.480	60	-	-	-
Corrected total	11104.567	59	-	-	-

Table 6. The date of the influence of infuse katu leaves which defferent of mean spermatid number

No.	Threatment	Mean of cell association number
5	Control	15.94± 4.71
2	P1 5%	15.66±2.76
3	P2 10%	15.22±4.62
4	P3 15%	13.89±2.23
5	P4 20%	10.40±3.88
6	P5 25%	9.64±2.89

## DISCUSSION

The result of the experiment are following:

1. Based on anova test, the number of cell associatioan in testis seminiferous tubules between groups of control, P1, P2, P3, P4, and P5 was very significantly different on level 95% ( $p = 0.00 < 0.05$ ). But the number of cell association of P3, P4, and P5 groups were not significantly different on 95% ( $p = 0.00 < 0.05$ ) from one another by Duncan's test, but all of the groups significantly different toward control group, P1, and P2 groups. Beside of those results. It was found that the number of cell association on control groups, P1, and P2 groups were significantly different on level of significant 95% ( $p = 0.00 < 0.05$ ).
2. There were high correlation between the number of cell associatioan toward the increase of infuse katu concentration were treated to mice by index correlation  $0.843 > 0.600$ .
3. There were significant difference the spermatosit number in testis seminiferous tubules of mice between all groups on level of significant 95% ( $p = 0.00 < 0.05$ ).
4. There were correalation between the spermatosit number in testis seminiferous tubules of mice toward the increase of infuse concentration were treated to mice in any groups by index correlation ( $r$ )  $0.914 > 0.600$ .
5. The different of the spermatid number in testis seminiferous tubules of mice between control group, P1, P2, and P3 were not sinificant; in such a manner for P1 to P2, P2 to P3, not sinificant on level of significant 95% ( $p = 0.00 < 0.05$ ). but spermatid number in control group, P1, P2, and P3 were different toward the spermatid number in P4 and P5 groups on level of significant 95% ( $p = 0.00 < 0.05$ ).
6. There were light correlation between the spermetid number in tubules seminiferous toward the increases of infuse concentration were treated to mice in any groups by index correlation ( $r$ )  $0.553 < 0.60$ .
7. There were high correlation between cell association number toward spermatosit number in testis seminiferous tubules of mice by index correlation  $0.839 > 0.60$ .
8. There were light correlation between cell association number toward spermatid number in testis seminiferous tubules of mice by index correlation ( $r$ )  $0.408 < 0.60$ .
9. There were light correlation between spermatid number toward spermatid number in testis seminiferous tubules of mice by index correlation ( $r$ )  $0.490 < 0.60$ .

## CONCLUSION

The conclusion of this experiment are following:

1. The infuse of *Sauropus andogynus* leaves affected to decreaces of the cell association number in testis seminiferous tubules of mice which treated 0.5 cc infuse per oral per day.
2. The heightening of the infus consentration of *Sauropus andogynus* leaves were treated 0.5 cc infus per oral per day, caused decreases of cell association number in testis seminiferous tubules of mice.
3. The infuse of *Souropus androgynus* leaves affected to decreases of spermatosit number in testis seminiferous tubules of mice which treated 0.5 cc infus per oral per day.
4. The heightening of the infuse consentration of *Saouropus androgynus* leaves were treated 0.5 cc infuse per oral per day, caused decreases of the spermatosit number in testis seminiferous tubules of mice.
5. The infuse of saouropus andorynus leaves affected to decreases of the spermatid number in testis seminiferous tubules of mice.
6. The heightening of the infuse consentration were treated to the male mice, caused to decreases of the spermatid number in testis seminiferous tubules.

## SUGGESTION

Based on the results, we suggest that if the infuse of *Saouropus andogynus* leaves will be consumed as contraceptive substance for men, there are studies shuold be conducted about:

1. Optinum dose of *Saouropus andogynus* leaves for men.
2. Infuse dose of *Saouropus andogynus* leaves which has reversible effect to the spermatogenesis procces for men.

## REFERENCES

- Evans R.M., 1988, The Steroid and Thyroid Hormone Receptor Superfamily, *Science*, 240, 889-896.
- Heyne K., 1987, Tumbuhan Berguna Indonesia II, Cetakan Ke I, Jakarta: penerbit Yayasan Sarana Wana Jaya.
- Pradjonggo T.S., 1983, Penelitian Pendahuluan Pengaruh Daun Katu (*Sauropus adrogynus*, Merr) Terhadap Gambaran Histologi Kelenjar Susu Mencit Betina Yang Menyusui, Skripsi, Fakultas Farmasi, Unair Surabaya, 1983.
- Sudarman M., Harsono R., 1985, Cabe Payung Warisan Nenek Moyang I, PN. Balai Pustaka.
- Wade G.N., 1996, Control of Fertility by Metabolic cues, *Am.J.Physiology*, 270, 1-19.
- Wignyosastro H., 1997, Ilmu Kandungan, Edisi Ke-2, Jakarta: Yayasan Bina Pustaka Sarwonoprawirohardjo.

## ORIGINALITY REPORT

5%

SIMILARITY INDEX

4%

INTERNET SOURCES

1%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.scribd.com](http://www.scribd.com)

Internet Source

2%

2

[erepository.uwks.ac.id](http://erepository.uwks.ac.id)

Internet Source

1%

3

[www.seceng.informatik.tu-darmstadt.de](http://www.seceng.informatik.tu-darmstadt.de)

Internet Source

1%

4

Submitted to Universitas Negeri Semarang

Student Paper

1%

5

Submitted to Universitas Jenderal Soedirman

Student Paper

1%

6

Submitted to Cita Hati Christian High School

Student Paper

&lt;1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On