

# 2. Repelence of Bintaro Plant Extract (Cerbera Manghas) Against Pod-sucking Insects (Riptortus Linearis) (Hemiptera

*by Dwi Haryanta*

---

**Submission date:** 11-Jan-2021 11:59PM (UTC-0800)

**Submission ID:** 1486214043

**File name:** has\_Against\_Pod-sucking\_Insects\_Riptortus\_Linearis\_Hemiptera.pdf (1.38M)

**Word count:** 5893

**Character count:** 28424

# Repelence of Bintaro Plant Extract (*Cerbera manghas*) against pod-sucking insects (*Riptortus linearis*) (Hemiptera)

Dwi Haryanta<sup>1</sup>, Achmadi Susilo<sup>2</sup>, Tatuk Tojibatus Sa'adah<sup>3</sup>

<sup>1,2,3</sup>Faculty of Agriculture, University of Wijaya Kusuma Surabaya

Email: [dwi.haryanta@yahoo.com](mailto:dwi.haryanta@yahoo.com)

46

Received: November 11, 2020. Revised: December 10, 2020. Accepted: December 11, 2020.

Published: December 14, 2020.

65

**Abstract**—The mechanism of the organic pesticides of Bintaro plant extract (*Cerbera manghas*) in reducing the insect pest population needs to be continuously studied. The study aims to determine the Repelence (ability to resist) of bintaro plant extracts against brown ladybugs (*Riptortus linearis*). The factorial experiment with the first-factor treatment was extracted Bintaro plant organs, namely leaf extract, stem bark extract, fruit pulp extract and bintaro seed extract, while the second treatment factor was the extract concentration (w/v), namely a concentration of 2,5%, a concentration of 7,5%, and a concentration of 12,5%. The experimental variables were the chemical compound content of each Bintaro plant part extract, insect mortality, length of each instar and stadia, and development into the next instar or stadia. The research results showed that Bintaro plants (leaves, fruit, seeds and bark) contain ingredients that can be used as organic pesticides. Bintaro plant extract has an effect on the mortality and development of *Riptortus linearis* insects. The results of the Bintaro plant extract test on the mortality and development of *Riptortus linearis* insects are less significant if it will be used to suppress the population, but the application of organic pesticides in the field is proven to be able to suppress the population, reduce the level of crop damage and increase crop production.

**Keywords**— Agricultural Biology, Entomology, Zoology, Botanology, Ecology

## I. INTRODUCTION

Pod-sucking pests (*Riptortus linearis*) attack post-flowering soybean plants, are important pests that can cause a decrease in yield both in quality and [1], [2]. The nymphs and adults damage the soybean pods by sucking the liquid of the seeds on the young pods, causing deformation of the seeds, the young pods drying out and falling out, which in turn causes a decrease in production both in quality and quantity [3]. The

pod-sucking pest, *Riptortus linearis* is the most dangerous pest among post-flowering soybean pests with an average intensity of 61,37% of the total affected pods and 61,67% of the total seeds affected. [4-5]. Riptortus insects destroy the pods and seeds of cowpeas. Adult insects and nymphs suck the liquid from the young shoots and pods, causing the plant shoots to dry out and the young pods to drop [6].

The pod sucking pest control that is relatively safe, cheap, and easy is to use botanical/vegetable pesticides. Depend on synthetic pesticides should be avoided because of their detrimental effects on human health, the environment, and the development of resistant strains of pathogens and pests. Botanical pesticides are effective in dealing with various plant pests, cheap, easily biodegradable, have a variety of ways of working, easily available sources and have low toxicity to non-target organisms [7]. The use of vegetable pesticides is effective in controlling sucking pests on cotton, and is proven to be non-toxic to natural enemies of pests and environmentally friendly [8]. Several plants that have been studied as biopesticides among others the plant extracts of *Spondias mombia*, *Momordica charantia*, *Mitrocarpus villosus* and *Chenopodium ambrosioides* for pod sucking pests. [9], the plant extracts of *Xylopia aethiopica* (Dunal) A. Rich., *Zingiber officinale* L. and *Capsicum annum* L.) for pest control of *Maruca vitrata* Fab. (Lepidoptera: Pyralidae) and *Clavigralla tomentosicollis* Stal (Hemiptera: coreidae) [10], plant extracts of *Annona muricata* and *Piper guineense* are very effective at controlling thrips [11], papaya and babadotan leaf extract for *Riptortus linearis* insects [5], *Azadirachta indica*, *Cymbopogon nardus* and the rhizome of *Alpinia galanga* to control soybean pests [12], cashew nut shells, Albasia bark, Siamese weed, candle bush leaves, for cacao pod borer [13], kamandrah seed extract and spinosad biopesticide for brown ladybugs [14],[6],[15], water extract of mango parasite of *Dendrothoe pentandra* (L.) Miq) affects the biology of *Spodoptera litura* F [16]. *Annona squamosa* seed powder, castor seed powder with *L. lecanii* fungi are able to control brown ladybugs [17].

Vegetable pesticides are less toxic to predators of coccinellidae than synthetic pesticides [18]. The efficacy of biopesticides is influenced by variations in the content of active ingredients in plants, concentrations, and its manufacturing process [19].

In Bintaro plants are found antioxidants, cytotoxics, antimicrobials, stabilizers of membrane activity and reduction of thrombolytic activity which can be used as a reference for *Cerbera odollam* plants as biopesticides. [20]). Bintaro plant extract has the potential to control sucking pests. Bintaro plant extracts have been studied to control the *Spodoptera litura* pest [21-22], to control caterpillars in tamarind [23], for the *Aedes aegypti* mosquito [24]. The methanol extract of *Cerbera manghas* fruit, twigs and leaves was able to control *Panonychus citri* directly [25].

The mechanism of organic pesticides (plant extracts) in reducing pest insect populations needs to be continuously studied. Organic pesticides (plant extracts) have no effect on insect pests in obtaining their hosts, as the results of research by [26] which concluded that the giving of Bintaro leaf extract (*Cerbera manghas*) had no effect on the response of brown ladybugs (*Riptortus linearis*, Hemiptera) behavior in getting the host. According to [8] that vegetable pesticides begin to decrease its toxicity after 96 hours. However, [25] effective pest reduction was noted for up to one week. The use of plant compounds (essential oils, flavonoids, alkaloids, glycosides, esters and fatty acids) has an anti-insect effect and is important as an alternative to chemical compounds in pest control in various ways, namely repellents. [27]. This study aims to determine the Repelence (ability to resist) of bintaro plant extracts against brown ladybugs (*Riptortus linearis*).

## II MATERIAL AND METHOD

The research on "The Repelence of Bintaro Plant Extracts (*Cerbera manghas*) against pod-sucking insects (*Riptortus linearis*, Hemiptera)" is an integral part of the research of [22] on Response of *Riptortus linearis* towards the application of Bintaro (*Cerbera manghas*) leaf extract. Bintaro plant material is developed into 4 parts, namely leaves, bark, pulp and seeds. The research was conducted at the Plant Protection Laboratory, Faculty of Agriculture, Wijaya Kusuma University, Surabaya.

## III RESULTS AND DISCUSSION

**Phytochemical Content of Bintaro Plant Organs as Biopesticides Material** - The results of phytochemical analysis of Bintaro plants showed that all parts of the plant contained polyphenols, saponins, steroids, flavonoids and tannins. The part of the Bintaro fruit pulp plant contains saponins of 3,01% the highest compared to other parts, the leaves contain 3,04% polyphenols the highest compared to other plant parts and the bark contains 3,08% tannins the highest compared to other plant parts. The data of phytochemical content of Bintaro plants more details presented in Table 1.

**The Mortality of *Riptortus linearis* F. Insects**-The results of the analysis of variance on the riptortus insect mortality data concluded that there was an interaction between the extracted

plant organ treatment factors and the concentration treatment, so that the presented is a combination treatment between two treatment factors. The experiment used a two-day old first instar nymph, which was small, soft and still very sensitive. Insect mortality occurred starting on the first day although only occurred in two treatments. The mortality rate continues to increase and the highest mortality is in Bintaro seed extract treatment with a concentration of 7,5%. On the 15th day of observation of the control treatment, no died yet, the mortality rate in Bintaro plant extract treatments increased with the highest mortality rate of 58,33% in the leaf extract treatment with a concentration of 2,5%. There are still many treatments with a mortality rate below 25%, some are even 0% (still all alive). Observations were conducted up to day 24, when all the insects had become adults. Mortality rate data for riptortus insects which given the treatment of the concentration of the Bintaro plant part extract presented in Tables 2a, 2b, and 2c.

Riptortus insects are very active, rarely stay in long bean pods (feed). The insect takes the food by jabbing the stylet then sucks the liquid of long bean pods. Bintaro extract that exists on the surface of long bean pods is less likely to be sucked into the insect's body. Physical contact between insects and Bintaro extract can occur when the insects are on the surface of long bean pods for a relatively long time.

### The development of *Riptortus linearis* F. Insects from instar 1 to imago-

In the ages of riptortus insects instar 2 to instar 5 there were no significant difference among the treatments, including comparison with control. In the ages of adult riptortus insect there were significant differences among the treatments. Riptortus insects with the treatment of Bintaro seed extract with a concentration of 7,5% and 12,5% have a relatively short age, namely 35,33 days for a concentration of 7,5% and 26,67 days for a concentration of 12,5%, while the other treatments its ages were over 40 days old. Data on the mean age for each instar, preadult/mature stadia and adult stadia of riptortus insects for each treatment are presented in Table 3.

The age of the riptortus insects from hatching eggs to death due to old age was significantly different among the treatments. Riptortus insects with the treatment of Bintaro seed extract with a concentration of 7,5% were able to live for 47,67 days, with a concentration of 12,5% were able to live for 44,33 days shorter than other treatments with an average of more than 55 days. The age of the riptortus insects until the mortality 50% significantly differ among the treatments. The fastest 50% mortality occurred in the treatment of Bintaro leaf extract with a concentration of 2,5%, namely at the age of 15 days, then the seed extract treatment with a concentration of 7,5% died 50% at the age of 19 days. Data on the total age and time of 50 percent mortality of the *Riptortus linearis* insects with the treatment of the concentration of extracts of leaf, bark, fruit and seeds of Bintaro are presented in Table 4.

The treatment of Bintaro plant extract did not significantly affect the development of riptortus insects from first instar to third instar, and into fourth instar, but had a significant effect on the development into fifth instar and into imago. The percentage of the development of the first instar caterpillar to imago, the lowest amounted to 62,50% occurred in the seed extract treatment with a concentration of 7,5% which was significantly different compared to the control treatment of



the development of first instar to imago amounted to 100%. Data on the development of *Riptortus linearis* insects from the first instar to the next until the imago with the treatment of various concentrations of the extracts of leaf, bark, fruit and seeds of Bintaro presented in Table 5.

**Discussion**—The treatment of the extract of leaf, fruit and stem bark of bintaro did not kill the *Riptortus* insects, and only Bintaro seed extract caused 33,33% *Riptortus linearis* insect mortality on the seventh day after treatment. The results of this study are in line with the results of research by [5] that the application of the extracts of papaya leaves and babadotan leaves with a concentration range of 30% - 70% causes *Riptortus linearis* mortality by 21,57% on 7 days after application and this not different from the control. If the Bintaro extract is eaten by insects it will cause death. Susilo, et al. 2019 [26] concluded that the provision of Bintaro leaf extract (*Cerbera manghas*) had no effect on the behavioral response of the brown ladybugs Insect (*Riptortus linearis*, Hemiptera) in getting a host and did not cause the significant mortality. Bintaro plant extract that is sprayed on the feed does not enter the digestive system because the *riptortus* takes food by sticking the stylet in the tissue. According [28] Bintaro leaf extract granules (*Cerbera odollam* Gaertn.) can cause the mortality of armyworms (*Spodoptera litura* Fab.) because the extract material is eaten and enters the digestive system of the caterpillar, the higher the granular concentration of Bintaro leaf extract (*Cerbera odollam* Gaertn.), then the higher the mortality of armyworms (*Spodoptera litura* Fab.).

Bintaro plant extract treatment has no significant effect on the development period of the preadult/mature phase, but seed extract can shorten the lifespan of adult insects. Bintaro plant extract treatment had an effect on the development of pre-adult/mature insects (nymph) into adult insects (imago). In line with the research results of [22] which concluded that Bintaro plant extract reduced caterpillar weight, inhibited ecdysis causing failure of 2nd instar development to 3rd instar and development into pupa [39] from *Spodoptera litura*. Plant compounds in the form of essential oils, flavonoids, alkaloids, glycoside [28] esters and fatty acids have anti-insect effects in various ways, namely repellents, food inhibitors/antifeedants, toxic, growth inhibitors, chemosterilants, and attractants [27].

The use of plant extracts proven to be able to control pests in the field, although [16] the mortality test is not as effective as synthetic pesticides. Petroleum ether extracts of *A. muricata* and *P. guineense* seeds are very effective in controlling thrips and pod borer pests in beans. [11]. The combination of seed powder botanical pesticide of *Annona squamosa*, Castor seed powder with the entomopathogenic fungus of *L. lecanii* was able to control *Riptortus linearis* [17]. Leaf extracts of *Tephrosia vogelii* and *Azadirachta indica* can reduce the intensity of pod damage by sucking [55] and borer pests and increase soybean seed yields. [1]. Extracts of *Azadirachta indica* A. Juss. and *Chromolaena odorata* (L.) were able to suppress the damage by pod sucking pests and increase the yield of cowpea seeds. [29]. Neem seed [3] oil extract and garlic were able to reduce the population of *Riptortus dentipes* in cowpea and bitter [31] on pods during the two rainy seasons [3]. Plant extracts of *Spondias mombia*, *Momordica charantia*, *Mitrocarpus villosus* and *Chenopodium ambrosioides* can be

used as biopesticides that effectively control pod-sucking insect [44] in cowpeas [9]. The treatment of water extracts from *Azadirachta indica* A. Juss., *Chromolaena odorata* (L.) and *Ricinus communis* (L.) plants can suppress the pod sucking pest population of *Riptortus linearis* and increase cowpea production [29]. The water extracts of *Hyptis suaveolens*, *Azadirachta indica*, *Manihot esculenta*, *Thevetia nerifolia* and *Cymbopogon nardus* plants were able to significantly reduce the pod sucker population. [15]. Vegetable insecticide formulations have the prospect to be developed further and have the potential to reduce the use of synthetic insecticides [13]. Plant extracts work slowly, but can be considered as an alternative pesticide that is environmentally friendly and applicable [28] developing countries [30]. Vegetable insecticides only affect target insects, do not kill beneficial natural enemies, provide residue-free food and a safe environment for humans [27].

#### IV CONCLUSION

Bintaro plants (leaves, fruit, seeds and bark) contain ingredients that can be used as organic pesticides. Bintaro plant extract has an effect on the mortality and development of *Riptortus linearis* insects. The results of the Bintaro plant extract test on the mortality and development of *Riptortus linearis* insects are less significant if it will be used to suppress the population, but the application of organic pesticides in the field is proven to be able to suppress the population, reduce the level of crop damage and increase crop production. From this phenomenon, it should be assumed that there is a possibility that plant which sprayed with organic pesticides tend to be avoided by insect pests. Research on the Repelence of Bintaro plant extracts against insect pests needs to be developed in the field.

There are needs for empirical proof of the existence of absolute Repelence of biopesticide ingredients, namely insect pests are really not interested in visiting plants that containing biopesticides so that the population is low, or there is an apparent/quasi Repelence, namely insect pests actually come to plants but because they are exposed to biopesticide treatment, the insects do not develop so that their population low

#### V ACKNOWLEDGEMENTS

Publication [62] material is part of the results of excellent basic research of higher education/ penelitian dasar unggulan perguruan tinggi (PDUPT) the 3rd year [17] in the 3-year plan. We would like to take this opportunity to thank the Directorate of Research and Community Service, Directorate General of Research and Development Strengthening, Ministry of Research and Technology who has provided funds for the implementation of research.

#### References

- [1] Hendriani, Latifah, and Nisa, A. Efficacy Some Botanical Insecticide [30] for Controlling Pest Pod Sucking in Soybean Fields. Jurnal Agrista Vol. 17 No. 1, 2013.

- [2] Krisnawati, A., and Adie, M.M. 2018. Evaluation of Soybean Resistance to Pod-Sucking Bug, *Riptortus linearis* and Performance of its Agronomic Characters. *3* *osaintifika* 10 (1) (2018) 213-222.
- [3] Agri, M. M., Maina, Y. T. and Sharah, H. A. 2013. Control of Pod-sucking bug *Riptortus dentipes* (Hemiptera: Alydidae) of Cowpea with aqueous Plant Extracts and Cymbush Super EC in Maiduguri, Nigeria. *Journal of Health, Medicine and Nursing- An Open Access International Journal* Vol.2 2013.
- [4] Bayu, M.S.Y.I. (2015). Attack level of various pod pests on soybean germplasm. National Seminar Proceeding of Masyarakat Biodiversitas Indonesia. 1: 878-883. [12] [Indonesian].
- [5] Amalia, E. R., Hariri, A.M., Lestari, P. dan Purnomo. 2017. Uji MORTALITAS PENGHISAP POLONG KEDELAI (*Riptortus Linearis* F.) (HEMIPTERA : ALYDIDAE) SETELAH APLIKASI EKSTRAK DAUN PEPAYA, BABADOTAN DAN MIMBA DI LABORATORIUM. *Jurnal Agrotek Tropika* J. Agrotek Tropika. Vol. 5, No. 1: 46 – 50, Januari 2017 ISSN 2337-4993 5(1): 46-50, 2017.
- [6] Dahal, S., Gautam, B., Sharma, B., Neupane, K., [29] Adel, S., Sedhai, S., Tiwari, G., and Ojha, L.N.2020. Management Options Against Pod Borer (*Maruca Testulalis* Geyer) And Pod Sucking Bugs (*Riptortus Dentipes*) Of Cowpea (*Vigna Unguiculata* (L.) WALP.). *Tropical Biodiversity (TRAB)* 1(1) (2020) 24-30.
- [7] Lengai, G.M.W. , James W. Muthomi , Ernest R. Mbega. 2020. Phytochemical activity and role of botanical pesticides in pest management for sustainable agricultural crop production . *Scientific African* 7 (2020) 4 0239.
- [8] Noonari, A.M., Abro, G.H.I, Khuhro, R.D. and Buriro, A.S. 2016. Efficacy of Bio-Pesticides for Management of Sucking Insect Pests of Cotton, *Gossipium hirsutum* (L.). *Journal of Basic & Applied Sciences*, 2016, 12, 306- [11] .
- [9] Adesina, J.M. 2013. Field assessment of insecticidal efficacy of some plant aqueous extracts in reducing cowpea pod-sucking bug *Acanthomia tomentosicollis* Stål (Hemiptera: Coreidae) infestation and damage. *Archives of Phytopathology and Plant Protection*, 2014 9 <https://doi.org/10.1080/03235408.2013.871106>.
- [10] Oparaeke A. M., Dike M. C., Amatobi C. I. 2005. Field evaluation of extracts of five Nigerian spices for control of post-flowering insect pests of cowpea, *Vigna unguiculata* (L.) Walp. *Protect. Sci.*, 41: 14–20.
- [11] Ugwu, JA. 2013. Insecticidal activity of some botanical extracts against legume flower thrips and legume pod borer on cowpea *Vigna unguiculata* L. Walp *The Journal of Basic and Applied Zoology* (2020) 81:13 <https://doi.org/10.126/s41936-020-00153-3>.
- [12] Asmanizar, Syamsafitri, Arianto, A. 2019. Response of Insect Pests to Botanical Insecticide Application on Leaf and Pod [34] bean (*Glycine max* L.) of Organic Farming System. *Advances in Biological Sciences Research, volume 8* International Conference and the 10th Congress of the Entomological Society of Indonesia (ICCESI [54] 9).
- [13] Bande, L.O.S., Gusnawaty HS [7] Mariadi, and Nuriadi. 2019. EFFECTIVENESS OF BOTANICAL INSECTICIDE FORMULATIONS AGAINST C [7] COA POD BORER *Conopomorpha cramerella* (Snell.). *J. HPT Tropika* Vol. 19, No. 1, 2019: 1 – 7.
- [14] Illah, I.N., Ramadhan, A., and Dhafi, A. 2017. The Ability to Kill of Kamandrah Seed Extract (*Croton tiglium* L) Against Ladybugs Chocolate Pea Pod Sucking (*Riptortus linearis*) And Its Implementations as a Medium of Learning. *e-JIP BIOL* Vol.5 (1): 48-57, Juni 2017 48 ISSN 2338-1795.
- [15] Bello Saliou, Babalakoun Owonola Adonis, Coulibaly K. Amadou, Zoundji [61] on Jeanne. 2020. Efficacy of aqueous plant extracts for control of bedbugs on cowpea (*Vigna unguiculata* (L.) Walpers) cultivars in northwestern Benin. *Agricultural Science Research Journal Volume (10) Issue (1): 15 – 30 January – 2020*.
- [16] Haryanta, D., Susilo, A., and Kusuma, W.H. 2020. Effect of mango's mistletoe (*Dendrophthoe entandra* (L.) miq) leaf extract on the biology of *Spodoptera litura* F. *Eco. Env. & Cons.* 26 (2) : 24 [33] pp. (471-479).
- [17] Suharsono & Prayogo, P. 2014. Integration Of Botanical Pesticide And Entomopathogenic Fungi To Control The Brown Stink Bug *Riptortus Linearis* F. (Hemiptera: Alydidae) In Soybean. *J. HPT Tropika*. ISSN 1411- [18] 5 Vol. 14, No. 1: 41 – 50 , Maret 2014.
- [18] Kunbhar, S., Rajput, L.B., Gilal, A.A., Channa, G.A., and Sahito, J.G.M. 2018. Impact of botanical pesticides against sucking insect pests and their insect predators in brinjal crop. *Journal of Entomology and Zoology Studies* [51] 8; 6(2): 83-87.
- [19] Doug [22] J., Toepfer, S., Bateman, M., and Jenner, W.H. 2019. Efficacy of homemade botanical insecticides based on traditional knowledge. A review. *Agronomy for Sustainable Development* (2019) 39:37 <https://doi.org/10.1007/s13593-019-0583-1>
- [20] Rahman, M.S., Faisal, A., Hasan, C.M., Ahsan, M., and Masud, M. 2017. Chemical and Biological Investigations of *Cerbera odollam* Gaertn. *J. Pharm. Sci.* 16(2): 179-186, 2017 (December).
- [21] Somsroi, P., and Chaibong, S. 2016. Effect of Suicide Tree Crude Extract (*Cerbera odollam* Gaertn.) on Common Cutworm (*Spodoptera litura* Fabricius). *JABHAT AGRIC.* 15 (1) : 16-21 (2016).
- [22] Purwani, K.I., Wijayawati, L., Nurhatika, S., Sa' Diyah, N.A., and Arifiyanto, A. 2014. Bintaro (*Cerbera odollam*) Leaf Extract As a Potential Biological Pest Control toward *Spodoptera litura* F. Mortality. *J. Appl. Environ. Biol. Sci.*, 4(4) 18-23, 2014. [15]
- [23] Juliati, Mardhiansyah, M., Arlita, T. 2016. TEST OF SOME CONCENTRATION *Cerbera manghas* LEAF EXTRACT AS A BOTANICAL PESTICIDE FOR [15] CONTROLLING *Plusia* sp. OF THE *Samanea saman*. *Jom Faperta UR* Vol 3 No 1 Februari 2016.
- [24] Aziz, A., Maig [13] T.C., Alza, Y., Ikhwan, Z., and R. Sahknan. 2019. An Effectiveness Test Analysis of Sea Mango Seeds Extract (*Cerbera manghas*) and papaya Leaves Extract (*Carica Papaya*) In Controlling The

**Creative Commons Attribution License 4.0  
(Attribution 4.0 International, CC BY 4.0)**

This article is published under the terms of the Creative Commons Attribution License 4.0

[https://creativecommons.org/licenses/by/4.0/deed.en\\_US](https://creativecommons.org/licenses/by/4.0/deed.en_US)

- Vector of Aedes Aegypti Mosquitos. World Journal of Pharmaceutical and Life Sciences 2019, Vol. 5, Issue 4 23-33.
- [25] Deng, Y., Liao, Y., Jingjing Li, Yang, L., Zhong, H., Zhou, Q. And Qing, Z., 2014. Acaricidal Activity against *Panonychus citri* and Active Ingredient of the Mangrove Plant *Cerbera manghas*. Natural Product Communications Vol. 9 (9) 2014.
- [26] Susilo, A.; Haryanta, D., Sa'adah, TT. 2019. Response of *Riptortus linearis* towards the application of Bintaro (*Cerbera manghas*) leaf extract. Eurasia J Biosci 13, 7-2224 (2019).
- [27] Hikal, W.M., Baeshen, R.S., and Said-Al Ahl, H.A.H. 2017. Botanical insecticide as simple extractives for pest control. Cogent Biology (2017), 3: 1404274. <https://doi.org/10.1080/23312025.2017.1404274>.
- [28] Sholahud, AH; Subchan, W., and Prihatin, J. 2018. Toxicity of Granules of Bintaro Leaf Extract (*Cerbera odollam* Gaertn.) on Armyworm (*Spodoptera litura* Fab.) Bioedukasi Vol. XVI. No.1 April 2018.
- [29] Degri, M.M., Mailafiya, D.M., Wabekwa, J.W. 2013. Efficacy of aqueous leaf extracts and synthetic insecticide on pod-sucking bugs infestation of cowpea (*Vigna unguiculata* (L.) Walp) in the Guinea Savanna Region of Nigeria. Advances in Entomology 1 (2013) 10-14.
- [30] Olaitan, A.F., and Abiodun, A.T. 2011. Comparative Toxicity of Botanical and Synthetic Insecticides Against major Field Insect Pests of Cowpea (*Vigna unguiculata* (L.) Walp). J. Nat. Prod. Plant Resour., 2011, 1 (3): 86-95.



Table 1. Results of the Analysis of Phytochemical Active Ingredients of Bintaro plant organs (*Cerbera manghas*) as a Source of Biopesticides Material

No	Material	Phytochemical Types				
		polyphenols %	saponins %	steroids %	Flavonoids %	Tannins %
1	young fruit pulp (coir)	1,35	3,01	1,32	2,15	0,21
2	seeds (ripe fruit)	1,03	2,18	1,24	0,38	0,34
3	Leaf	3,04	2,63	1,05	1,42	0,48
4	Bark stem/twigs	0,53	0,26	2,11	0,14	3,08

Table 2a. The Mean Percentage of *Riptortus linearis* mortality with the treatment of concentration of the extract of leaf, stem bark, fruit pulp and seeds of Bintaro (*Cerbera manghas*)

Treatment	% Mortality at day ---							
	1	2	3	4	5	6	7	8
Control	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
K <sub>1</sub> B <sub>1</sub>	0,00	0,00 b	4,17 b	4,17 b	4,17b	4,17b	4,17b	16,67ab
K <sub>3</sub> B <sub>1</sub>	0,00	0,00 b	4,17 b	4,17 b	4,17b	4,17b	4,17b	4,17b
K <sub>5</sub> B <sub>1</sub>	0,00	0,00 b	4,17 b	4,17 b	4,17b	4,17b	4,17b	4,17b
K <sub>1</sub> B <sub>2</sub>	0,00	0,00 b	0,00 b	0,00 b	0,00b	0,00b	4,17b	4,17 b
K <sub>3</sub> B <sub>2</sub>	0,00	0,00 b	4,17 b	4,17 b	4,17b	4,17b	4,17b	4,17b
K <sub>5</sub> B <sub>2</sub>	0,00	4,17 b	4,17 b	8,33 b	16,67b	16,67b	12,50b	16,67ab
K <sub>1</sub> B <sub>3</sub>	0,00	0,00 b	0,00 b	0,00 b	0,00b	0,00b	0,00b	0,00b
K <sub>3</sub> B <sub>3</sub>	0,00	0,00 b	0,00 b	0,00 b	0,00b	0,00b	0,00b	0,00b
K <sub>5</sub> B <sub>3</sub>	4,17	4,17 b	4,17 b	4,17 b	4,17b	4,17b	8,33b	8,33 b
K <sub>1</sub> B <sub>4</sub>	0,00	0,00 b	4,17 b	4,17 b	4,17b	4,17b	4,17b	4,17b
K <sub>3</sub> B <sub>4</sub>	8,33	20,83 a	29,17 a	29,17 a	29,17a	29,17a	29,17a	33,33 a
K <sub>5</sub> B <sub>4</sub>	0,00	0,00 b	0,00 b	0,00 b	0,00b	0,00b	0,00b	0,00b
LSD	TN	5,83	12,60	13,88	15,79	15,79	16,15	19,34

Table 2b. The Mean Percentage of *Riptortus linearis* mortality with the treatment of concentration of the extract of leaf, stem bark, fruit pulp and seeds of Bintaro (*Cerbera manghas*)

Treatment	% Mortality at day ---						
	9	10	11	12	13	14	15
Control	0,00	0,00	0,00	0,00	0,00	0,00	0,00
K <sub>1</sub> B <sub>1</sub>	16,67ab	16,67ab	16,67abc	16,67abc	50,00 a	54,17 a	58,33a
K <sub>3</sub> B <sub>1</sub>	4,17 b	4,17 b	4,17 bc	4,17bc	4,17 d	4,17 cd	4,17cd
K <sub>5</sub> B <sub>1</sub>	4,17 b	4,17 b	8,33 bc	8,33bc	8,33 cd	8,33 cd	8,33cd
K <sub>1</sub> B <sub>2</sub>	4,17 b	4,17 b	4,17 bc	4,17bc	4,17 d	4,17 cd	8,33cd
K <sub>3</sub> B <sub>2</sub>	4,17 b	4,17 b	4,17 bc	4,17bc	4,17 d	4,17 cd	8,33cd
K <sub>5</sub> B <sub>2</sub>	16,67ab	16,67ab	25,00 ab	25,00ab	25,00 bc	25,00 bc	25,00bc
K <sub>1</sub> B <sub>3</sub>	0,00 b	0,00 b	0,00 c	0,00c	0,00 d	0,00 d	0,00 d
K <sub>3</sub> B <sub>3</sub>	4,17 b	4,17 b	4,17 bc	4,17bc	4,17 d	4,17 cd	4,17cd
K <sub>5</sub> B <sub>3</sub>	8,33 b	8,33 b	8,33 bc	8,33bc	8,33 cd	12,50 cd	12,50cd
K <sub>1</sub> B <sub>4</sub>	4,17 b	4,17 b	8,33 bc	16,67abc	16,67 cd	16,67bcd	16,67bcd
K <sub>3</sub> B <sub>4</sub>	33,33 a	33,33 a	33,33 ac	37,50a	37,50 ab	37,50 ab	37,50ab
K <sub>5</sub> B <sub>4</sub>	0,00 b	0,00 b	0,00 c	0,00 c	0,00 d	0,00 d	0,00 d
LSD	19,63	19,63	22,58	22,33	20,06	22,08	22,83

Table 2c. The Mean Percentage of *Riptortus linearis* mortality with the treatment of concentration of the extract of leaf, stem bark, fruit pulp and seeds of Bintaro (*Cerbera manghas*)

Treatment	% Mortality at day ---				
	16	18	20	22	24
Control	0,00	4,17	4,17	4,17	4,17
K <sub>1</sub> B <sub>1</sub>	66,67 a	66,67 a	66,67 a	70,83 a	70,83 a
K <sub>3</sub> B <sub>1</sub>	4,17 de	4,17 d	4,17 d	4,17 d	4,17 de
K <sub>5</sub> B <sub>1</sub>	8,33 cde	48,33 d	8,33 d	12,50 d	16,67 cde
K <sub>1</sub> B <sub>2</sub>	8,33 cde	8,33 d	16,67 cd	16,67 cd	20,83 cde
K <sub>3</sub> B <sub>2</sub>	8,33 cde	8,33 d	8,33 d	8,33 d	12,50 de
K <sub>5</sub> B <sub>2</sub>	25,00 bc	23,60 bcd	25,00 bcd	25,00 bcd	25,00bcde
K <sub>1</sub> B <sub>3</sub>	0,00 e	0,00 63	0,00 d	0,00 d	0,00 e
K <sub>3</sub> B <sub>3</sub>	4,17 de	4,17 d	4,17 d	4,17 d	4,17 de
K <sub>5</sub> B <sub>3</sub>	12,50 cde	12,50 cd	20,83 bcd	20,83 bcd	20,83 cde
K <sub>1</sub> B <sub>4</sub>	20,83 bcd	37,50 bc	41,67 abc	41,67 bc	41,67 bc
K <sub>3</sub> B <sub>4</sub>	37,50 b	45,83 ab	45,83 ab	45,83 b	50,00 ab
K <sub>5</sub> B <sub>4</sub>	0,00 e	0,00 d	8,33 d	20,83 bcd	29,17 bcd
LSD	20,48	25,19	27,14	26,40	26,72

Note: The average value in the same column followed by the same letter is not significantly different based on the 5% LSD test

K<sub>0</sub> : treatment of the extract concentration 0%      B<sub>1</sub> : Bintaro leaf extract  
 K<sub>1</sub>: treatment of the extract concentration 2,5%      B<sub>2</sub>: Bintaro stem bark extract  
 K<sub>3</sub>: treatment of the extract concentration 7,5%      B<sub>3</sub>: Bintaro fruit extract  
 K<sub>5</sub>: treatment of the extract concentration 12,5%      B<sub>4</sub>: Bintaro seed extract



Table 3. Length of development of the *Riptortus linearis* insect per instar with the treatment of concentration of the extract of leaf, stem bark, fruit pulp and seeds of Bintaro (*Cerbera manghas*)

Treatment	Length of development (days) per instar / stadia					
	instar 2	instar 3	instar 4	instar 5	Total preadult	adult
Control	4,67	3,67	3,33	2,00	13,33	40,67
K <sub>1</sub> B <sub>1</sub>	4,67 ab	4,67	3,33	1,00	12,33	45,67 a
K <sub>3</sub> B <sub>1</sub>	5,33 a	4,00	2,67	2,00	13,33	44,67 a
K <sub>5</sub> B <sub>1</sub>	3,33 bcd	4,00	3,33	2,33	12,33	43,67 a
K <sub>1</sub> B <sub>2</sub>	4,33 abc	3,67	3,33	1,33	13,00	44,33 a
K <sub>3</sub> B <sub>2</sub>	2,67 cd	3,33	3,33	1,67	12,00	45,33 a
K <sub>5</sub> B <sub>2</sub>	4,67 ab	4,00	2,00	2,33	14,33	42,67 a
K <sub>1</sub> B <sub>3</sub>	2,00 d	4,00	3,00	1,33	10,33	45,67 a
K <sub>3</sub> B <sub>3</sub>	1,67 d	3,33	3,67	1,33	10,33	47,67 a
K <sub>5</sub> B <sub>3</sub>	3,33 bcd	3,67	2,67	1,67	11,00	46,00 a
K <sub>1</sub> B <sub>4</sub>	4,33 abc	3,33	2,33	2,67	13,00	43,00 a
K <sub>3</sub> B <sub>4</sub>	5,67 a	3,67	2,00	2,67	14,00	35,33 b
K <sub>5</sub> B <sub>4</sub>	5,67 a	4,67	2,33	3,33	14,33	26,67 c
LSD	1,87	TN	TN	TN	TN	6,04

Note : The average value in the same column followed by the same letter is not significantly different based on the 5% LSD test

K<sub>0</sub> : treatment of the extract concentration 0%      B<sub>1</sub> : Bintaro leaf extract  
K<sub>1</sub> : treatment of the extract concentration 2,5%      B<sub>2</sub> : Bintaro stem bark extract  
K<sub>3</sub> : treatment of the extract concentration 7,5%      B<sub>3</sub> : Bintaro fruit extract  
K<sub>5</sub> : treatment of the extract concentration 12,5%      B<sub>4</sub> : Bintaro seed extract

Table 4. The total age and time of 50 percent mortality of the *Riptortus linearis* insects with the treatment of the concentration of extracts of leaf, bark, fruit and seeds of Bintaro (*Cerbera manghas*)

Treatment	Total length of life (day)	length of 50% mortality (day)
Control	55,00	49,33
K <sub>1</sub> B <sub>1</sub>	58,33 a	15,00 d
K <sub>3</sub> B <sub>1</sub>	59,00 a	46,33 ab
K <sub>5</sub> B <sub>1</sub>	57,33 a	54,33 a
K <sub>1</sub> B <sub>2</sub>	58,33 a	53,67 a
K <sub>3</sub> B <sub>2</sub>	58,33 a	42,00 ab
K <sub>5</sub> B <sub>2</sub>	58,33 a	35,00 bc
K <sub>1</sub> B <sub>3</sub>	57,00 a	53,67 a
K <sub>3</sub> B <sub>3</sub>	59,00 a	56,33 a
K <sub>5</sub> B <sub>3</sub>	58,00 a	44,67 ab
K <sub>1</sub> B <sub>4</sub>	57,33 a	33,33 bc
K <sub>3</sub> B <sub>4</sub>	47,67 b	19,00 d
K <sub>5</sub> B <sub>4</sub>	44,33 b	26,33 cd
LSD	5,93	15,02

Note : The average value in the same column followed by the same letter is not significantly different based on the 5% LSD test

K<sub>0</sub> : treatment of the extract concentration 0%      B<sub>1</sub> : Bintaro leaf extract  
K<sub>1</sub> : treatment of the extract concentration 2,5%      B<sub>2</sub> : Bintaro stem bark extract  
K<sub>3</sub> : treatment of the extract concentration 7,5%      B<sub>3</sub> : Bintaro fruit extract  
K<sub>5</sub> : treatment of the extract concentration 12,5%      B<sub>4</sub> : Bintaro seed extract

Table 5. The percentage development of *Riptortus linearis* insects from the first instar to the next instar, and the imago with the treatment of the given of various concentrations of the extracts of leaf, bark, fruit and seeds of Bintaro (*Cerbera manghas*)

Treatment	Percentage development from first instar to instar...			
	instar 3	instar 4	instar 5	imago
Control	100,00	100,00	100,00	100,00
K <sub>1</sub> B <sub>1</sub>	93,75	91,67	83,33	70,83 bc
K <sub>3</sub> B <sub>1</sub>	93,75	95,83	95,83 ab	95,83 ab
K <sub>5</sub> B <sub>1</sub>	100,00	95,83	95,83 ab	91,67 ab
K <sub>1</sub> B <sub>2</sub>	100,00	95,83	95,83 ab	91,67 ab
K <sub>3</sub> B <sub>2</sub>	93,75	95,83	95,83 ab	95,83 ab
K <sub>5</sub> B <sub>2</sub>	100,00	79,17	75,00 bc	75,00 abc
K <sub>1</sub> B <sub>3</sub>	100,00	100,00	100,00 a	100,00 a
K <sub>3</sub> B <sub>3</sub>	100,00	100,00	95,83 ab	95,83 ab
K <sub>5</sub> B <sub>3</sub>	100,00	91,67	91,67 ab	91,67 ab
K <sub>1</sub> B <sub>4</sub>	100,00	95,83	91,67 ab	83,33 abc
K <sub>3</sub> B <sub>4</sub>	93,75	70,83	62,50 c	62,50 c
K <sub>5</sub> B <sub>4</sub>	100,00	100,00	100,00 a	100,00 a
LSD	TN	TN	21,56	24,51

Note: The average value in the same column followed by the same letter is not significantly different based on the 5% LSD test

K<sub>0</sub> : treatment of the extract concentration 0%

K<sub>1</sub> : treatment of the extract concentration 2,5%

K<sub>3</sub> : treatment of the extract concentration 7,5%

K<sub>5</sub> : treatment of the extract concentration 12,5%

B<sub>1</sub> : Bintaro leaf extract

B<sub>2</sub> : Bintaro stem bark extract

B<sub>3</sub> : Bintaro fruit extract

B<sub>4</sub> : Bintaro seed extract



Figure 1 Bintaro leaves




Figure 2 Bintaro bark




Figure 3 Bintaro fruit (still young)



Figure 4 Bintaro seed



## 2. Repelence of Bintaro Plant Extract (Cerbera Manghas) Against Pod-sucking Insects (Riptortus Linearis) (Hemiptera)

### ORIGINALITY REPORT

24%

SIMILARITY INDEX

19%

INTERNET SOURCES

15%

PUBLICATIONS

7%

STUDENT PAPERS

### PRIMARY SOURCES

1

[www.naun.org](http://www.naun.org)

Internet Source

2%

2

[arpgweb.com](http://arpgweb.com)

Internet Source

1%

3

[iiste.org](http://iiste.org)

Internet Source

1%

4

[krishikosh.egranth.ac.in](http://krishikosh.egranth.ac.in)

Internet Source

1%

5

[www.ejobios.org](http://www.ejobios.org)

Internet Source

1%

6

Muneer Abbas, Faisal Hafeez, Muhammad Farooq, Amjad Ali. "Dusky Cotton Bug *Oxycarenus* spp. (Hemiptera: Lygaeidae): Hibernating Sites and Management by using Plant Extracts under Laboratory Conditions", Polish Journal of Entomology, 2015

Publication

1%

7

La Ode Santiaji Bande, Gusnawaty HS, Mariadi

Mariadi, Nuriadi Nuriadi. "EFFECTIVENESS OF BOTANICAL INSECTICIDE FORMULATIONS AGAINST COCOA POD BORER *Conopomorpha cramerella* (Snell.)", JURNAL HAMA DAN PENYAKIT TUMBUHAN TROPIKA, 2020

Publication

1 %

8

[www.wseas.org](http://www.wseas.org)

Internet Source

1 %

9

[www.agriculturejournals.cz](http://www.agriculturejournals.cz)

Internet Source

1 %

10

[www.bpk-palembang.org](http://www.bpk-palembang.org)

Internet Source

1 %

11

[doi.org](http://doi.org)

Internet Source

1 %

12

[bbp2tp.litbang.pertanian.go.id](http://bbp2tp.litbang.pertanian.go.id)

Internet Source

1 %

13

[sinta3.ristekdikti.go.id](http://sinta3.ristekdikti.go.id)

Internet Source

<1 %

14

Submitted to University of Mosul

Student Paper

<1 %

15

[media.neliti.com](http://media.neliti.com)

Internet Source

<1 %

16

[basicandappliedzoology.springeropen.com](http://basicandappliedzoology.springeropen.com)

Internet Source

<1 %

17

[pps.uny.ac.id](https://pps.uny.ac.id)

Internet Source

&lt;1 %

18

[www.frontiersin.org](https://www.frontiersin.org)

Internet Source

&lt;1 %

19

[citeseerx.ist.psu.edu](https://citeseerx.ist.psu.edu)

Internet Source

&lt;1 %

20

[www.ijlpr.com](https://www.ijlpr.com)

Internet Source

&lt;1 %

21

F.O. Alao, T.A. Adebayo. "Comparative Efficacy of *Tephrosia vogelii* and *Moringa oleifera* against Insect Pests of Watermelon (*Citrullus lanatus* Thumb)", International Letters of Chemistry, Physics and Astronomy, 2015

Publication

&lt;1 %

22

Submitted to University of Southern California

Student Paper

&lt;1 %

23

Min Zhao, Han Xiao, Dong Sun, Shunshan Duan. "Investigation of the Inhibitory Effects of Mangrove Leaves and Analysis of Their Active Components on *Phaeocystis globosa* during Different Stages of Leaf Age", International Journal of Environmental Research and Public Health, 2018

Publication

&lt;1 %

24

Titumeer Al Fahad, Md Ruhul Kuddus,

&lt;1 %



Choudhury M Hasan. "Phytochemical and Biological Studies of Bark Extract of Miliusa velutina (Dunal) Hook. f. & Thomson", Dhaka University Journal of Pharmaceutical Sciences, 2020

Publication

25

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1 %

26

[download.atlantis-press.com](http://download.atlantis-press.com)

Internet Source

<1 %

27

Submitted to UIN Raden Intan Lampung

Student Paper

<1 %

28

[www.cogentoa.com](http://www.cogentoa.com)

Internet Source

<1 %

29

[econpapers.repec.org](http://econpapers.repec.org)

Internet Source

<1 %

30

[www.jurnal.unsyiah.ac.id](http://www.jurnal.unsyiah.ac.id)

Internet Source

<1 %

31

Jacobs Mobolade Adesina. " Field assessment of insecticidal efficacy of some plant aqueous extracts in reducing cowpea pod-sucking bug Stâl (Hemiptera: Coreidae) infestation and damage ", Archives Of Phytopathology And Plant Protection, 2014

Publication

<1 %

- |    |   |      |
|----|---|------|
| 32 | Ewelina Lukaszewska, Elwira Sliwinska. "Most organs of sugar-beet ( <i>Beta vulgaris</i> L.) plants at the vegetative and reproductive stages of development are polysomatic", Sexual Plant Reproduction, 2007<br>Publication | <1 % |
| 33 | S.K. Dara. "Microbial Control of Arthropod Pests in Small Fruits and Vegetables in Temperate Climate", Elsevier BV, 2017<br>Publication   | <1 % |
| 34 | Submitted to Universitas Brawijaya<br>Student Paper   | <1 % |
| 35 | biosains.mipa.uns.ac.id<br>Internet Source  | <1 % |
| 36 | balingtang.litbang.pertanian.go.id<br>Internet Source   | <1 % |
| 37 | J. T. Shaw, M. E. Gray, K. L. Steffey, J. W. Finger. "Efficacy of Soil Insecticides in Controlling Wireworms in Field Corn, 1997", Arthropod Management Tests, 1998<br>Publication  | <1 % |
| 38 | Submitted to Universiti Malaysia Sabah<br>Student Paper   | <1 % |
| 39 | Priyantini Widiyaning, Dyah Rini Indri, Bambang Priyono, Nur Asiyah, Pramita Lulu Febri.  | <1 % |

"Antifeedant Effect of Some Medicinal Plant Extracts Against Rice Weevil", Pakistan Journal of Biological Sciences, 2020

Publication

40

E Joeniarti, S Purwanti, S Hidayati, N Nurlina. "Mixed formula of neem leaves extract and curcumin as botanical insecticides for sustainable agriculture", Journal of Physics: Conference Series, 2020

Publication

<1 %

41

Tatiana E. Marinchenko. "Greening of agricultural production is the main vector of development of the Russian agricultural sector", E3S Web of Conferences, 2020

Publication

<1 %

42

Juliana Amaka Ugwu. "Insecticidal activity of some botanical extracts against legume flower thrips and legume pod borer on cowpea *Vigna unguiculata* L. Walp", The Journal of Basic and Applied Zoology, 2020

Publication

<1 %

43

"Experimental Techniques in Host-Plant Resistance", Springer Science and Business Media LLC, 2019

Publication

<1 %

44

[file.scirp.org](https://file.scirp.org)

Internet Source

<1 %



45	<a href="http://www.entomoljournal.com">www.entomoljournal.com</a> Internet Source	<1 %
46	Hyo-Seong Yeo, Jae-Young Lim, Na-Young Ahn. "Effects of Aging on Angiogenic and Muscle Growth-Related Factors in Naturally Aged Rat Skeletal Muscles", Annals of Geriatric Medicine and Research, 2020 Publication	<1 %
47	<a href="http://ciat-library.ciat.cgiar.org">ciat-library.ciat.cgiar.org</a> Internet Source	<1 %
48	<a href="http://www.repo.uni-hannover.de">www.repo.uni-hannover.de</a> Internet Source	<1 %
49	Submitted to KYUNG HEE UNIVERSITY Student Paper	<1 %
50	<a href="http://doczz.net">doczz.net</a> Internet Source	<1 %
51	Shujie Ma, Ran Jia, Menglei Guo, Kaitao Qin, Lihui Zhang. "Insecticidal activity of essential oil from Cephalotaxus sinensis and its main components against various agricultural pests", Industrial Crops and Products, 2020 Publication	<1 %
52	<a href="http://karimzadeh.ijent.ir">karimzadeh.ijent.ir</a> Internet Source	<1 %
53	"Zählsynopse", Walter de Gruyter GmbH, 2003	

54

L O S Bande, Mariadi, HS Gusnawaty, Nuriadi, L Trisulpa, Rahmania. " Botanical pesticides effect from shells of bean's cashew nut on biological agents of sp. and sp. ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

55

E.O. Egho. "Seeds of Neem Tree (*Azadirachta indica* A. Juss). Promising Biopesticide in the Management of Cowpea Insect Pests and Grain Yield in the Early Cropping Season at Asaba and Abraka, Delta State, Nigeria", Journal of Agricultural Science, 11/28/2011

Publication

<1 %

56

Bulto Tufa, Dawd Mohammad, Kumar Singh Surendra. "Evaluation of five botanical powders against maize weevil, *Sitophilus zeamais* (Motsch.) (Coleoptera: Curculionidae) on maize (Jibat variety) under laboratory condition", Journal of Plant Breeding and Crop Science, 2017

Publication

<1 %

57

OYEDEJI, Stephen, ISICHEI, Augustine Onwuegbukiwe and OGUNFIDODO, Adekunle. "Performance of some local nigerian turfgrasses

<1 %

in sole and mixed stands", Tarla Bitkileri Bilimi Derneği, 2014.

Publication

58

Fatima Shirin. "In vitro clonal propagation of mature *Tectona grandis* through axillary bud proliferation", Journal of Forest Research, 12/01/2005

Publication

<1 %

59

D. L. Kerns, T. Tellez. "Efficacy of Experimental Insecticides to Citrus Thrips on Lemons, 1996", Arthropod Management Tests, 1997

Publication

<1 %

60

Yecheng Deng, Yongmei Liao, Jingjing Li, Linlin Yang, Hui Zhong, Qiuyan Zhou, Zhen Qing. "Acaricidal Activity against and Active Ingredient of the Mangrove Plant ", Natural Product Communications, 2014

Publication

<1 %

61

F. Kusi, J.A. Nboyine, M. Abudulai, A. Seidu, Y.R. Agyare, I. Sugri, M. Zakaria, R.K. Owusu, S.K. Nutsugah, L. Asamoah. "Cultivar and insecticide spraying time effects on cowpea insect pests and grain yield in northern Ghana", Annals of Agricultural Sciences, 2019

Publication

<1 %

62

Eka Nurfani, Muhammad A.K. Purbayanto, Takashige Aono, Kouichi Takase, Yudi Darma.

<1 %

## "Origin of fast-response photocurrent in ZnO thin film", Optical Materials, 2018

Publication

63

Chengyue Liu, Shuanglin Dong, Yangen Zhou, Kunpeng Shi, Zhe Pan, Dajiang Sun, Qinfeng Gao. "Temperature-Dependent Fatty Acid Composition Change of Phospholipid in Steelhead Trout (*Oncorhynchus mykiss*) Tissues", Journal of Ocean University of China, 2019

Publication

<1%

64

"Nanopesticides", Springer Science and Business Media LLC, 2020

Publication

<1%

65

Ruchi Sood. "Asafoetida (*Ferula asafoetida*): A high-value crop suitable for the cold desert of Himachal Pradesh, India", Journal of Applied and Natural Science, 2020

Publication

<1%

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off

## 2. Repelence of Bintaro Plant Extract (Cerbera Manghas) Against Pod-sucking Insects (Riptortus Linearis) (Hemiptera)

### GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10