

Benalu mangga Journal EEC vol 26 issue 2 2020 turnitin

by Abin Suarsa

Submission date: 17-Nov-2020 09:37PM (UTC+1100)

Submission ID: 1442626524

File name: Benalu_mangga_Journal_EEC_vol_26_issue_2_2020.pdf (898.3K)

Word count: 6133

Character count: 29475

Effect of mango's mistletoe (*Dendrophthoe pentandra* (L.) miq) leaf extract on the biology of *Spodoptera litura* F.

Dwi Haryanta, Achmadi Susilo and Wahyu Hendrawan Kusuma

^{1,2,3}Faculty of Agriculture, University of Wijaya Kusuma, Surabaya, Indonesia

(Received 10 November, 2019; accepted 11 December, 2019)

ABSTRACT

Waste of mistletoe plants resulting from the pruning of urban forest plants in Surabaya is abundant, and has the potential to be used as an organic pesticide. The study aimed to determine the effect of mango's mistletoe water extract on the biology of *Spodoptera litura* F. Experiments with concentration (weight/ volume) treatment of mistletoe leaf extracts with six levels namely concentration of 0.0% (100% water); concentration of 2.5% (25 g of mistletoe in one liter of extract); concentration of 5.0% (50 g mistletoe in one liter of extract); concentration of 7.5% (75 g of mistletoe in one liter of extract); concentration of 10.0% (100 g mistletoe in one liter of extract); and a concentration of 12.5% (125 g of mistletoe in one liter of extract). Experimental variables are larvae mortality, the development of each instar and stage, and the amount of food consumption. The results showed that *Dendrophthoe pentandra* (L.) Miq) mango mistletoe water extract significantly affected on the biology of *Spodoptera litura* F, namely (1) is toxic, with a concentration of 12.5% can kill up to 95% on the 8th day significantly differ from the concentration treatment at underneath; (2) extending the larval period (larvae), but shortening the pre-pupil and pupa period; (3) reduce appetite (decreased amount of feed consumption); (4) foiling the development of larvae to become the next instar, or the development from larvae to pupae, or from pupa to imago

Key words: Mango's Mistletoe Leaf Extract, Biology of *Spodoptera litura* F.

Introduction

Urban forest plants or green open spaces in the city of Surabaya are attacked by mistletoe. There are several types of mistletoe that attack, but most are *Dendrophthoe pentandra* (L.) from the Loranthaceae familia as is commonly found attacking mango plants (Haryanta *et al.*, 2018). Mistletoe has been known as a detrimental plant, plants that are attacked its growth can be inhibited; the branches are deciduous, dry and dead, so that it can reduce the beauty of city parks. At the time of pruning of urban forest plants there will be a lot of waste from plant leaves, including leaves from mistletoes which have

not been used optimally. During this time the leaves are transported to a compost house or recycling center to be composted, or in certain places burned

The mistletoe leaf extract has been widely used in the healing of various types of diseases. The *Viscum album* mistletoe leaves contain many ingredients, minerals and vitamins. Many of the ingredients contained are affected by the host where mistletoes live. Mistletoe in the avocado plants contains higher tannins, vitamin B1, vitamin C and calcium than the other two host plants (Ishiwu *et al.*, 2014).⁴³ The mistletoe extract can improve healing in cancer treatment (Ishiwu *et al.*, 2013; Marvibaigi *et al.*, 2014; Schad *et al.*, 2013; Patrick *et al.*, 2013), and can im-

prove clinical response and patient safety level and chemotherapy process of cancer treatment (Yang *et al.*, 2013; Pelzer and Troger *et al.*, 2016). Flavonoid compounds found in the leaves of *Dendrophthoe pentandra* L. Miq is thought to be able to dissolve calcium kidney stones in vitro (Sasmitho *et al.*, 2001). The mistletoe leaf water extract is hypoglycemic and has the potential to be used in the treatment of diabetes militus (Osadolo *et al.*, 2014). Flavonoid compounds found in the water and ethyl acetate fraction of *Dendrophthoe pentandra* L. Miq mistletoes is thought to be able to dissolve calcium kidney stones (Sasmitho *et al.*, 2001), as well as antibacterial, antioxidant and antidibetes (Moghadamtousi *et al.*, 2013; Fitrilia *et al.*, 2013). Mistletoe extract of viscum album subspace abietis contains many polyphenols and has potential as an antioxidant (Yulian and Safrijal, 2018; Pietrzak *et al.*, 2014).

The mistletoe leaf extract contains terpenoids, carbohydrates, reducing sugars, saponins, glycosides, steroids and resins (Hasanah *et al.*, 2017; Benjamin *et al.*, 2016; Sembiring *et al.*, 2016; Fitrilia *et al.*, 2013). Plant stem extracts containing polyphenols, flavonoids, alkaloids, tannins and sapanosides can be used as plant-based insecticides (Attaullah *et al.*, 2019; Tando *et al.*, 2018; Diouf *et al.*, 2016). Plant ingredients such as flavonoids, saponins, tannins, steroids, cardiaglycosides, alkaloids, anthraquinones, and terpenoids are toxic to *Drosophila melanogaster* (Kiaz *et al.*, 2018). Mistletoe plant extracts of the *Loranthus micranthus* contain antibacterial ingredients with tests on the bacteria of *Escherichia coli*; *Staphylococcus aureus*; *Pseudomonas aeruginosa*; *Proteus vulgari*; *Salmonela typhii* (Benjamin *et al.*, 2016), and effective to be developed as an antimicrobial agent (Fahmi *et al.*, 2018). Plant chemical components such as flavonoid, phenolic, steroid, tannin, and saponin can kill third-instar gray worms of *Spodoptera litura* (Aini and Tukiran, 2014).

The use of plant extracts as plant-based insecticides or biopesticides has been widely recommended to farmers. Information on biopesticides is generally still limited to plant raw materials and target pests. The technology recommended is still conventional or traditional. There are twelve types of plants recommended by agricultural extension workers in 20 countries, namely garlic (*Allium sativum*), neem (*Azadirachta indica*), chili (*Capsicum spp.*), Weeds (*Chromolaena odorata*), gliriside (*Gliricidia sepium*), *Melia azedarach*, Noni (*Moringa oleifera*), tobacco (*Nicotiana tabacum*), Ocimum free

maximum), tephrosia (*Tephrosia vogelii*), marigold tree (*Tithonia diversifolia*), and bitter leaf (*Vernonia amygdalina*) (Dougoud *et al.*, 2019). Neem seed extract concentration of 50% is the most effective, environmentally friendly, inexpensive and effective method of controlling future pests (Lawal *et al.*, 2015). Botanical insecticides continue to attract attention for small farmers and the environment worldwide, considered an alternative to synthetic insecticide. The use of secondary metabolites by extracting plants is a tradition more than 3000 years ago. Intensive research on various commercial products still cannot meet global demand for biopesticides (Pavela, 2016). The use of plant extracts and botanical insecticides are biopesticides that do not pollute the environment, and can be included in integrated pest control (IPM) and organic agriculture programs (Ghosh *et al.*, 2016).

The mistletoe plants are almost never attacked by pests, so it is hoped that host plants that are sprayed (treated) with mistletoe extracts will be protected from insect pests as the mistletoes. Need to look for biopesticide ingredients with abundant availability so that they are easily available and inexpensive. This study aims to examine the ability of the mistletoe leaf extract in controlling the insect population of *Spodoptera litura* F. The results of the study can be the information or a reference for the utilization of waste management in urban forest plants, including many abundant mistletoes in the city of Surabaya as an organic pesticide in urban farming practices.

Materials and Methods

Research on the effect of mango mistletoe leaf extract (*Dendrophthoe pentandra* (L.) Miq) on the Biology of *Spodoptera litura* F insect., conducted in the laboratory of the Faculty of Agriculture, University of Wijaya Kusuma Surabaya. The mistletoe is taken from mango plants that grow on the edge of a residential road in the South Surabaya area, while *Spodoptera* eggs are obtained from rice fields in Karangpulo, Malang Regency.

Experiment with treatment of concentration (weight/volume) of extracts of mistletoe leaves with six levels, namely concentration 0.0 (100% water); : concentration of 2.5% (25 g of mistletoe in one liter of extract); concentration of 5.0% (50 g mistletoes in one liter of extract); concentration of 7.5% (75 g of mistletoes in one liter of extract); concentration of 10.0% (100 g mistletoes in one liter of extract); and a



Fig. 1a. Mango plants are attacked by Mistletoe Fig. 1b. The mistletoe leaves are ready to be extracted

concentration of 12.5% (125 g of mistletoe in one liter of extract). The experiment was repeated four times. The experimental unit was ten *spodoptera* larvae which were kept in plastic jars of 20 cm diameter and 10 cm height as in Figure 2a. Larva feed uses sunflower leaves that have been planted 7 weeks before the experiment begins.

Preparation of the *Spodoptera litura* Test Insect. Eggs or larvae are obtained from the field of legumes and then kept in plastic jars with a diameter of 16 cm and a height of 17 cm covered with gauze. The larvae are fed with sunflower leaves or *jatropha* leaves every day until the larvae enter the precupa stage. At the bottom of the jar is given a layer of fine sand as thick as 5-8 cm as a media form. After the larvae enter the sand, the remnants of the feed are cleaned, then in the jar given the cotton which dipped in honey then hung and pieces of manila paper as a place to lay eggs. Imago arising from pupae will suck honey as an additional nutrient, mate and lay eggs on manila paper. The eggs taken

then placed in another jar fed with sunflower leaves until they hatch and are used as stocks of test insect as shown in Fig. 2b.

Preparation of mistletoe leaf extract. Leaf extraction of *Dendrophthoe falcata* L can be carried out with four solvents namely water, ethanol, chloroform and petroleum ether (Tando, 2018). In this experiment the production of mistletoe leaf extracts using water solvent. Fresh and healthy mistletoe leaves are washed then air dried. The leaf that has been clean then cut into small sizes and extracted using a homogenizer or blender for 15 minutes. The solution is then filtered with a cloth and the liquid is an extract that will be used for experiments. To be more effective into the extract liquid need to be added 2 g of soap per liter of liquid. Concentration (as a treatment) uses a weight ratio of leaves and water as a solvent, for example to get a concentration of 5%, then as much as 50 g of mistletoe leaves are extracted into 1000 cc of crude extract solution (before filtering).



Fig. 2a. Trial unit, The plastic box with sunflower leaves that is ready infested larvae



Fig. 2b. The 2nd-3th instar *Spodoptera litura* larvae as experimental material

The experiment was started by dipping the sunflower leaves (leaf size adjusted to easily enter the jar) into the mistletoe leaf extract with the appropriate concentration of treatment. The dried leaves were placed in a jar, then infested the first instar *Spodoptera* larvae with 2-day-old totaling 10 heads for each experimental unit (each jar). The jar is covered with gauze as in Figure 2a.

Experimental variables were the number of dead larvae, the amount of feed consumption, namely the area of sunflower leaves lost by larvae, instar changes in larva stages marked by skin changes, changes in larva stages into pupa stages and imago stages (adults). Observations were made every day, except the variable of length of the larva was observed once in the pre pupa stage.

Results

Mortality of *Spodoptera litura* F. larvae

Experiments using the first instar larvae with two-day-old, which are small, soft and still very sensitive. Death of a larva occurs starting on the first day after treatment and occurs in all treatments. More deaths at the beginning of the trial were due to the adjustment period of the larva and the impact of moving it from the hatchery to the trial site. Death data until the 4th day tend to increase and among the treatments still showed no significant difference. Observations starting from the 5th day of the treat-

ment concentration of 10% and 12.5% experienced a greater increase than the other treatments, and on the observation of the 8th day treatment of the concentration of 12.5% the larvae remain 3 from 40 initial numbers, meaning 92,50% of larvae are dead. Movement of larva mortality data from 1st day to 8th day for all treatments is presented in Table 1.

The process of larva death at the beginning of the experiment (up to the 4th-5th day) occurred suddenly, which in the previous, the movements were still the normal on the observation the next day was dead. The cause of death at the beginning of the trial may have been due to the impact of the feed that being replaced daily or by the effects of the treatment of poisonous mistletoe leaf extract. The process of death on observation the 6th day until the end of the trial of death begins with slow movements and larva sizes tend to be smaller. The cause of larva death may be the effect of the treatment of mistletoe leaf extracts affecting the decline in appetite and the effect of chemicals in the mistletoe leaf extract on the physiological processes in the larva's body.

Total feed's consumption of *Spodoptera litura* F larvae

Spodoptera litura larvae eat sunflower leaves in the epidermis and mesophyll leaves, leaving leaves bones. Data on the amount of larva feed consumption is obtained by calculating the area of the leaf

Table 1. The mean percentage of larva Mortality of *Spodoptera litura* by the treatment of the giving of various concentrations of mango mistletoe leaf extract

Treatment	Mortality of <i>Spodoptera litura</i> on day....							
	1	2	3	4	5	6	7	8
K ₀	17.50	32.50	52.50	52.50	52.50 bc	52.50 bc	52.50 c	67.50 bc
K ₁	27.50	30.00	37.50	42.50	42.50 c	45.00 c	45.00 c	45.00 c
K ₂	45.00	52.50	60.00	60.00	62.50 abc	62.50 bc	62.50 bc	62.50 bc
K ₃	27.50	42.50	52.50	60.00	60.00 bc	62.50 bc	62.50 bc	62.50 bc
K ₄	20.00	35.00	52.50	67.50	72.50 ab	72.50 ab	75.00 ab	77.50 ab
K ₅	27.50	35.00	67.50	77.50	85.00 a	92.50 a	92.50 a	92.50 a
LSD 5%	NS	NS	NS	NS	23.23	21.80	21.87	18.11

Information : The average value in the same column followed by the same letter is not significantly different based on the LSD test 5%

K₀ : concentration of 0.0% (100% water)

K₁ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₂ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₃ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₄ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₅ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

with a hole or leaf mesophyll layer that is lost and only the bones of the leaf. The amount of feed consumption on the first day was at most on the 0% concentration treatment, namely control treatment (not given mistletoe leaf extract). The next day's feed consumption data fluctuated and among the treatments were not significantly different. Observation of feed consumption data from the 4th day to the end of observation day that is the 16th day remained fluctuating but consistently larva at a concentration of 12.5% consumed the least amount of feed consumption compared to other treatments. Data on feed consumption up to the 22nd day for each treatment are presented in Table 2a, 2b and 2c.

Insect Development of *Spodoptera litura* F from 1st instar larva to imago

The level of larva starts at the 1st instar, but because the study begins with the 1st instar at 2 days old, the measurement started at the 2nd instar. The duration of the 2nd instar and the 3rd instar development did not differ significantly among the treatments, while the 4th instar and the 5th instar as well as the total stages of larvas differed significantly among the treatments. The treatment of the mistletoe extract concentration of 12.5% was longer than the other treatments, but the pre-pupa and pupa stages were shorter. Overall insect development from instar one to adulthood for the treatment of concentrations of

Table 2a. The amount of leaves that have been consumed by *Spodoptera litura* larva's with the treatment of the giving of various concentrations of mango mistletoe leaf extract (mm²/larvae) on observations from the 1st day to the 8th day

Treatment	Amount of feed consumption on day...							
	1	2	3	4	5	6	7	8
K ₀	4.06 a	5.93	7.66 c	13.56	17.36	18.78	19.19 b	55.29 a
K ₁	2.75 b	8.93	11.11 bc	15.35	15.13	16.65	44.75 a	37.42 ab
K ₂	2.97 b	12.17	16.05 ab	16.12	14.23	19.66	24.23 b	33.97 abc
K ₃	2.45 b	11.80	18.52 ab	14.80	16.10	15.75	20.30 b	36.19 ab
K ₄	2.95 b	11.35	11.01 bc	13.48	16.04	18.96	15.88 b	20.42 cd
K ₅	2.16 b	9.80	23.50 a	10.59	8.96	9.38	12.75 b	7.38 d
LSD 5%	0.89	NS	8.91	NS	NS	NS	20.0	25.12

Table 2b. The amount of leaves that have been consumed by *Spodoptera litura* larva's with the treatment of the giving of various concentrations of mango mistletoe leaf extract (mm²/larvae) on observations from the 9th day to the 16th day

Treatment	Amount of feed consumption on day...							
	9	10	11	12	13	14	15	16
K ₀	38.99	55.29	38.99	28.96	31.46	113.00 a	89.88	98.75
K ₁	33.75	37.42	35.07	31.75	31.75	42.65 b	57.09	81.13
K ₂	44.08	33.97	44.08	38.35	38.35	50.13 b	61.15	90.27
K ₃	34.53	36.19	39.14	27.90	27.90	50.65 b	85.33	79.20
K ₄	25.33	29.17	34.38	40.13	40.13	29.25 b	69.63	85.54
K ₅	17.50	7.38	17.50	15.75	15.75	21.38 b	39.00	28.25
LSD 5%	NS	NS	NS	NS	NS	35.61	NS	NS

Information: The average value in the same column followed by the same letter is not significantly different based on the LSD test 5%

K₀ : concentration of 0.0% (100% water)

K₁ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₂ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₃ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₄ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₅ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

12.5% tends to be longer. Data on average development for each instar or each stage of insect *spodoptera litura* for each treatment presented in Table 3.

Spodoptera litura larvae which are treated with a concentration of mistletoe leaf extract fail to develop into the next phase. The failure also occurred in the control treatment that is 0% concentration. The results of the analysis of variance showed that the

highest treatment concentration of 10.00% and 12.5% experienced a failure, namely for the treatment of the 10% concentration that succeeded in becoming 3rd instar only 15%, while for the treatment of the concentration of 12.5% which succeeded in becoming the 3rd instar only 5%, even those that became imago did not exist (0%). Data on the success of the *Spodoptera litura* insect in its development from instar to next instar, from larva to prepupa,

Table 2c. The amount of leaves that have been consumed by *Spodoptera litura* larva's with the treatment of the giving of various concentrations of mango mistletoe leaf extract (mm²/larvae) on observations from the 17th day to the 22nd day

Treatment	Amount of feed consumption on day...					
	17	18	19	20	21	22
K ₀	94.69 a	73.50 a	44.31	37.40	16.90	4.70
K ₁	73.32 a	56.17 a	71.38	49.21	40.65	4.00
K ₂	76.60 a	47.33 a	32.23	25.31	12.56	5.13
K ₃	62.25 a	60.00 a	57.31	53.06	31.13	23.13
K ₄	80.33 a	69.00 a	58.75	36.92	31.13	27.75
K ₅	16.13 b	15.50 b	14.38	10.13	7.88	6.88
LSD	32.37	30.09	NS	NS	NS	NS

Information :The average value in the same column followed by the same letter is not significantly different based on the LSD test 5%

K₀ : concentration of 0.0% (100% water)

K₁ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₂ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₃ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₄ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₅ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

Table 3. The duration of development of *Spodoptera litura* of each instar, the period of larva stage, pre-pupa, and pupae with the treatment of the giving of various concentrations of mango'smistletoe leaf extract

Treatment	The duration of development of each instar or each stage (day)						
	2nd instar	3rd instar	4th instar	5th instar	larvae	prepupa	pupa
K ₀	3.25 b	8.50	4.00 cd	3.75 bc	22.50 b	2.50 bc	8.75 a
K ₁	4.00 a	6.50	5.50 b	4.25 bc	22.75 b	2.50 bc	8.00 a
K ₂	3.75 b	7.00	5.00 bc	3.25 bc	21.00 b	2.25 bc	9.00 a
K ₃	4.25 ab	6.75	4.50 bd	2.50 c	22.50 b	4.50 a	8.50 a
K ₄	5.00 a	6.50	3.50 d	4.75 b	23.50 b	3.75 ab	4.50 b
K ₅	3.50 b	8.00	6.50 a	9.00 a	28.00 a	1.00 c	2.00 c
LSD 5%	1.06	NS	1.21	2.04	3.0	2.17	2.34

Information: The average value in the same column followed by the same letter is not significantly different based on the LSD test 5%

K₀ : concentration of 0.0% (100% water)

K₁ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₂ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₃ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₄ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₅ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

pupa and Imago with the treatment of the giving of various concentrations of mango mistletoe leaf extract presented in Table 4.

Discussion

Death of larva *Spodoptera litura* occurs from the first day after the treatment. Starting on day 5, the greatest mortality rates were on the treatment with concentrations of 10.00% and 12.50%, although up to the 10th day the mortality rate of 100% do not exist yet. The mistletoe leaf extract at a concentration of 500 ppm and 100 ppm is toxic to the *Artemia Salina* Leach larvae. The chemicals contained in plant extracts (including mistletoes) consisting of flavonoids, steroids, saponins, glycosides, tannins, alkaloids, terpenoids and anthraquinone can cause death of house flies through the decrease in activity of the enzyme ² acetyl kholin esterase (ACE), acyl carier protein (ACP), AKP, a-Carboxyl, and b-Carboxyl (Attaullah *et al.* ² 2019). Chemical compounds in plants including flavonoids, steroids, saponins, glycosides, tannins, alkaloids, terpenoids and anthraquino can be used as natural pesticides, can reduce pest attacks but do not directly kill target pest organisms quickly (Tando, 2018). According to the results of research (Fahmi and Hamonangan, 2018) mistletoe leaf extract is effective to be developed as an antimicrobial agent. The chemical com-

ponents of flavonoids, phenolics, steroids, tannins, and saponins ca²³ill the third-instar *Spodoptera litura* gray larvae with LC50 values of 98.6339 mg/L (Aini and Tukiran, 2014). Plant stem extracts containing polyphenols, flavonoids, alkaloids, tannins and sapanosides can be used as plant-based insecticides against test insect of *Callosobruchus* ⁵²ulatus and *Sitophilus zeamais* (Diouf *et al.*, 2016). Leaf and stem extracts of mistletoes (*Viscum album*) taken from the host plants of *Azadirachta indica*, *Acacia albida* and *Psidium guajava* contain tannins, alkaloids, saponins, glycosides, flavonoids, phenols, a combination of anthraquinones and reduction sugars, can cause mortality in test animals namely albino rats (John *et al.*, 2018). Extracts of several weed plants are toxic to the *Drosophila melanogaster* fly. Ingredients contained in the extract are flavonoids, saponins, tannins, steroids, cardiacglycosides, alkaloids, anthraquinones, and terpenoids that can drivethe decrease in function of the enzymes of AChE, AcP AkP, 5-Carboxyl, and 5-Carboxyl. Compounds that are direct insecticide is phenol compound, can be used in plant pest management (Riaz *et al.*, 2018). The mistletoe leaf extract of *Dendrohtoe pentandra* (L.) Miq contains flavonoids which are poisonous to the larvae with LC50 values of 13.95 µg/mL. This shows that the ⁴⁵stletoe leaf extract is poisonous to the larva and has the potential to be developed as an insecticide in pest control

Table 4. Percentage of the success of the *Spodoptera litura* insect in the development from instar to next instar, from larva to prepupa, pupa and Imago with the treatment of the giving of various concentrations of mango'smistletoe leaf extract

Treatment	Percentage of instar or stage						
	⁵¹ 2nd instar	3th instar	4th instar	5th instar	pre pupa	pupa	imago
K ₀	37.50	30.00	30.00	30.00	30.00	30.00	15.00
K ₁	62.50	50.00 a	45.00	27.50	27.50	27.50	17.50
K ₂	37.50	32.50 a	32.50	30.00	30.00	27.50	17.50
K ₃	47.50	37.50 a	37.50	25.00	20.00	27.50	15.00
K ₄	47.50	15.00	15.00	15.00	10.00	10.00	7.50
K ₅	30.00	5.00	5.00	2.50	2.50	2.50	0.00
BNT	TN ¹	17.68	16.61	13.33	11.35	14.75	8.75

Information: The average value in the same column followed by the same letter is not significantly different based on the LSD test 5%

K₀ : concentration of 0.0% (100% water)

K₁ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₂ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₃ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₄ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

K₅ : concentration of 7.5% (75 g of mistletoe in one liter of extract)

(Hardiyanti *et al.*, 2018).

Feed consumption data from observation of the 4th day to end of observation the 16th day remained fluctuating but consistently larva at a concentration of 12.5% consumed the least amount of feed consumption compared to other treatments. In general, larva feed consumption decreases during the process of skin replacement (changing instars), and maximum feed consumption in larvae of 3th-5th instar. Pharmacological screening results showed that the ethanol extract of mango mistletoe leaves had an effect on the central nervous system by reducing the effect of retablisment and hanging on male mice, as well as providing a catalytic effect on female mice (Nurfaat and Indriyati, 2016).

The mistletoe leaf extract influences on the duration of development of the *Spodoptera litura* larva stadia. The treatment of the mistletoe extract concentration of 12.5% at the larvae stadia stage was longer than the other treatments. In accordance with opinion (Attaullah *et al.*, 2019) chemicals material contained in plant extracts namely Flavonoids, steroids, saponins, cardiac glycosides, tannins, alkaloids, terpenoids and anthraquinone can prolong the stages of larvae and pupae of house fly through the decrease in activity of the enzyme acetyl cholin esterase (ACE), acyl protein carrier (ACP), AKP, a-Carboxyl, and b-Carboxy.

Conclusion

The extract of *Dendrothoe pentandra* (L.) Miq significantly affected on the biology of *Spodoptera litura* F, which is (1) poisonous, with a concentration of 12.5% can be lethal to 95% on the 8th day significantly differ from the concentration treatment below; (2) extending the larval period (larvas), but shortening the prepupa and pupa periods; (3) reduce appetite (decreased amount of feed consumption); (4) foiling the development of larvae to become the next instar, or the development from larva to pupae, or from pupa to imago

Acknowledgments

Manuscript material is part of the institution's internal research scheme. We would like to thank the Rector and the Chair of the Institute of Research and Community Services at the University of Wijaya Kusuma Surabaya who have provided funds for conducting research and have given opportunities

to participate in international journal preparation training.

References

- Aini, N. and Tukiran. 2014. Fitokimia Dan Uji Bioinsektisida Isolat Dari Ekstrak Kulit Batang Tumbuhan Nyiri Batu Terhadap Ulat Grayak. *Journal of Chemistry*. 3 (2).
- Attaullah, Zahoor, M.K., Zahoor, M.A., Mubarik, M.S., Rizvi, H., Majeed, H.N., Zulhussnain, M., Ranian, K., Sultana, K., Imran, M. and Qamer, S. 2019. Insecticidal, biological and biochemical response of *Musca domestica* (Diptera: Musci) to some indigenous weed plant extracts. *Saudi Journal of Biological Sciences*. <https://doi.org/10.1016/j.sjbs.2019.05.009>
- Benja, E. Ezema, Eze, F.U. and Ezeofor, C.C. 2016. Phytochemical and Antibacterial Studies of Eastern Nigerian Mistletoe (*Loranthus micranthus*) Mistletoe on *Pentacletra Macrophylla* and *Parkia Biglobosa*. *International Journal of Pharm Tech Research*. 9 (5) : 360-365.
- Diouf, E.H.G., Diop, M., Sene, A., Samb, A. and Gueye, S. 2016. Comparison of the Insecticidal Activities of Three Plants against Two Devastating Insects: *Callosobruchus maculatus* and *Sitophilus zeamais*. *Open Access Library Journal*. 3 (e2966).
- Dougoud, J., Toepfer, S., Bateman, M. and Jenner, W.H. 2019. Efficacy of homemade botanical insecticides based on traditional knowledge. A review. *Agronomy for Sustainable Development*. 39: 37.
- Fahmi, A., Bulan, R. and Hamonangan, N. 2018. Uji Aktivitas Toksisitas Dan Antimikroba Flavonoid Total Daun Benalu (*Dendrophthoe pentandra* (L.) Miq) Dari Pohon Glodokan (*Polyalthia longifolia*). *Chempublish Journal*. 3(1).
- Fitrilia, T., Bintang, M. and Safithri, M. 2013. Phytochemical screening and antioxidant activity of clove mistletoe leaf extracts (*Dendrophthoe pentandra* (L.) Miq). *IOSR Journal of Pharmacy (e)-ISSN: 2250-3013*
- Ghosh, S.Kr. 2016. Integrated Field Management of Aphids (*Myzus persicae* Sulz. and *Aphis gossypii* Glov. Together) on Potato (*Solanum Tuberosum* L.) Using Bio-Pesticides. *International Journal of Science, Environment and Technology*. 4(3) : 682-689.
- Haque, Zaman, A., Hossain, M., Sarker, I. and Islam, S. 2014. Evaluation of Antidiarrhoeal and Insecticidal activities of Ethanol Extract and its Fractions of *Dendrophthoe falcata* (L.) Leaves. *IJPSR*. 5(9).
- Hardiyanti, R., Marpaung, L., Adnyana, I.K., Simanjuntak, P. 2018. Duku's Mistletoe Leaves (*Dendrothoe pentandra* (L.) Miq) Collected from North Sumatera as Botanical Insecticides. *IOP Conf. Series. Journal of Physics: Conf. Series*. 1232 (2019) 012016
- Haryanta, D. and Susilo, A. 2018. Pola Distribusi Dan Identifikasi Jenis Benalu Pada Tumbuhan Ruang

- Terbuka Hijau Kota Surabaya. *Journal of Research and Technology*. 4(2) Desember 2018
- 31 Hasanah, U., Rosdiana, D. and Syaefudin, 2017. Aktivitas Antibakteri Ekstrak Etanol Kulit Batang dan Daun Berenuk (*Crescentia cujete* L.). *Current Biochemistry*. 4 (1): 1- 14
- 13 Ishiwu, C.N., Obiegbuna, J.E. and Aniagolu, N.M. 2013. Evaluation of Chemical Properties of Mistletoe Leaves from Three Different Trees (Avocado, African Oil Bean and Kola). *NIFOJ*. 31(2): 1-7.
- 34 John Wasagwa, Buratai Bala Lawan, Biu Abubakar Abdullahi, Luka Joshua, Ngulde Ibrahim Saidu, Onyi Emmanuel Thank God. 2018. Phytochemistry and Toxicity Studies on Aqueous Extracts of Leaf and Stem of Mistletoe (*Viscum album*) in Albino Rats. *International Journal of Homeopathy & Natural Medicines*. 4(1): 1-5. doi: 10.11648/j.ijhnm.20180401.11
- 48 Lawal, F., Aliyu, R.E. and Adamu, A.K. 2015. Efficacy of Aqueous Neem Seed Extract in the Control of Green Peach Aphids (*Myzus Persicae* Sulzer) on Chili Pepper (*Capsicum Annum* L.). *Journal of Agriculture and Crops*. 1(5): 57-62
- 21 Marvibaigi, M., Supriyanto, E., Amini, N., Majid, F.A.A., and Jaganathan, S.K.. 2014. Preclinical and Clinical Effects of Mistletoe against Breast Cancer. *BioMed Research International*. Volume 2014, Article ID 785479, 15 pages.
- 20 Moghadamtousi, S.Z., Hajrezaei, M., Kadir, H.A. and Zandi, K. 2013. *Loranthus micranthus* Linn.: Biological Activities and Phytochemistry. Evidence-Based Complementary and Alternative Medicine Volume 2013, Article ID 273712, 9 pages <http://dx.doi.org/10.1155/2013/273712>
- 37 Nurfaat, D.L. and Indriyati, W. 2016. Uji Toksisitas Akut Ekstrak Etanol Benalu Mangga (*Dendrophthoe petandra*) Terhadap Mencit Swiss Webster. *IJPST*. Volume 3, Nomor 2, Juni 2016.
- 33 Osadolor, H.B. and Ojewu, D.D. 2014. Aqueous extracts of African mistletoe (*Loranthus bengwensis*) leaves exert hypoglycaemic effects in normal rabbits. *Biokemistri* 26(3): 85-87.
- 5 Patrick J. Mansky, Dawn B. Wallerstedt, Timothy S. Sannes, Jamie Stagl, Laura Lee Johnson, Marc R. Blackman, Jean L. Grem, Sandra M. Swain, and Brian P. Monahan, 2013. NCCAM/NCI Phase 1 Study of Mistletoe Extract and Gemcitabine in Patients with Advanced Solid Tumors. *Evidence Based Complementary and Alternative Medicine*. 2013, Article ID 964592, 11 pages <http://dx.doi.org/10.1155/2013/964592>
- 17 Pavela, R. 2016. History, Presence and Perspective of Using Plant Extracts as Commercial Botanical Insecticides and Farm Products for Protection against Insects – a Review. *Plant Protect. Sci.* 52(4): 229-241.
- 11 Pelzer, F. and Troger, W. 2018. Complementary Treatment with Mistletoe Extracts During Chemotherapy: Safety, Neutropenia, Fever, and Quality of Life Assessed in a Randomized Study. *The Journal of Alternative and Complementary Medicine*. 24(9) : 954-961.
- 15 Pietrzak, W., Nowak, R. and Olech, M. 2014. Effect of extraction method on phenolic content and antioxidant activity of mistletoe extracts from *Viscum album* sub sp. *abietis*. *Chemical Papers*. 68 (7): 976-982.
- 2 Riaz, B., Zahoor, M.K., Zahoor, M.A., Majeed, H.N., Javed, I., Ahmad, A., Jabeen, F., Zulhussnain, M. and Sultana, K. 2018. Toxicity, Phytochemical Composition, and Enzyme Inhibitory Activities of Some Indigenous Weed Plant Extracts *Drosophila melanogaster*. *Evidence-Based Complementary and Alternative Medicine* Volume 2018, Article ID 2325659, 11 pages <https://doi.org/10.1155/2018/2325659>.
- 9 Sasmid, Darsono, Zainul Kamal, Matrozi. 2001. Kemampuan Fraksi Ekstrak Air Dan Etil Asetat Daun Benalu Petai (*Dendrophthoe pentandra* L. Miq) Melarutkan Batu Ginjal Kalsium Invitro Yang Diuji Dengan Metode Aktivasi Neutron Cepat. *Majalah Farnasi Indonesia*. 12(4) : 186-193.
- 3 Schad, F., Atxner, J., Buchwald, D., Happe, A., Popp, S., Kröz, M. and Matthes, H. 2013. Intratumoral Mistletoe (*Viscum album* L) Therapy in Patients With Unresectable Pancreas Carcinoma: A Retrospective Analysis. *Integrative Cancer Therapies*. 13(4) : 332-340.
- 16 Sembiring, H.B., Lenny, S. and Marpaung, L. 2016. Aktivitas Antioksidan Senyawa Flavonoida dari Daun Benalu Kakao (*Dendrophthoe pentandra* (L.) Miq.) *Chimica et Natura Acta*. 4(3) : 117-122.
- 19 Tando, E. 2018. Review: Potensi Senyawa Metabolit Sekunder dalam Sirsak (*Annona Murricata*) dan Srikaya (*Annona squamosa*) sebagai Pestisida Nabati untuk Pengendalian Hama dan Penyakit pada Tanaman. *Jurnal Biotropika*. 1 Vol. 18 No. 1
- Wongkar, J.S., Max R. J. Runtuwene and Abidjulu, J. 2015. Uji Toksisitas Ekstrak Daun Benalu Langsung (*Dendrophthoe petandra* (L) Miq) dengan Metode Brine Shrimp Lethality Test (BSLT) LC50. *Jurnal Mipa Unsrat Online*. 4 (2) : 157-160.
- 28 Yang, P., Jiang, Y., Pan, Y., Ding, X., Rhea, P., Ding, J., Hawksworth, H., Felsher, D. Narla, G., Lu, Z. and Richard T. Lee. 2019. Mistletoe extract Fraxini inhibits the proliferation of liver cancer by down-regulating c-Myc expression. *Scientific Reports* | 2019 9:6428 | <https://doi.org/10.1038/s41598-019-41444-2>
- 27 Yulian, M. and Safrijal, 2018. Uji Aktivitas Antioksidan Daun Benalu Kopi (*Loranthus Ferrugineus* Roxb.) dengan Metode Dpph (1,1 - Difenil -2- Pikrilhidrazil). *Lantanida Journal*. 6(2) : 103-202.

ORIGINALITY REPORT

21%

SIMILARITY INDEX

17%

INTERNET SOURCES

14%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1

mafiadoc.com

Internet Source

1%

2

Attaullah, Muhammad Kashif Zahoor, Muhammad Asif Zahoor, Muhammad Samee Mubarik et al. "Insecticidal, biological and biochemical response of *Musca domestica* (Diptera: Muscidae) to some indigenous weed plant extracts", Saudi Journal of Biological Sciences, 2020

Publication

1%

3

parasiticplants.org

Internet Source

1%

4

repository.unair.ac.id

Internet Source

1%

5

www.brmi.online

Internet Source

1%

6

Hira Afzal, Sohail Ahmed, Rashad Rasool Khan, Muhammad Sufian, Muhammad Arshad, Muhammad Qasim. "Management of house fly,

1%

Musca domestica L. (Muscidae: Diptera),
through botanical baits", Revista Brasileira de
Entomologia, 2020

Publication

7	Submitted to Universitas Brawijaya Student Paper	1 %
8	www.envirobiotechjournals.com Internet Source	1 %
9	www.repository.uinjkt.ac.id Internet Source	1 %
10	file.scirp.org Internet Source	1 %
11	Submitted to Queen's University of Belfast Student Paper	1 %
12	link.springer.com Internet Source	1 %
13	www.scielo.org.mx Internet Source	1 %
14	dergipark.org.tr Internet Source	<1 %
15	onlinelibrary.wiley.com Internet Source	<1 %
16	www.scribd.com Internet Source	<1 %

17

Submitted to Universidad Estadual Paulista

Student Paper

<1 %

18

media.neliti.com

Internet Source

<1 %

19

ejournal.uksw.edu

Internet Source

<1 %

20

hdl.handle.net

Internet Source

<1 %

21

aip.scitation.org

Internet Source

<1 %

22

www.cabi.org

Internet Source

<1 %

23

id.portalgaruda.org

Internet Source

<1 %

24

www.premilife.com

Internet Source

<1 %

25

www.itb.ac.id

Internet Source

<1 %

26

Submitted to Sriwijaya University

Student Paper

<1 %

27

sinta.ristekbrin.go.id

Internet Source

<1 %

28

med.stanford.edu

Internet Source

<1 %

29

econpapers.repec.org

Internet Source

<1 %

30

www.wjgnet.com

Internet Source

<1 %

31

ppjp.ulm.ac.id

Internet Source

<1 %

32

T. Agber Cyprian, Shaakaa Sewuese, Linus U. Akacha. "Preliminary Phytochemical Screening and Thin Layer Chromatography Analysis of Stem Bark Extracts of African Mistletoe Parasitic on *Vitellaria paradoxa*, *Piliostigma thonningii* and *Combretum fragrans*", Asian Journal of Applied Chemistry Research, 2019

Publication

<1 %

33

B. E. Orimadegun, E. B. Bolajoko, A. A. Onyeaghala, O. O. Ademola-Aremu.

"Quantitative analyses of phytochemical and trace elements contents of daily detox, herbal tea consumed in Nigeria", Journal of Medicinal Plants Research, 2018

Publication

<1 %

34

sciencepg.org

Internet Source

<1 %

35

Jhones Luiz de Oliveira, Estefânia Vangelie Ramos Campos, Leonardo Fernandes Fraceto. "Recent Developments and Challenges for Nanoscale Formulation of Botanical Pesticides for Use in Sustainable Agriculture", Journal of Agricultural and Food Chemistry, 2018

Publication

<1 %

36

www.hindawi.com

Internet Source

<1 %

37

journal.ugm.ac.id

Internet Source

<1 %

38

pp.qau.edu.pk

Internet Source

<1 %

39

Patrick J. Mansky, Dawn B. Wallerstedt, Timothy S. Sannes, Jamie Stagl et al. "NCCAM/NCI Phase 1 Study of Mistletoe Extract and Gemcitabine in Patients with Advanced Solid Tumors", Evidence-Based Complementary and Alternative Medicine, 2013

Publication

<1 %

40

Rini Hardiyanti, Lamek Marpaung, I Ketut Adnyana, Partomuan Simanjuntak. "Duku's Mistletoe Leaves (Dendrohtoe pentandra (L.) Miq) Collected from North Sumatera as Botanical Insecticides", Journal of Physics: Conference Series, 2019

Publication

<1 %

41	"Nanopesticides", Springer Science and Business Media LLC, 2020 Publication	<1 %
42	ejournal.undip.ac.id Internet Source	<1 %
43	www.frontiersin.org Internet Source	<1 %
44	Submitted to Associatie K.U.Leuven Student Paper	<1 %
45	Shujie Ma, Ran Jia, Menglei Guo, Kaitao Qin, Lihui Zhang. "Insecticidal activity of essential oil from <i>Cephalotaxus sinensis</i> and its main components against various agricultural pests", Industrial Crops and Products, 2020 Publication	<1 %
46	Pavel Minář. "Effect of late summer treatments by strobilurines on storage diseases of apples", Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 2006 Publication	<1 %
47	Mantu Bhuyan, Jiban J. Mahanta, Pranab R. Bhattacharyya. " Biocontrol potential of tortoise beetle () (Coleoptera: Chrysomelidae) on in Assam, India ", Biocontrol Science and Technology, 2008 Publication	<1 %

48	arpgweb.com Internet Source	<1 %
49	Sameer Sharma, Sudhakar Malla. "<i>In silico</i> Characterization and Selection of Epitope-based Peptide Vaccines Against Ebola Viruses", International Journal of Homeopathy & Natural Medicines, 2020 Publication	<1 %
50	docplayer.info Internet Source	<1 %
51	revistas.censa.edu.cu Internet Source	<1 %
52	Mohsen Marvibaigi, Neda Amini, Eko Supriyanto, Fadzilah Adibah Abdul Majid et al. "Antioxidant Activity and ROS-Dependent Apoptotic Effect of Scurrula ferruginea (Jack) Danser Methanol Extract in Human Breast Cancer Cell MDA-MB-231", PLOS ONE, 2016 Publication	<1 %
53	downloads.hindawi.com Internet Source	<1 %
54	www.mdpi.com Internet Source	<1 %
55	Julien Dougoud, Stefan Toepfer, Melanie Bateman, Wade H. Jenner. "Efficacy of	<1 %

homemade botanical insecticides based on traditional knowledge. A review", Agronomy for Sustainable Development, 2019

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off