

The Role of Zinc Supplementation on the level of MDA and the number of Mycobacterium tuberculosis colonies in male tuberculosis Wistar rats

by Sukma Sahadewa

Submission date: 07-Jun-2021 03:11PM (UTC+0700)

Submission ID: 1602008722

File name: erium_tuberculosis_colonies_in_male_tuberculosis_Wistar_rats.pdf (203.32K)

Word count: 4544

Character count: 23919

can cause lipid peroxidation. The end result of lipid peroxidase is malondialdehyde (MDA) which is a marker of oxidative damage and MDA levels correlate with the progression of pulmonary lesions (granulomas) (Palasinamy et al., 2011).

Study Method:

This is a true experimental study (true experimental laboratory) which was carried out in a laboratory in vivo by using a post-test control group design with research subjects as male Wistar (*Rattus norvegicus*) rats aged 2-3 months, obtained from the PUSVETNA unit (Surabaya). Dependent variables in this study are levels of MDA and the number of *Mycobacterium tuberculosis* and control variables in this study are temperature, humidity, close body, age, physical activity, and stress.

The induction of tuberculosis rats refers to previous studies (Mustika et al., 2014; Rodrigues et al., 2009; Heng et al., 2011; Kumar et al., 2014). Mice were infected with *Mycobacterium tuberculosis* through the trachea. The rat was anesthetized before being infected with *Mycobacterium tuberculosis*. Anesthesia is performed by injection with ketamine HCl 50 mg / kg weight and Xylazine 0.2 ml intramuscularly or subcutaneously. The mice were fixed supine and incised in the median line of the cervical area so that the trachea was seen. *Mycobacterium tuberculosis* dose was 10^8 / ml injected into the trachea as much as 50 μ l with a vertical position tuberculin needle, then stitched again. On the 29th day (week 4) after infection with *Mycobacterium tuberculosis*, mice in the negative control group (K1) were killed to prove the existence of tuberculosis infection in rat lung tissue. The left lung tissue was taken aseptically and cultured on Middlebrook 7H10 media, while the right lung was put in 10% formalin buffer for MDA examination.

After the mice were adapted 7 days before treatment, 5 rats were set aside as a negative / healthy control group (K0), and the remaining 30 mice were infected with *Mycobacterium tuberculosis* H37Rv and waited for 29 days. Rats were divided into 6 groups randomly namely Group 1 (K0) as a negative control group / healthy group of 5 animals, group 2 (K1) as a positive control group 1 / group of TB rats as much as 5 animals, group 3 (K2) as a control group positive 2 received INH dose of 6 mg / day and rifampicin 10 mg / day for 5 heads, group 4 (P1) as a treatment group 1 received INH dose of 6 mg / day, rifampicin 10 mg / day and zinc 50 mg / kg / day for 5 tails, group 5 (P2) as treatment group 2 received INH dose 6 mg / day, rifampicin 10 mg / day and zinc 100 mg / kg weight / day as many as 5 tails and group 6 (P3) as treatment group 3 received INH 6 mg / day, rifampicin 10 mg / day and zinc 200 mg / kg / day as many as 5 tails. Groups K0 and K1 were euthanized on the 30th

day to prove that K1 had been infected with TB. The left lung tissue was taken aseptically and cultured on Middlebrook 7H10 media, while the right lung was put in 10% buffered formalin to be tested for MDA. Research for groups K1, and P1-P3 continued with treatment according to the group for 8 weeks. After 8 weeks of treatment of rats in euthanasia for their organ harvesting for MDA and the number of *Mycobacterium tuberculosis* colonies. MDA measurements were performed using pulmonary tissue by the TBARS method using the rat Malondialdehyde kit.

RESULT:

In this study there were 2 variables observed, namely MDA level and the number of *Mycobacterium tuberculosis* H37Rv. While there are 6 observation groups, each consisting of 5 male Wistar rats (*Rattus norvegicus*) aged 2-3 months. Table 1 shows the average data values and standard deviations. Of the 6 groups of samples the most prominent was the number in the positive control group (K1), namely the group of mice infected with H37Rv bacteria. On MDA levels showed the highest value compared to the other observation groups. Likewise in the variable number of *Mycobacterium tuberculosis* H37Rv, the positive control group (K1) (mice infected with H37Rv bacteria) showed the only group found to be as many as *Mycobacterium tuberculosis* H37Rv while in the other group no *Mycobacterium tuberculosis* H37Rv was found.

Table 1. Comparison of MDA levels and bacteria count among groups.

Kelompok Pengamatan	Rerata \pm SD	
	Kadar MDA	Jumlah bakteri
Kontrol (-)	166 \pm 21	0
Kontrol (+) (bakteri)(K1)	356 \pm 18	83.2 \pm 4.56
Kontrol (+) (bakteri+INH)(K2)	308 \pm 16	0
P1	272 \pm 16	0
P2	232 \pm 14	0
P3	195 \pm 13	0

After the Shapiro-Wilk test on 2 variants in 6 groups, all data have met the parametric prerequisite test, which is the data proven to be spread following the normal distribution. Furthermore, the data is ready to be further analyzed by parametric statistical tests. One way Anova test was then performed between the control groups namely positive control (K1) (mice infected with H37Rv bacteria), positive control group (K2) (mice infected with H37Rv bacteria and received INH treatments 6mg / day + Rifampicin 10mg / day), to prove the effect of the treatment of *Mycobacterium tuberculosis* H37Rv and the combination of *Mycobacterium tuberculosis* H37Rv + INH treatment at a dose of 6 mg / day, Rifampicin 10 mg / day. Significant differences were found in levels of MDA, IL-10 and IL-17 between control groups.

In the one way ANOVA test between treatment groups namely group P1 (mice infected with H37Rv bacteria and receiving INH treatment 6mg / day + Rifampicin 10mg / day + Zinc 50mg / kg weight / day), P2 group (mice infected with H37Rv bacteria and receiving H37Rv bacteria) INH treatment 6mg / day + Rifampicin 10mg / day + Zinc 100mg / kg weight / day), and group P3 (mice infected with H37Rv bacteria and receive INH treatment 6mg / day + Rifampicin 10mg / day + Zinc 200mg / kg weight / day).

Table 2. Duncan test results among groups

Kelompok pengamatan	Rerata±SD	p-value
Control(+) (bakteri)	356±18 ^a	0.000<α
Control (+) (bakteri+INH)	308±16 ^b	
P1	272±16 ^c	
P2	232±14 ^d	
P3	195±13 ^e	

35

Table 2 shows the results of the Duncan test between the positive control group and 34 three treatment groups. The apparent value of MDA levels in the positive control group (K1) is greater than the three treatment groups P1, P2, or P3. This means that the combination treatment of *Mycobacterium tuberculosis* H37Rv + INH treatment 6mg / day + Rifampicin 10mg / day + Zinc in male wistar rats is proven to reduce MDA levels when 35 compared with mice infected with H37Rv bacteria. There was also a significant difference in the mean MDA level between the positive control group (K2) (308 ± 16b nmol / mL) and the P1 group (272 ± 16c nmol / mL), with the P2 group (232 ± 14d nmol / mL), and also different from the P3 group (195 ± 13e nmol / mL). The apparent value of MDA levels in the positive control group (K2) is greater than the three treatment groups P1, P2, or P3. This means that the combined treatment of *Mycobacterium tuberculosis* H37Rv + INH treatment 6mg / day + Rifampicin 10mg / day + Zinc in male wistar rats is proven to reduce MDA levels when compared with mice infected with H37Rv bacteria and receive INH treatments 6mg / day + Rifampicin 10mg / day proven to reduce MDA levels when compared with mice infected with H37Rv bacteria and receive INH treatments 6mg / day + Rifampicin 10mg / day/day. So it was proven the first sub-hypothesis, there were differences in MDA levels between the control group compared with the tuberculosis group in male wistar rats that had been given zinc supplementation




Figure 1.

Figure 1 shows the highest MDA levels in the control group (+) (bacteria) compared to the other groups. This means that the treatment of *Mycobacterium tuberculosis* H37Rv in male wistar rats will cause high MDA levels, or in other words mice will become tuberculosis. While the average stem MDA levels were slightly lower than the average stem of the control group (+) (bacteria) was the control group (+) (bacteria + INH + Rifampicin). This means that the treatment of INH treatment dose of 6 mg / day + Rifampicin dose of 10 mg / day can suppress the increase in MDA levels in male Wistar tuberculosis rats. Also visible rods and MDA levels in group P1, group P2, and group P3 showed lower than the control group (+) (bacteria) and the control group (+) (bacteria + INH + Rifampicin). Means that there is an effect of zinc supplementation can reduce MDA levels in tuberculosis male wistar rats. The higher the dose given, the lower the MDA level in tuberculosis male wistar rats.

On the lowest mean bar of MDA level is the control group (-) (healthy mice). Whereas the closest bar of MDA level to the control group (-) which is (166 ± 21a nmol / mL) the mean bar of group P3 (195 ± 13f nmol / mL). So the zinc dose of 200mg / kg / day is considered as the most optimum dose in reducing MDA levels in tuberculosis male Wistar rats.

Table 3. Mean number of *Mycobacterium tuberculosis* among groups

Kelompok pengamatan	Rerata ± SD
Control (-)	0
Control (+) (bakteri)	83.2±4.66
Control (+) (bakteri+INH)	0
P1	0
P2	0
P3	0

Table 3 shows that in male Wistar rats with tuberculosis a number of *Mycobacterium tuberculosis* were found with an average ± standard deviation of 83.2 ± 4.66. Whereas in the positive control group K2, a number of *Mycobacterium tuberculosis* was not found. It is suspected that the administration of INH treatment dose of 6mg / day + Rifampicin 10mg / day can suppress the growth of *Mycobacterium tuberculosis* in male Wistar tuberculosis rats. Likewise in the P1 group, P2 group, and P3 group there were no *Mycobacterium tuberculosis* found. It is suspected that the administration of a combination of 6 mg / d 14 INH treatment + 10 mg / day Rifampicin + zinc dose 50 mg / kg / day, 100 mg / kg / day dose, and 200 mg / kg / day dose can suppress *Mycobacterium tuberculosis* growth in male wistar rats tuberculosis. So the dose of zinc supplement 32 on that can reduce the number of bacteria is a dose of 50 mg / kg / day, a dose of 100 mg / kg / day, and a dose of 200 mg / kg / day. In other words, all these doses are able to suppress the growth of *Mycobacterium tuberculosis* in male tuberculosis wistar rats. So the twelfth sub-

hypothesis is proven, that is suspected to be found the optimum dose of zinc supplementation able to reduce the number of *Mycobacterium tuberculosis* in male Wistar tuberculosis rats.

DISCUSSION:

In this study, Zinc was given a dose of Zinc at a dose of 50 mg / kg / day, a dose of 100 mg / kg / day, and a dose of 200 mg / kg / day proved to be able to reduce levels of MDA and the number of *M. tuberculosis* bacteria in male wistar rats infected with *Mycobacterium tuberculosis* H37Rv + INH treatment 6mg / day + Rifampicin 10mg / day. Provision of Zinc in TB patients has been shown to reduce levels of oxidative stress in TB patients in several studies. In another study conducted by Suparno et al, a comparison of MDA and Zinc levels was performed in multi drug resistant tuberculosis and sensitive tuberculosis patients. In this study, there were no significant differences in MDA levels in the MDR-TB group and the TB-sensitive group. In this study, MDA levels were probably influenced by high zinc levels in patients with MDR-TB, where Zinc inhibits the nicotinamide oxidase of Adenine Dinucleotide Phosphate (NADPH). Zinc acts as an antioxidant that stops free radical reactions (Suparno, et al., 2018). Different results obtained in the study conducted by Visser et al in 2010, found no significant difference in the success of the treatment of TB patients. This study provides Zinc and vitamin A to TB patients, then monitors the progress of treatment for 8 weeks. However, the researchers stated that there were many confounding factors in this study, especially in the severity of patients and when uncontrolled sampling (Visser, et al., 2011).

The number of tuberculosis bacteria is one of the parameters used in research to assess the response of treatment. This study showed that in male Wistar rats with tuberculosis, a number of *Mycobacterium tuberculosis* was found with an average \pm standard deviation of 83.2 ± 4.66 . Whereas in the positive control group K2, a number of *Mycobacterium tuberculosis* was not found. It is suspected that the administration of INH treatment dose of 6mg / day + Rifampicin 10mg / day can suppress the growth of *Mycobacterium tuberculosis* in male Wistar tuberculosis rats. Likewise in the P1 group, P2 group, and P3 group there were no *Mycobacterium tuberculosis* found. It is suspected that the administration of a combination of 6 mg / day INH treatment + 10 mg / day Rifampicin + zinc dose 50 mg / kg / day, 100 mg / kg / day dose, and 200 mg / kg / day dose can suppress *Mycobacterium tuberculosis* growth in male wistar rats tuberculosis. So the dose of zinc supplementation that can reduce the number of bacteria is a dose of 50 mg / kg / day, a dose of 100 mg / kg / day, and a dose of 200 mg / kg / day. In other words, all these

doses are able to suppress the growth of *Mycobacterium tuberculosis* in male tuberculosis wistar rats. So the twelfth sub-hypothesis is proven, that is suspected to be found the optimum dose of zinc supplementation able to reduce the number of *Mycobacterium tuberculosis* in male Wistar tuberculosis rats.

These results are consistent with research conducted by Knekt et al in 2017, where experimental animals in the form of Wistar BALB / c rats given OAT showed a lower number of *Mycobacterium tuberculosis* colonies than those without. Identification of *Mycobacterium tuberculosis* subpopulation is one of the crucial parameters that shows the success of treatment and is widely used in research on the management of *Mycobacterium tuberculosis* infection (Knekt, et al., 2017). In this study, the result of a significant decrease in the number of *Mycobacterium tuberculosis* colonies in experimental animals given Zinc supplementation. It can be concluded that zinc supplementation can increase the immunity of experimental animals and decrease the number of *Mycobacterium tuberculosis* bacteria.

CONCLUSION:

In this study, there was a significant difference in MDA levels as a parameter of oxidative stress in TB infected mice in both the control group and the treatment group. Lower MDA levels in rats with treatment groups showed the ability of Zinc to reduce levels of oxidative stress in patients with tuberculosis.

The results of this study are in accordance with previous studies that show the function of Zinc as an antioxidant that can reduce oxidative stress in infections. This research also shows the function of Zinc in modulating the immune system. In this case, there is a response that shows an increase in the immune system in the body, thus helping in healing tuberculosis infections. However, in actual clinical settings, other comorbid conditions such as immunodeficiency, severity of tuberculosis infection and the age of the patient should be considered in determining the prognosis for successful therapy.

REFERENCE:

1. Abasaliport R, Moradi H, Zarei S, Asadi S, Salehzadeh A. 2015. Toxicity of nanoparticles on adult male Wistar rats Jour of Food and mical Toxicology.
2. T.B.R, Montiel J.L, Fernandez G, Valecillo A. (2015). The of Cytokines and other Factors Involved in the *Mycobacterium tuberculosis* Infection. World Journal of Immunology. Vol 5 (1). Pp 16-50.
3. Adriani M, Wirjatmadi B. 2014. Gizi dan Kesehatan Balita, Peranan Mikro Zinc pada Pertumbuhan Balita. Ed 1. Jakarta : Kencana Prenada
4. Beamer G.L, Flaherty D.K., Assogba B.D. (2008). Interleukin-10 notes *Mycobacterium tuberculosis* Diseases Progression in CBA/J mice. Journal of Immunology. 181 (8). 5545-5550.
5. Bharadwaj, S. 2016. Malnutrition : laboratory markers vs nutritional assesment. *Gastroenterology report*, 4(4), pp. 272-280.

6. Bonaventura P, Benedetti G, Albarède F, Miossec P. 2015. Zinc and its role in immunity and inflammation. *Autoimmunity Review*; 14:277-285.
7. Bosco, D.M., Mohanasundaram, M.D., Drogemuller, J.C., Lang, J.C., Zalewski, D.P., Coates, T.P. 2010. Zinc and Zinc Transporter Regulation in Pancreatic Islets and the Potential Role of Zinc in Islet Transplant. *Rev Diabet Stud*; 7: 263-274.
8. Calvacanti Y.V.N, Brelaz M.C.A, Neves J.K.A, Ferraz J.C. 2012. Role of TNF- α , IFN- γ , and IL-10 in the development of pulmonary tuberculosis. *Pulmonary Medicine*. Vol 2012. pp 1-10.
9. Chandrasekaran, P., Saravanan, N., Bethunaickan, R. and Tripathy, S., 2017. Malnutrition : molecular of immune responses in tuberculosis. *Frontiers in immunology*, 8(1316), pp. 1-8.
10. Chen YY, Chang JR, Huang WF, Hsu CH, Cheng HY, Sun JR, Kuo SC, Su JJ, Lin MS, Chen W, Dou HY.(2015). Genetic diversity of the Mycobacterium tuberculosis EastAfrican-Indian family in three tropical Asian countries. *Journal of Microbiology, Immunology and Infection*; xx:1-7.
11. Dembic Z. (2015). Chapter 4: The Role and Regulation of the Immune Responses. The Cytokines of the Immun System. *British Library*. 16:106-111.
12. Dunn J.D, Alvarez L.A, Zhang X, Soldati T. 2015. Reactive oxygen species and mitochondria: A nexus of cellular homeostasis. Review article of *Redox Biology*.
13. Duzguner V, Kaya S. (2007). Effect of Zinc on the Lipid Peroxidation and the Antioxidant Defense Systems of the Alloxan-Induced Diabetic Rabbits.
14. Efimova O, Szankasi P, Kelley TW. 2011. Ncf1 (p47phox) is essential for direct regulatory T cell mediated suppression of CD4+ effector T cells. *PLoS One* 11:e16013.
15. Fenelly, K. P. 2012. Variability of infectious aerosols produced during coughing by patients with pulmonary tuberculosis. *American journal of respiratory and critical care medicine*, 186(5), pp. 450-457.
16. Forbs B, Sahn D, Weissfield AS. 2007. Diagnostic microbiology. St Louis, Missouri. Mosby Elsevier: 843-859.
17. Hoivo, S. and Fukada, T., 2016. Roles of zinc signaling in the immune system. *Journal of Immunology Research*, 10(1155), pp. 1-22.
18. Hunter, R. L., 2011. Pathology of post primary tuberculosis of the lung : an illustrated clinical review. *Tuberculosis*, 91(6), pp. 497-509.
19. Jasenosky L, Scriba T, Hanekom W, Goldfield A. (2015). T cells and Adaptive Immunity to Mycobacterium tuberculosis in Human. *Immunology Reviews*. Vol 264(1). Pp 74-87.
20. Jin W, Dong C. (2013). Review: IL-17 Cytokines in Immunity and Inflammation. *Journal Emerging Microbes and Infection*. Vol 2013 (2). pp 1-5.
21. Jindal, S. K., 2016. Oxidative stress and antioxidant imbalance : respiratory disorders. Dalam: *Oxidative stress and antioxidant protection the science of free radical biology and disease*. Hoboken: John Wiley and Sons Inc. 46 p. 307-318.
22. Khan N, Vidayanti A, Lee N.S, Cho M.Y, Eom M, Kim H.Y. (2016). Innate Immunity Holding the Flanks until Reinforced by Adaptive Immunity against Mycobacterium tuberculosis Infection. *Frontiers in Microbiology* vol 7 pp 1-9.
23. Kitabayashi C, Fukada T, Kanamoto M, Ohashi W, Hojyo S, Atsumi T. (2010). Zinc suppresses Th17 Development via Inhibiting of STAT3 Activation. *Immunology J* vol 22 no 5, pp 375-386.
24. Kulchavya, Ekaterina. (2013). Innate and Acquired Response on Tuberculosis. Review Article. *Journal Clinical and Cellular Immunology* S13-005. Pp 1-6.
25. Lamsal H, Gaultam N, Bhatta N, Toora B.D. 2007. Evaluation of lipid peroxidation product, nitrite, and antioxidant level in newly diagnosed and two months follow up patients with pulmonary TB. *Asian Journal of Biochemistry*.
26. Lyadova I.V, Panteleev A.V. 2015. Th1 and Th17 cells in tuberculosis: Protection, Pathology, and biomarkers. Review article mediators of inflammation. Volume 2015, article ID 854507.
27. Maggini S, Wintergerst E.S, Beveridge S, Hornig D.H. 2007. Selected vitamin and trace elements support immune function by strengthening epithelial barriers and cellular and humoral immune responses. *British journal of nutrition* 98 (1)29-35.
28. Marreiro D. D. N. et al., 2017. Zinc and oxidative stress : current mechanism. *Antioxidants*, 6(2), p. 24.
29. Mustika A, Rahaju A.S, Indrawati R. 2014. Penurunan kerusakan jaringan paru terinfeksi tuberkulosis oleh ekstrak pegagan melalui peningkatan ekspresi *tissue inhibitor of matrix metalloproteinase-1*. *Jurnal Veteriner* 2014 Vol.15 No.4 pp.530-540 ref.39.
30. Neyrolles O, Mintz E, Catty P. 2015. Zinc and copper toxicity in host defense against pathogens: Mycobacterium tuberculosis as a model example of an emerging paradigm. *Metal Economy Host-Microbe Interaction*; 3:1-4.
31. Overbeck S, Rink L, Haase H. 2008. Modulating the immune response by oral zinc supplementation: a single approach for multiple diseases. *Arch Immunol Ther Exp*; 56:15-30.
32. Palanisamy GS, Kirk NM, Ackart DF. 2011. Evidence for oxidative stress and defective antioxidant response in guinea pigs with tuberculosis. *PLoS One*; 6:e26254.
33. Pfander S, Fohr K, Lutz AK, Putz S, Achberger K, Linta L, et al. 2016. Cellular zinc homeostasis contributes neuronal differentiation in human induced pluripotent stem cells. *Neural Plasticity*; 2016:3760702.
34. Powers S.K. 2011. Reactive oxygen species: Impact on skeletal muscle. *Comprehensive Physiology* 1 (2): 941-969.
35. Pratomo IP, Burhan E, Tambunan V. 2012. Malnutrisi dan Tuberkulosis. *Indon Med Assoc* vol 62, Nomor: 6.
36. Radak Z. 2013. Oxygen consumption and usage during physical exercise: The balance between oxidative stress and ROS-dependent. *Antioxidants and Redox signaling* volume 18, Number 10.
37. Redford P.S, Murray PJ, O'Garra A. 2011. The Role of IL-10 in immune Regulation during Mycobacterium tuberculosis infection. *Mucosal Immunology*; 4 (3). 261-270.
38. Schneider J.M, Fujii M.L, Lamp C.L, Lönnerdal B, Zidenberg-Cherr S. 2007. The prevalence of low serum zinc and copper levels and dietary habits associated with serum zinc and copper in 12- to 36-month-old children from low-income family at risk for iron deficiency. *Journal of African Dietetic Associations*; 107:1924-1929.
39. Schwander S, Dheda K. 2011. Human lung immunity against Mycobacterium tuberculosis. *American Journal Respiratory*. Vol 183. Pp 696-702.
40. Sia K.J, Georgiva M, Rengarajan J. (2015). Review Article: Innate Immune Defenses in Human Tuberculosis: An Overview of the Interactions between Mycobacterium tuberculosis and Immune Cells. *Journal of Immunology Research* Vol 2015 pp 1-2.
41. Stensland I, Kim J.C, Bowring B, Collins A.M, Mansfield J.P, Pluske J.R. A comparison of diets supplemented with a feed additive containing organic acids, cinnamaldehyde and a permeabilizing complex, or zinc oxide, on post-weaning diarrhoea, selected bacterial populations, blood measures and performance in weaned pigs experimentally infected with enterotoxigenic E. coli. *Animals* 5;5:1147-68.
42. Sterling T.R. 2015. Treatment of pulmonary tuberculosis in HIV-infected patients. *Uptodate Wolters Kluwer*. Pp 1-23.
43. Stokowa-Soltys K, Barbosa N.A, Kasprowicz A, Wiczorek R, Gaggelli N, Gaggelli E. 2016. Studies of viomycin, an anti tuberculosis antibiotic: copper(ii) coordination, DNA degradation and the impact on ribozyme cleavage activity. *Dalton Transaction*; 45:8645-58.
44. Su W.L, Peng W.C, Huang C.H, Yang C.y. 2010. Association of reduced tumor necrotic factor alpha, gamma interferon, and interleukin 1 β but increased IL-10 expression with improved chest radiography in patients with pulmonary tuberculosis. *Clinical and Vaccine Immunology*. 17(2).223-231.
45. Sulis, G., Roggi, A., Matteelli, A. and Raviglione, M. C., 2014. Tuberculosis : Epidemiology and Control. *Mediterranean Journal of Pathology and Infectious Diseases*.
46. Turner, R. D. and Bothamley, G. H., 2014. Cough and the transmission of tuberculosis. *The journal of infectious diseases*, 21(1), pp. 1367-1372.
47. Zhang Z W, Wang Q H, Zhang J.L, Li S, Wang X.L, Xu S.W. 2012. Effects of oxidative stress on immuno-suppression induced by selenium deficiency in chickens. *Biological Trace Element Research*; 149:352-361.

The Role of Zinc Supplementation on the level of MDA and the number of Mycobacterium tuberculosis colonies in male tuberculosis Wistar rats

ORIGINALITY REPORT

22%

SIMILARITY INDEX

18%

INTERNET SOURCES

15%

PUBLICATIONS

16%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Derby Student Paper	1 %
2	K. P. Rini Vijayan, A. V. Raghu. "Embelin: an HPTLC method for quantitative estimation in five species of genus Embelia Burm. f.", Future Journal of Pharmaceutical Sciences, 2021 Publication	1 %
3	www.labome.org Internet Source	1 %
4	Submitted to Anglia Ruskin University Student Paper	1 %
5	cdn.intechopen.com Internet Source	1 %
6	Submitted to SUNY, Binghamton Student Paper	1 %
7	Submitted to University of South Australia Student Paper	1 %

8	academic.oup.com Internet Source	1 %
9	Submitted to Erasmus University of Rotterdam Student Paper	1 %
10	Submitted to University of Wales Institute, Cardiff Student Paper	1 %
11	researchonline.lshtm.ac.uk Internet Source	1 %
12	Mursel Bicer, Mustafa Akil, Abdullah Sivrikaya, Ersan Kara, Abdulkerim Kasim Baltaci, Rasim Mogulkoc. "Effect of zinc supplementation on the distribution of various elements in the serum of diabetic rats subjected to an acute swimming exercise", Journal of Physiology and Biochemistry, 2011 Publication	1 %
13	Submitted to University of Birmingham Student Paper	1 %
14	bmccomplementmedtherapies.biomedcentral.com Internet Source	1 %
15	Submitted to Universitas Brawijaya Student Paper	<1 %
16	Submitted to University of KwaZulu-Natal Student Paper	<1 %

17

Submitted to University of Salford

Student Paper

<1 %

18

W.-L. Su, W.-C. Perng, C.-H. Huang, C.-Y. Yang, C.-P. Wu, J.-H. Chen. "Association of Reduced Tumor Necrosis Factor Alpha, Gamma Interferon, and Interleukin-1 (IL-1) but Increased IL-10 Expression with Improved Chest Radiography in Patients with Pulmonary Tuberculosis", Clinical and Vaccine Immunology, 2009

Publication

<1 %

19

indonesianjpharm.farmasi.ugm.ac.id

Internet Source

<1 %

20

Nnodim Johnkennedy, Anyadoh Sylvia Onyinyechi, NwosuNjoku Emmanuel Chukwunyere. "The antioxidant status and lipid peroxidation product of newly diagnosed and 6 weeks follow-up patients with pulmonary tuberculosis in Owerri, Imo state, Nigeria", Asian Pacific Journal of Tropical Disease, 2011

Publication

<1 %

21

d-scholarship.pitt.edu

Internet Source

<1 %

22

ejnnpn.springeropen.com

Internet Source

<1 %

23

Internet Source

<1 %

24

repository.unair.ac.id

Internet Source

<1 %

25

Submitted to University of Abertay Dundee

Student Paper

<1 %

26

www.potravinarstvo.com

Internet Source

<1 %

27

Ivan Tochitsky, Sooyeon Jo, Nicholas Andrews, Masakazu Kotoda et al. "Inhibiting cough by silencing large pore-expressing airway sensory neurons with a charged sodium channel blocker", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

28

trepo.tuni.fi

Internet Source

<1 %

29

kvk.ubka.uni-karlsruhe.de

Internet Source

<1 %

30

openaccesspub.org

Internet Source

<1 %

31

repository.ias.ac.in

Internet Source

<1 %

32

eprints.whiterose.ac.uk

Internet Source

<1 %

33	Submitted to University of Liverpool Student Paper	<1 %
34	journals.sagepub.com Internet Source	<1 %
35	www.oapen.org Internet Source	<1 %
36	Dinia Rizqi Dwijayanti, Widodo, Mansur Ibrahim, Muhaimin Rifa'i. "EMSA Eritin polyherbal can suppress NF-κB activation and decrease IL-17 cytokine in an irradiated mice model", Food and Agricultural Immunology, 2016 Publication	<1 %
37	id.scribd.com Internet Source	<1 %
38	"HIV and Tuberculosis", Springer Science and Business Media LLC, 2019 Publication	<1 %
39	Submitted to Buckinghamshire Chilterns University College Student Paper	<1 %
40	Submitted to Institute of International Studies Student Paper	<1 %
41	Submitted to The University of Manchester Student Paper	<1 %

42	res.mdpi.com Internet Source	<1 %
43	ejournal.unsrat.ac.id Internet Source	<1 %
44	Submitted to Creighton University Student Paper	<1 %
45	onlinelibrary.wiley.com Internet Source	<1 %
46	yonsei.pure.elsevier.com Internet Source	<1 %
47	www.mdpi.com Internet Source	<1 %
48	dspace.library.uvic.ca Internet Source	<1 %
49	www.frontiersin.org Internet Source	<1 %
50	Agustin Iskandar, Karomah Sriwedari, Indah Adhita Wulanda, Muhammad Rasjad Indra, Hartojo, Novi Khila Firani, Ery Olivianto. "The level of chemerin and adipocyte fatty acid binding protein in Toxoplasma gondii seropositive obese individuals", Asian Pacific Journal of Tropical Biomedicine, 2017 Publication	<1 %

51 Submitted to Endeavour College of Natural Health <1 %
Student Paper

52 Mercedes Gonzalez-Juarrero. "Local Pulmonary Host-Directed Therapies for Tuberculosis via Aerosol Delivery", Wiley, 2016 <1 %
Publication

53 dokumen.pub <1 %
Internet Source

54 "Oxidative stress and antioxidant imbalance: respiratory disorders", Oxidative Stress and Antioxidant Protection, 2016. <1 %
Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off